

Review

Tragia L. Genus: Ethnopharmacological Use, Phytochemical Composition and Biological Activity

Rodrigo Duarte-Casar ^{1,2}  and Juan Carlos Romero-Benavides ^{2,*} 

¹ Maestría en Química Aplicada, Facultad de Ciencias Exactas y Naturales, Universidad Técnica Particular de Loja, Loja 110108, Ecuador; rduarte@utpl.edu.ec

² Departamento de Química, Facultad de Ciencias Exactas y Naturales, Universidad Técnica Particular de Loja, Loja 110108, Ecuador

* Correspondence: jcromerob@utpl.edu.ec; Tel.: +593-98-770-8487

Abstract: *Tragia* L. is a genus of plants belonging to the *Euphorbiaceae* family with worldwide intertropical distribution, composed of more than 150 species. In this literature review, 26 species of the genus used as medicinal plants were found, mainly in East Africa and the Indian subcontinent, with a variety of uses among which antibacterial, anti-inflammatory, anticancer and reproductive health are most common. Research has been done on a few of the species, mostly those of the Old World, with emphasis on four of them: *Tragia involucrata* Linn., *Tragia spathulata* Benth., *Tragia benthamii* Baker and *Tragia plukenetii* Radcl.-Sm., confirming several ethnomedicinal claims. Moreover, a variety of active phytochemicals have been isolated, mainly ethers, hydrocarbons, flavonoids and sterols. There is ample field for the evaluation of the activity of *Tragia* extracts and essential oils and the identification of their active compounds, particularly of the New World species, for which there is still very little research.

Keywords: *Tragia*; ethnopharmacology; phytochemicals; *Euphorbiaceae*; biological activities



Citation: Duarte-Casar, R.; Romero-Benavides, J.C. *Tragia* L. Genus: Ethnopharmacological Use, Phytochemical Composition and Biological Activity. *Plants* **2021**, *10*, 2717. <https://doi.org/10.3390/plants10122717>

Academic Editor: Antonella Smeriglio

Received: 24 September 2021
Accepted: 7 December 2021
Published: 10 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Plants have been used as a source of medicinal substances for a long time, with a use that amply predates history and presumably even mankind [1–3], and the discovery of active species and their use has historically been characterized by a trial-and-error approach [4]. This empirical knowledge has been and is being alidated by systematic research and is used as a guideline to direct the search for better and new drugs, integrating ancestral knowledge and modern methods [5].

Among the plant families considered medicinal, *Euphorbiaceae* is well regarded. The ample geographical distribution of the family and the variety of stress conditions the plants grow in, which trigger the production of secondary metabolites [6], partially explain the abundance and variety of biologically active compounds found in the family and thus its medicinal activity [7,8].

This review endeavors to summarize the current knowledge about species of the *Tragia* genus, which belongs to the *Euphorbiaceae* family, concerning their medicinal properties, phytochemical basis, and in vitro and in vivo evidence and envisioning future research prospects.

2. Genus

The genus *Tragia* is one of the 317 genera in the *Euphorbiaceae* family. There are 161 accepted names belonging to 154 species in the *Tragia* genus, with “pantropical and warm temperate distribution” [9,10]. The etymology for the name of this genus comes from the Greek *tragos*, meaning goat. This name may stem either from the name of the German botanist Hieronymus Bock—Bock means “ram” or “he-goat” in German, or from the hairy appearance of the plant that would resemble a male goat [11].

Tragia species exhibit very ample morphological characters: they are perennial plants with herb, shrub, subshrub and twining vine growth habits, with lanceolate leaves presenting either entire or serrated margins. Plants belonging to this genus sting when touched due to the presence of leaf hairs with a needle-shaped crystal of calcium oxalate (raphide) in the terminal cells that is expelled on contact and punctures the skin, allowing irritants to enter and cause transient stinging [12,13], presumably a defense mechanism against herbivores [14]. Several common names for *Tragias*, such as noseburn (*Tragia* spp.), Indian stinging nettle (*T. involucrata*), fireman (*T. volubilis*) or stinging nettle creeper (*T. durbanensis*), are due to this stinging property. Figure 1 shows *T. involucrata* leaf hairs with raphides visible, taken in Kerala, India, and *T. ramosa* with clearly visible raphides, taken in Nevada, USA.



Figure 1. *Tragia involucrata* leaves, left. *Tragia ramosa* showing leaf and stem, covered by long, rough Scheme 3.0 license; right, Stan Shebs, GDFL license.

Species belonging to *Euphorbiaceae* in general and to *Tragia* in particular are still not fully settled [8], as new species are being discovered [15] and species are being reassigned to other genera [9,16], so the number of species in the genus is still subject to change.

3. Distribution and Localization

Species belonging to the *Tragia* genus are present in subtropical America, Eastern and Southern Africa, the Indian subcontinent and Northeastern Australia. Of the 154 species listed in the genus [17], 94 are found in Africa, 48 in America, 10 in Asia and 3 in Oceania, with some species such as *T. arabica* and *T. plukenetii* present both in Africa and Asia. The map in Figure 2 shows the intertropical distribution of *Tragia* species by country.




Figure 2. Worldwide *Tragia* species distribution, by country.

4. Methodology

Published works (articles and patents) were searched on scientific databases—Science Direct, Google Scholar and Scopus—for each species of the genus, using inverted commas for an exact match, e.g., “*Tragia acalyphoides*”. Relevant articles were selected after removing search terms unrelated to the area of interest such as corrosion, reforestation or hare diet. When abundant results were obtained, the search was refined with more specific terms, for example “*Tragia involucrata medicinal*” or “*Tragia involucrata ethnopharmacology*”. Duplicate articles were removed, and the remaining articles were reviewed with a focus on ethnopharmacological uses, phytochemical composition and biological activity, both in vitro and in vivo. When possible, the latest articles, no older than 10 years, have been cited. Preprints were not included.

The research interest in *Tragia* species in medical and health sciences has increased during the last twenty years. Figure 3 shows the number of publications that include the word *Tragia* in their text in the fields mentioned. Even though the subject is not a very popular one, a steady increase in appearances can be seen, with a marked increase between 2019 and 2020 and the first half of 2021.




Figure 3. Publications containing the word *Tragia* since the year 2000 in Medical and Health sciences and in Chemical sciences. Data source: [18].

Compared to the other genera in the *Plukenieteae* tribe, *Tragia* concentrates 67% of the research, compared to 12% for *Cnesmone*, 10% for *Acidoton*, 4% for *Sphaerostylis* and 1% each for *Megistostigma*, *Pachystylidium*, *Platygyne* and *Tragiella* [18].

5. Ethnopharmacological Usage

Of the more than 150 species of the genus, few appear in the scientific literature, and even fewer are mentioned from an ethnopharmacological perspective. Notwithstanding, *Tragia* species are a part of traditional medicinal systems of East Africa and the Indian subcontinent, such as Siddha and Ayurveda [19], with documented uses of *T. involucrata* appearing as early as the 1st century CE [20] and with only a handful of mentions of *Tragia* species in the New World pharmacopoeia, concerning mostly topical applications. There is concern over an excessive use of *Tragia* species, e.g., *Tragia bicolor*, which poses a conservation hazard [21,22].

Most of the interest in this genus has been focused on four species: *Tragia involucreta*, *Tragia spathulata*, *Tragia plukenetii* and *Tragia benthamii* [23], with the bulk of the research focused on *T. involucreta*. Nevertheless, several more species and their medicinal uses appear in literature. Table 1 summarizes the species with reported medicinal use along with their stated ethnopharmacological uses, when available. The Anatomical Therapeutic Chemical (ATC) Classification by the World Health Organization (WHO) is used to classify the uses for each species [24]. Figure 4 shows the geographical distribution of the documented uses. The ethnomedical uses of *Tragia* spp are most abundant in the Indian subcontinent and East and Southern Africa.

Table 1. *Tragia* species and their ethnopharmacological use. Species are listed in alphabetical order and validated against [25].

Species	Region	Plant Organs Used	Use	Form of Usage	ATC Category	References
<i>Tragia aliena</i> Pax and K.Hoffm.	Brazil	NS	Medicinal (not specified)	NS	V	[26]
<i>Tragia benthamii</i> Baker	Nigeria, Cameroon	Whole plant	Abortifacient	Decoction	G	[27]
		Leaves, roots	Antimalarial	NS	P	[28]
		Whole plant	Filaricidal		P	[23]
<i>Tragia bicolor</i> Miq.	India, Sri Lanka	NS	Medicinal	NS	V	[21]
<i>Tragia brevipes</i> Pax.	Rwanda, Kenya	Leaves	Anticancer	Decoction Chewing Ash	L	[29]
			Antigonorrhoeic		G	[30]
			Aphrodisiac		G	[31]
			Erectile dysfunction		G	[32]
			Obesity		G	[33]
			Uterotonic		A	[34]
		G	[35]			
			G	[36]		
<i>Tragia cinerea</i> (Pax) M.G.Gilbert and Radcl.-Sm.	Ethiopia	Leaves	Antigonorrhoeic	Powdered plant, drunk mixed with butter/honey	G	[37]
		NS	Anti-inflammatory		M	[38]
			Aphrodisiac		G	
<i>Tragia cordata</i> Michx.	America, Ethiopia	Roots	Urinary tract and external parasites	Decoction Topical (powdered root)	G D	[39]
<i>Tragia dioica</i> Sond.	South Africa	Leaves	Fatigue Tuberculosis	NS	V J	[40]
<i>Tragia doryodes</i> M.G. Gilbert	Ethiopia	Leaves	Anthrax	Decoction	J	[41]
<i>Tragia durbanensis</i> Kuntze.	South Africa	NS	Skin rashes	NS	D	[42]
<i>Tragia furialis</i> Bojer	Tanzania, Madagascar	Roots	Abscess	Cold water maceration, drunk	J	[43]
			Analgesic		N	[44]
			Antimalarial		P	[45]
			Aphrodisiac		G	
			Paralysis		N	
<i>Tragia geraniifolia</i> Klotzsch ex Müll.Arg.	Argentina	Roots	Emollient	NS	D	[46]
			Rubefacient		D	[47]
		NS	Diuretic Antirheumatic		G M	
<i>Tragia gracilis</i> Griseb.	Cuba	NS	Not specified	NS	V	[48]
<i>Tragia hildebrandtii</i> Müll.Arg.	India	NS	Not specified	NS	V	[49]
<i>Tragia hispida</i> Willd.	Sri Lanka	NS	Tooth decay	NS	A	[50]
<i>Tragia insuavis</i> Prain.	Kenya	Endophytes	Antibacterial	NS	J	[51]
<i>Tragia involucreta</i> L.	Southern Asia (India, Sri Lanka, Bangladesh)	Whole plant, Leaves, Roots	Analgesic	Decoction Juice Poultice	N	[20,52]
			Antidiabetic		A	
			Anti-inflammatory		M	[53]
			Antimicrobial		J	[23]
			Antinociceptive		N	[54]
			Antioxidant		-	[55]
			Antiparasitic		D	[56]
			Antitumor		L	[57]
			Diuretic		G	[58]
			Hepatoprotective		N	
<i>Tragia meyeriana</i> Müll.Arg.	South Africa	NS	Aphrodisiac Antineoplastic Immune booster	Decoction	G	[59]
		Leaves, Stems			L	[60]
		NS (barks, stems and corms mentioned)			L	[61]

Table 1. Cont.

Species	Region	Plant Organs Used	Use	Form of Usage	ATC Category	References
<i>Tragia mitis</i> Hochst. ex A.Rich.	Ethiopia	Root	Antidiarrheal	Crushed, mixed with water and sugar	A	[62]
<i>Tragia mixta</i> M.G.Gilbert	Djibouti	Leaves	Analgesic Stomach aches Tonsillitis	Heated Poultice	N A A	[63] [64]
<i>Tragia okanyua</i> Pax	Namibia	NS Root	Dizziness Snake bite Cardiovascular problems Sexually transmitted diseases (STD)	Powdered, drunk with water	N V B G	[65] [66]
<i>Tragia plukenetii</i> Radcl.-Sm.	East Africa, India	Leaves	Antihyperglycemic Antitumor	Decoction	A L	[23]
<i>Tragia praetervisa</i> Chakrab. & N.P.Balakr.	India, Sri Lanka	NS	Not specified	NS	V	[49]
<i>Tragia preussii</i> Pax	Central African Republic	Leaves	Rheumatism	NS	M	[67]
<i>Tragia pungens</i> (Forssk.) Müll.Arg.	Yemen	Whole plant	Allergy and skin diseases Antirheumatic Cytotoxic Anti-impotence	Paste	D M L G	[68] [69] [70]
<i>Tragia ramosa</i> Torr.	U.S.A., Mexico	Leaves	Not specified	NS	V	[71]
<i>Tragia rupestris</i> Sond.	South Africa	Whole plant	Medicine (not specified)	NS	V V	[72] [73]
<i>Tragia senegalensis</i> Müll. Arg	Benin	Leaves	Azoospermia	NS	G	[74]
<i>Tragia sonderi</i> Prain	Swaziland	Root	HIV/AIDS	Decoction Topical	L	[75]
<i>Tragia spathulata</i> Benth.	West Africa	Leaves	Antibacterial	NS	J	[23] [76]
<i>Tragia subsessilis</i> Pax	Uganda	Root	Tuberculosis	NS	J	[77]
<i>Tragia uberabana</i> Müll. Arg	Brazil	NS	Medicinal Toxic	NS	V V	[78]
<i>Tragia vogelii</i> Keay	Burkina Faso	Whole plant	Abortifacient	Decoction	G	[79]
<i>Tragia volubilis</i> L.	Mexico, Antilles, Brazil	Leaves, Stem, Root	Diuretic Medicinal STDs	Decoction	G V G	[80] [26] [46,81]
<i>Tragia yucatanensis</i> Millsp.	Belize, Guatemala, Mexico	Leaves	Burns Rheumatism	Topical	D M	[82]

NS: not specified. ATC categories are as follows. A: alimentary tract and metabolism, B: blood and blood-forming organs, C: cardiovascular system, D: dermatological, G: genitourinary system and sex hormones, H: systemic hormonal preparations, excluding sex hormones and insulins, J: anti-infective for systemic use, L: antineoplastic and immunomodulating agents, M: musculo-skeletal system, N: nervous system, P: antiparasitic products, insecticides and repellents; R: respiratory system, S: sensory organs; V: various [24], not present in the classification. STDs: sexually transmitted diseases.

According to the ATC classification, the most frequent ethnopharmacological uses of *Tragia* spp. in ethnopharmacology are: genitourinary system and sex hormones, with 19% of occurrences (15 of 77); nervous system, with 12%; and alimentary tract and metabolism, anti-infective for systemic use and antineoplastic and immunomodulating agents with 10% of occurrences each. The “various” classification presents 17% of occurrences, which include non-specified and vague uses, such as “toxic” or “medicinal”.

As for the morphological structures used per species, the most common are the leaves, 38%; followed by “not specified”, 33%; whole plant, 15%; roots, 13% and a single occurrence of endophytes (3%).




Figure 4. Ethnomedicinal uses for *Tragia* spp. The circle diameter is proportional to the uses reported for each country.

6. Biological Activity

Biological activity tests of *Tragia*, both in vitro and in vivo, are performed mostly with plant extracts and to a much lesser degree with essential oils: leaf, root or the whole plant, although ethnopharmacological uses mostly employ the plant via infusions, decoctions or ashes [23,35]. Different solvents and solvent mixtures have been used for the extracts, mainly methanol and ethanol. Due to the presence of *Tragia* in ethnomedical traditions in Africa and Asia, there is a team of research about the bioactivity of Old World *Tragia* extracts that have confirmed their activity and potency in some cases. Not all the health claims or traditional uses recorded have been validated through research. Again, the bulk of the research is centered on *T. involucrata*.

6.1. In Vitro Activity

Extracts of *T. benthamii*, *T. brevipes*, *T. involucrata*, *T. pungens* and *T. spatulatha* have been tested to ascertain their in vitro activity for a variety of uses. The in vitro research is summarized in Figure 5.




Figure 5. Summary of in vitro activity of *Tragia* species.

Cases in which the efficacy has been shown in vitro are listed in Table 2.

Table 2. In vitro activity of *Tragia* extracts. Species are in alphabetical order.

Species	Extract	Plant Organs Used	Biological Activity	Biological Model	Effect	Methodology	Reference
<i>T. benthamii</i>	Methanol	Whole plant	Antibacterial	28 strains (sensitive and MDR) of <i>Pseudomonas aeruginosa</i> , <i>Klebsiella pneumoniae</i> , <i>Enterobacter aerogenes</i> , <i>Escherichia coli</i> , <i>Providencia stuartii</i>	Effective against 11/28 strains (39.3%)	256–1024 µg/mL INT colorimetric assay	[83]
<i>T. brevipes</i>	Methanol: water 9:1	Leaf	Antibacterial	<i>Escherichia coli</i> , <i>Salmonella spp.</i> , <i>Enterobacter aerogenes</i> , <i>Bacillus cereus</i> , <i>Serratia liquefaciens</i> , <i>Proteus vulgaris</i>	Inhibition zones (mm) +2 +10 +9 +24 +5 +8	500 mg/mL extract—well diffusion assay	[84]
<i>T. brevipes</i>	Methanol:DCM 1:1	Leaf	Antiproliferative	DU145 HCC HELA	+IC ₅₀ : 30 µg/mL - -	Extract MTT	[85]
<i>T. involucreta</i>	Chloroform	Root	Antidiabetic	Fertile eggs of white leghorn chicken	+	0.5, 1 mg/egg. Streptozotocin-induced diabetes	[86]
<i>T. involucreta</i>	Ethyl acetate	Root	Antibacterial Antifungal	<i>Staphylococcus aureus</i> <i>Bacillus subtilis</i> <i>Bacillus brevis</i> <i>Staphylococcus epidermidis</i> <i>Escherichia coli</i> <i>Shigella dysenteriae</i> <i>Pseudomonas aeruginosa</i> <i>Vibrio cholera</i> <i>Trichophyton rubrum</i> <i>Malassezia furfur</i>	Inhibition zones (mm) +18 +14 +5.7 +0.6 +17 +3.7 +9.4 +4.7 +3.7 +13.5	50–250 mg/mL. Disc diffusion	[53]
<i>T. involucreta</i>	Methanol	Leaf	Antifungal	<i>Rhizopus stolonifer</i> , <i>Aspergillus niger</i> , <i>Alternaria solani</i> , <i>Mucor indicus</i> , <i>Chaetomium globosum</i> , <i>Tilletia indica</i>	Inhibition zone +16 ± 0.3 mm +15 ± 0.2 mm +15 ± 0.6 mm - - +10 ± 0.5 mm	Agar disc diffusion	[87]
<i>T. involucreta</i>	Isolated hydrocarbons and ethers	-	Antibacterial	<i>Burkholderia pseudomallei</i> (TES21), <i>Burkholderia pseudomallei</i> (KHW), <i>Klebsiella pneumoniae</i> (ATCC15380) <i>Klebsiella pneumoniae</i> <i>Pseudomonas aeruginosa</i> (ATCC27853), <i>Vibrio damsela</i> , <i>Salmonella typhi</i> (ATCC51812)	Inhibition zone mm +23 +25 - +20 - +19 +28	Agar disc diffusion	[88]

Table 2. Cont.

Species	Extract	Plant Organs Used	Biological Activity	Biological Model	Effect	Methodology	Reference
<i>T. involucrata</i>	Methanol Ethyl acetate Chloroform Petroleum ether	Leaf	Antiproliferative	K562 cell lines	- - CHCl ₃ AcOEt	MTT	[89]
<i>T. involucrata</i>	Water +NP	Leaf	Antiuroliathatic	-	+Struvite crystal growth inhibitory effect	2% extract; AgNPs (200 µg mL ⁻¹)	[90]
<i>T. involucrata</i>	Methanol	Whole plant	Radioprotective	Cultured human peripheral lymphocytes	+Pretreatment (10 µg mL ⁻¹)	⁶⁰ Co gamma irradiation Comet assay	[91]
<i>T. meyeriana</i> and other plant species	Boiling water	Whole plant	Immunomodulatory	Isolated peripheral blood mononuclear cells	+	<i>S. aureus</i> stimulation. Inflammatory cytokine secretion in THP-1 monocytes	[61]
<i>T. pungens</i>	Methanol	NS	Antibacterial Cytotoxic	<i>Staphylococcus aureus</i>	+(8–14 mm)	Disk diffusion assay, Neutral red uptake assay	[69]
				<i>Bacillus subtilis</i>	-		
				<i>Micrococcus flavus</i>	-		
				<i>Pseudomonas aeruginosa</i>	-		
				<i>Candida maltosa</i>	-		
<i>FL cells</i>	+Cytotoxicity. IC ₅₀ : 70 µg/mL						
<i>T. spatulatha</i>	Ethanol Methanol Acetone	Leaf	Antibacterial Antifungal		MIC (mg/mL)	Agar well diffusion	[76]
				<i>Staphylococcus aureus</i> ,	+21		
				<i>Proteus mirabilis</i> ,	+21		
				<i>Klebsiella pneumoniae</i> ,	+25		
				<i>Salmonella typhi</i> ,	+25		
				<i>Streptococcus pneumoniae</i> ,	+25		
				<i>Escherichia coli</i> ,	-		
				<i>Candida albicans</i> ,	-		
<i>Aspergillus flavus</i> ,	-						
<i>Fusarium solani</i>	-						

MDR: multi-drug resistant. NP: nanoparticle. DCM: dichloromethane. NS: not specified; INT: p-Iodonitrotetrazolium chloride; MTT: 3-(4-5-dimethyl-2-thiazoly)-2,5-diphenyltetrazolium bromide; MIC; minimum inhibitory concentration; AcOEt: ethyl acetate; AgNP: silver nanoparticles; + active. - not active.

In vitro biological activity tests devote the most attention to leaves (36%), with whole plant and root used to a lesser extent, with both 14%. Extraction solvents are methanol (47%), DCM (5%), Ethyl acetate (10%), water (6%), chloroform (5%), petroleum ether (5%), ethanol (5%) and acetone (5%). This solvent usage supports the assumption that most active compounds are moderately polar and are thus extracted with polar solvents.

Testing centers on antibacterial (41%) and antifungal (18%) activity of the extracts, with antiproliferative (12%) and antidiabetic, antiurolithiatic, radioprotective, immunomodulatory and cytotoxic effects (6% each) behind. This is a different profile than what was found in the ethnomedicinal claims, which centers on the genitourinary system and sex hormones. This is justified because aphrodisiacs do not have the expected properties [92].

6.2. In Vivo Activity

Besides in vitro activity testing, research has been done in animal models, mostly mice and also chicks, with at least one clinical trial performed in humans. The *Tragia* extracts evaluated in vivo, summarized in Figure 6 and Table 3, are obtained from four species: *T. benthamii*, *T. furialis*, *T. involucrata* and *T. plukenetii*.




Figure 6. Summary of in vivo activity of *Tragia* extracts.

Table 3. In vivo activity of *Tragia* extracts.

Species	Extract	Plant Organs Used	Animal Model	Activity	Results	Reference
<i>T. benthamii</i>	Ethanol	Whole plant	Swiss albino mice	Antimalarial	–Very poor activity against <i>P. berghei</i> (NK-65) at 50 mg·kg ^{−1} bw.	[27]
<i>T. benthamii</i>	Water	NS	Chick	Anti-inflammatory	+Carrageenan-induced foot edema. Maximal inhibition 84.3% at 300 mg/kg bw.	[93]
<i>T. furialis</i>	Ethanol–water	NS	White albino mice	Antimalarial	+IC ₅₀ : 639.3 mg·kg ^{−1} bw against <i>P. berghei</i> .	[43]
<i>T. involucrata</i>		Root	Wistar rats	Hepatoprotective	+100–300 mg/kg bw. Hepatoprotective against CCl ₄ induced toxicity and antioxidant activity; Attenuation of biomarker alteration (SGOT, SGPT, ALP, TP).	[57]
<i>T. involucrata</i>	Benzene: Ethyl acetate 1:1	Root	<i>Culex quinquefasciatus</i>	Larvicidal	+0.1–0.4% w/v Oviposition and phagodeterrence, larvicidal.	[94]
<i>T. involucrata</i>	Ethanol	Leaf	Albino rats (male)	Nephroprotective	+250 and 500 mg/kg bw. Decrease in serum urea and creatinine in acetaminophen-induced toxicity.	[95]
<i>T. involucrata</i>	Hexane Ethyl acetate	Aerial parts	Swiss albino mice	Antitumor	+50–150 mg/kg bw. Ehrlich's Ascites Carcinoma. DD antitumor activity and increased life span for both extracts.	[96]
<i>T. involucrata</i>	Hot water	NS	Wistar rats (male)	Diuretic	+1650, 2200 mg/kg bw. Loop diuretic action.	[56]
<i>T. involucrata</i>	Hot water—freeze dried	Whole plant	Clinical trial	Antidiabetic	240 mL decoction/day. FPG decrease from 164.4 ± 20.4 to 130.9 ± 16.2 mg/dL.	[52]
<i>T. involucrata</i>	Methanol	Leaf	Swiss albino mice	Analgesic Anxiolytic Sedative	+200, 400 mg/kg bw. Acetic acid writhing and formalin-induced paw licking; behavioral tests; pentobarbital-induced sleep time.	[97]

Table 3. Cont.

Species	Extract	Plant Organs Used	Animal Model	Activity	Results	Reference
<i>T. involucrata</i>	Methanol	Leaf	Wistar rats	Antibacterial	+100, 200 mg/kg bw. Wound healing in <i>S. aureus</i> infections.	[98]
<i>T. involucrata</i>	Methanol	Leaf	Swiss albino mice	Antiepileptic	+400, 800 mg/kg bw MES, PTZ, PTX induced convulsions DD.	[99]
<i>T. involucrata</i>	Methanol	NS	Swiss albino mice	Radioprotective	+100 mg/kg bw. DD survival increase	[100]
<i>T. involucrata</i>	Methanol	Root	Charles-Foster rats Swiss albino mice	Analgesic Anti-inflammatory	+Carrageenan paw edema, cotton pellet granulomata, acetic acid writhing.	[101]
<i>T. involucrata</i>	Methanol	Root	Wistar rats	Antibacterial	+100, 200 mg/kg bw. Wound healing in <i>S. aureus</i> infections	[102]
<i>T. involucrata</i>	Methanol	Root	Charles–Foster rats Swiss albino mice	CNS depressant	+100–300 mg/kg bw. Behavioral pattern, spontaneous motility, pentobarbitone-induced sleep, body temperature, aggressive behavior pattern and conditioned avoidance response (CAR).	[103]
<i>T. involucrata</i>	Methanol Chloroform	Whole plant	Albino rats	Anti-inflammatory	+100, 300 mg/kg bw. Both extracts. Carrageenan paw oedema.	[54]
<i>T. involucrata</i>	Methanol Ethyl acetate	Whole plant	Swiss albino mice	Analgesic	+500 mg/kg bw. Acetic acid model; tail flick model analgesic activity.	[55]
<i>T. involucrata</i>	Water	Leaf	Wistar rats Swiss mice (male)	Anti-inflammatory	+50–400 mg/kg bw in carrageenan-induced hindpaw edema and cotton pellet granuloma models.	[104]
<i>T. involucrata</i>	Water +NP	Leaf	Wistar rats (male)	Antiuroliathatic	+200 mg/kg bw. CaOx stone formation inhibition in ethylene glycol-induced urolithiasis.	[90]
<i>T. plukenetii</i>	Ethanol	Aerial parts	Wistar rats (male)	Antihyperglycemic	+At an oral dose of 150 and 300 mg/kg bw. Oral glucose tolerance test in alloxan induced diabetic rats.	[105]
<i>T. plukenetii</i>	Ethanol	Whole plant	Wistar rats Guinea pigs Rabbits	Antipyretic Diuretic Antiasthmatic Analgesic Antispasmodic	+100 mg/kg bw. +Antipyretic: Brewer's yeast-induced hyperpyrexia method. +Diuretic: in vivo Lipschitz test method. +Antiasthmatic: Isolation of guinea pig ileum preparation; histamine-induced bronchoconstriction. +Analgesic: acetic acid writhing response. +Antispasmodic: studies on isolated rabbit jejunum.	[106]
<i>T. plukenetii</i>	Ethanol	Whole plant	Swiss albino mice (male)	Antitumor	+100–300mg/kg bw. Ehrlich ascites carcinoma survivability. Antioxidant parameters increased DD.	[107]
<i>T. plukenetii</i>	Methanol Benzene Chloroform	Leaf	Swiss albino mice	Anticonvulsant	+100 mg/kg bw. Methanol extract against PTZ-induced convulsions.	[108]

NS: not specified; –: no activity; +: activity present; DD: dose-dependent, bw: body weight; MES: maximal electroshock; PTZ: pentylenetetrazol; PTX: picrotoxin; FPG: fasting plasma glucose; SGOT: serum glutamic oxaloacetic transaminase; SGPT: serum glutamic pyruvic transaminase; ALP: alkaline phosphatase.

Most of the research (73%) centers on *T. involucrata*, with *T. plukenetii* (18%), *T. benthamii* (9%) and *T. furialis* (5%) behind. In vivo assay extracts were obtained from leaves (29%), whole plant (25%), root (21%) and aerial parts (8%). Solvents used are methanol (48%), ethanol (26%) and water (13%), which shows that most active compounds are polar and are thus extracted with polar solvents.

For both in vitro and in vivo testing, the most common effect is antibacterial and antimicrobial with 22% of the reviewed studies. This is higher than the 10% reported in the ethnopharmacological uses. Effects having to do with cancer prevention and treatment—antiproliferative, antitumor, cytotoxic immunomodulatory and radioprotective—add up to 17% of the reported effects, which makes it the second most frequent use. Analgesic and anti-inflammatory activity is equally reported in 10% of the tests.

The findings reported in literature validate several medicinal use cases for *Tragia* species and dismiss some claims, e.g., *T. meyeriana* as an antineoplastic [60].

7. Phytochemical Composition

Phytochemical studies allow for the identification, separation and isolation of compounds of interest [109]. Based on phytochemical screenings published in the literature, the main secondary metabolites found in *Tragia* species extracts are alkaloids, glycosides, flavonoids, and sterols [23,110].

Some compounds found in plants belonging to the *Tragia* genus, classified according to their chemical nature, are listed in Table 4. Where applicable, the biological activity of the identified compound has been mentioned.

Identification of the compounds relies heavily on spectroscopic and spectrometric methods [109], and chromatography retention times and comparison with the literature are also used for tentative identification.

Figure 7 shows the structure of some of the compounds identified in *Tragia* extracts and oils, mentioning their biological activity in bold when reported. As expected in plant extracts, there is a variety of secondary metabolites in the form of terpenoids and flavonoids. Ethers and non-terpenoid hydrocarbons are reported as having antibacterial activity, and they are not in any of the common groups of secondary metabolites. There is more information about the activity of the extracts and essential oils than about the activity of compounds on their own. The recent discovery of anti-inflammatory peptides in *Tragia benthamii* extracts [93] opens a new area of interest in the research of *Tragia* species.

A strength of the genus is its diversity and its pantropical distribution, which makes it readily available in most tropical countries. A weakness would be that, despite the interest shown concerning *T. involucreata* and other traditionally medicinal species, there appear to be no drugs derived from plants of these species, remaining in the realm of herbal remedies and plant extracts, entailing less medicinal interest than other genera of the *Euphorbiaceae* family, notably *Euphorbia* [8]. This can be attributed to the stage of research, with most work performed in vitro or in vivo and with a single clinical trial [52]. Hopefully the current research will advance into new drugs.

Table 4. Compounds isolated/identified in *Tragia* extracts and oils and their biological effect.

No.	Compound	Identified	Isolated	Methodology Used	Species	Collection area	Plant Organ Used	Use	Effect	Reference
Acetal										
1	1,1-diethoxy-2-methylpropane	X		Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
2	16-heptadecenal	X		Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
3	Hexanal	X		Hydrodistillation GC/GC-MS	<i>T. benthamii</i>	Ibadan, Nigeria	Leaves	NS	NS	[112]
4	(E)-4-(1-hydroxypropyl)-7,8-dimethyl-9-(prop-1-en-1-yl)-[1,3] dioxolo [4,5-g]quinolin-6(5H)-one	X	X	Acidified ethanol extract GC, MS, LC	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
Esters										
5	4-oxo-4H-pyran-2,6-dicarboxylic acid bis-[6-methyl-heptyl] ester	X	X	Ethanol extract IR ¹ H, ¹³ C NMR, MS	<i>T. involu-crata</i>	Salem, India	Roots	Antidiabetic	Blood glucose reduction	[86]
6	Ethyl linoleate	X	X	Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
7	Ethyl palmitate	X	X	Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
8	Vinyl hexyl ether	X	X	Aqueous extract GC, MS	<i>T. involu-crata</i>	Tamil Nadu, India	Leaf	Antibacterial <i>Escherichia coli</i> <i>Proteus vulgaris</i> <i>Staphylococcus aureus</i>	MBC 12.25 µg/mL	[98,113]
Flavonoids										
9	3-(2,4-dimethoxyphenyl)-6,7-dimethoxy-2,3-dihydrochromen-4-one	X	X	Ethyl acetate extract FTIR, MS, ¹ H NMR	<i>T. involu-crata</i>	Odisha, India	Root	Antibacterial Fungicidal	MIC 1.25-12.5 µg/mL	[53]
10	Iridin	X	X	Ethyl acetate extract FTIR, MS, ¹ H NMR	<i>T. involu-crata</i>	Odisha, India	Root	Toxic		[53]
11	Quercetin	X	X	Ethyl acetate extract FTIR, MS, ¹ H NMR	<i>T. involu-crata</i>	Odisha, India	Root	Antioxidant		[53]
12	Rutin	X	X	Ethyl acetate extract FTIR, MS, ¹ H NMR	<i>T. involu-crata</i>	Odisha, India	Root	Antioxidant		[53]
Heterocycle										
13	2,5-dithia-3,6-diazabicyclo[2.2.1]heptane	X	X	95% aqueous ethanol extraction ¹ H, ¹³ C NMR	<i>T. benthamii</i>	Ibadan, Nigeria	Whole plant	NS		[114]

Table 4. Cont.

No.	Compound	Identified	Isolated	Methodology Used	Species	Collection area	Plant Organ Used	Use	Effect	Reference
Hydrocarbons										
14	2,6-dimethylheptane	X	X	Aqueous extract GC, MS	<i>T. involu-crata</i>	Tamil Nadu, India	Leaf	Antibacterial <i>Proteus vulgaris</i>	MBC 10 µg/mL	[98]
15	2,4-dimethylhexane	X	X	Aqueous extract GC, MS	<i>T. involu-crata</i>	Tamil Nadu, India	Leaf	Antibacterial <i>Staphylococcus aureus</i>	MBC 12.25 µg/mL	[98]
16	2-methylnonane	X	X	Aqueous extract GC, MS	<i>T. involu-crata</i>	Tamil Nadu, India	Leaf	Antibacterial <i>Escherichia coli</i> <i>Proteus vulgaris</i>	MIC 5.0 µg/mL	[98]
17	Shellsol (2-methyldecane)	X	X	Aqueous extract GC, MS	<i>T. involu-crata</i>	Tamil Nadu, India	Leaf	Antibacterial <i>Proteus vulgaris</i> <i>Staphylococcus aureus</i>	MBC 25.0 µg/mL	[98]
18	3,5-di- <i>tert</i> -butyl-4-hydroxyanisole	X	X	95% aqueous ethanol extraction ¹ H, ¹³ C NMR	<i>T. benthamii</i>	Ibadan, Nigeria	Whole plant	Antioxidant		[114]
19	5-hydroxy-1-methylpiperdin-2-one	X	X	Methanol extract IR, ¹ H, ¹³ C RMN, LC	<i>T. involu-crata</i>	Kerala, India	Leaf	Antihistamine	Muscle relaxant, bronchodilating and anti-allergic effects	[115]
Polyols										
20	Erythritol	X	X	95% aqueous ethanol extraction ¹ H, ¹³ C NMR	<i>T. benthamii</i>	Ibadan, Nigeria	Whole plant	NS	NS	[114]
21	Glycerol	X	X	95% aqueous ethanol extraction ¹ H, ¹³ C NMR	<i>T. benthamii</i>	Ibadan, Nigeria	Whole plant	NS	NS	[114]
Terpenoids										
22	10,13-dimethoxy-17-(6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1 <i>H</i> -cyclopenta[α]phenanthrene.	X	X	Ethyl acetate extract FTIR, MS, ¹ H NMR	<i>T. involu-crata</i>	Odisha, India	Root	NS	NS	[53]
23	Stigmasterol	X		Aqueous extract GC, MS	<i>T. involu-crata</i>		Leaf	NS	NS	[98]
24	Caryophyllene	X		Hydrodistillation GC/GC-MS	<i>T. benthamii</i>	Ibadan, Nigeria	Leaves	Anti inflammatory		[112]
25	Citronellal	X	X	Ethanol extract IR, ¹ H RMN, LC	<i>T. ramosa</i>	Maharashtra, India	Leaves	Antibacterial		[71]
26	Clerodane	X	X	Ethanol extract IR, ¹ H RMN, LC	<i>T. ramosa</i>	Maharashtra, India	Leaves	NS	NS	[71]
27	Geranylacetone	X		Hydrodistillation GC/GC-MS	<i>T. benthamii</i>	Ibadan, Nigeria	Leaves	NS	NS	[112]

Table 4. Cont.

No.	Compound	Identified	Isolated	Methodology Used	Species	Collection area	Plant Organ Used	Use	Effect	Reference
28	Neophytadiene (2-(4,8,12-Trimethyltridecyl)buta-1,3-diene)	X	X	Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
29	Phytol	X	X	95% aqueous ethanol extraction ¹ H, ¹³ C NMR	<i>T. benthamii</i>	Ibadan, Nigeria	Whole plant	NS	NS	[114]
30	Squalene (all <i>trans</i>)	X	X	Ethanol extract GC, MS	<i>T. plukenetii</i>	NS	Whole plant	NS	NS	[111]
31	α -terpinene	X	X	Ethanol extract IR, ¹ H RMN, LC	<i>T. ramosa</i>	Maharashtra, India	Leaves	Antiinflammatory, Antimicrobial	NS	[71]

GC: gas chromatography; MS: mass spectrometry; LC: liquid chromatography; IR: infrared spectroscopy; NMR: nuclear magnetic resonance; FTIR: Fourier transform infrared spectroscopy; Q-TOF: quadrupole time of flight mass spectrometry; TLC: thin layer chromatography; NS: not specified.




Figure 7. Compounds identified in *Tragia* extracts and oils.

8. Conclusions

Species belonging to the *Tragia* genus are present in traditional medicine in several cultures and have multiple uses, among which antibacterial, anticancer and aphrodisiac are most frequent. There is scientific evidence that supports the use of these species in medicine, both at the extract level and at the active compound level, with in vivo tests in

rats and mice, but there are no drugs derived from the species yet. The activity reported most frequently for *Tragia* extracts is antimicrobial and cancer-related, which suggests further research in those areas.

Less than 20% of the *Tragia* species are considered medicinal. This implies vast potential for screening and discovery of active compounds.

Most ethnopharmacological reports come from Asia and Africa, mainly East Africa and the Indian subcontinent. New world *Tragia* species have not been sufficiently studied and may prove to be a rich source of extracts and phytochemicals for drug research. Future directions for research include nanoparticles, the research into peptides extracted from *Tragia* species and the validation of medicines containing *Tragia* extracts against SARS-CoV-2.

Author Contributions: Conceptualization, J.C.R.-B. and R.D.-C.; investigation, R.D.-C.; resources, J.C.R.-B.; writing, R.D.-C.; review and editing, J.C.R.-B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: We are grateful to Natalia Bailón-Moscoso for her many valuable suggestions that improved this work. We are also grateful to the Universidad Técnica Particular de Loja (UTPL) for supporting this research and open access publication.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Solecki, R.S. Shanidar IV, a Neanderthal Flower Burial in Northern Iraq. *Science* **1975**, *190*, 880–881. [CrossRef]
2. Ur Rehman, F.; Kalsoom, M.; Adnan, M.; Fazeli-Nasab, B.; Naz, N.; Ilahi, H.; Ali, M.F.; Ilyas, M.A.; Yousaf, G.; Toor, M.D.; et al. Importance of Medicinal Plants in Human and Plant Pathology: A Review. *Int. J. Pharm. Biomed. Res.* **2021**, *8*, 1–11. [CrossRef]
3. Marrelli, M. Medicinal Plants. *Plants* **2021**, *10*, 1355. [CrossRef]
4. Jamshidi-Kia, F.; Lorigooini, Z.; Amini-Khoei, H. Medicinal Plants: Past History and Future Perspective. *J. Herbmmed Pharmacol.* **2018**, *7*, 1–7. [CrossRef]
5. Katiyar, C.; Kanjilal, S.; Gupta, A.; Katiyar, S. Drug Discovery from Plant Sources: An Integrated Approach. *AYU Int. Q. J. Res. Ayurveda* **2012**, *33*, 10–19. [CrossRef]
6. Li, Y.; Kong, D.; Fu, Y.; Sussman, M.R.; Wu, H. The Effect of Developmental and Environmental Factors on Secondary Metabolites in Medicinal Plants. *Plant Physiol. Biochem.* **2020**, *148*, 80–89. [CrossRef]
7. Coy Barrera, C.A.; Gómez, D.C.; Castiblanco, F.A. Importancia Medicinal Del Género Croton (Euphorbiaceae). *Rev. Cuba. Plantas Med.* **2016**, *21*, 234–247.
8. Mwine, J.T.; Damme, P.V. Why Do Euphorbiaceae Tick as Medicinal Plants? A Review of Euphorbiaceae Family and Its Medicinal Features. *J. Med. Plants Res.* **2011**, *5*, 652–662.
9. Gillespie, L.J.; Cardinal-McTeague, W.M.; Wurdack, K.J. Monadelphina (Euphorbiaceae, Plukenetieae), a New Genus of Tragiinae from the Amazon Rainforest of Venezuela and Brazil. *PhytoKeys* **2020**, *169*, 119–135. [CrossRef]
10. POWO *Tragia* Plum. Ex L. | Plants of the World Online | Kew Science. Available online: <http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:327688-2> (accessed on 25 October 2021).
11. Urtecho, R. *Tragia*—FNA. Available online: <http://dev.semanticfna.org/Tragia> (accessed on 26 April 2021).
12. Naudé, T.W.; Naidoo, V. Oxalates-containing plants. In *Veterinary Toxicology*; Elsevier: Amsterdam, The Netherlands, 2007; pp. 880–891. ISBN 978-0-12-370467-2.
13. Prasad, R.; Shivay, Y.S. Oxalic Acid/Oxalates in Plants: From Self-Defence to Phytoremediation. *Curr. Sci.* **2017**, *112*, 1665–1667. [CrossRef]
14. Ensikat, H.-J.; Wessely, H.; Engeser, M.; Weigend, M. Distribution, Ecology, Chemistry and Toxicology of Plant Stinging Hairs. *Toxins* **2021**, *13*, 141. [CrossRef]
15. Freire de Sá Cordeiro, W.P.; Athiê-Souza, S.M.; Laurênio de Melo, A.; Ferreira de Sales, M. A New Endangered Species of *Tragia* (Euphorbiaceae) from the Brazilian Atlantic Forest. *Syst. Bot.* **2020**, *45*, 839–844. [CrossRef]
16. Freire de Sá Cordeiro, W.P.; Athiê-Souza, S.M.; Buriel, M.T.; de Melo, A.L.; de Sales, M.F. Chicomendes (Euphorbiaceae, Tragiinae): A New Amazonian Genus Segregated from *Tragia*. *Plant Syst. Evol.* **2021**, *307*, 46. [CrossRef]
17. Govaerts, R. *World Checklist and Bibliography of Euphorbiaceae (and Pandaceae)*; Royal Botanic Gardens: Kew, UK, 2000; Volume 1, ISBN 978-1-900347-83-9.

18. Digital Science Dimensions [Software]. Available online: <https://app.dimensions.ai/analytics/publication/overview/timeline> (accessed on 18 June 2021).
19. Kar, A.; Choudhary, B.K.; Bandyopadhyay, N.G. Comparative Evaluation of Hypoglycaemic Activity of Some Indian Medicinal Plants in Alloxan Diabetic Rats. *J. Ethnopharmacol.* **2003**, *84*, 105–108. [[CrossRef](#)]
20. Pallie, M.S.; Perera, P.K.; Kumarasinghe, N.; Arawwawala, M.; Goonasekara, C.L. Ethnopharmacological Use and Biological Activities of *Tragia involucreta* L. *Evid. Based Complement. Altern. Med.* **2020**, *2020*, 1–17. [[CrossRef](#)] [[PubMed](#)]
21. Abdulhameed, S.; Pradeep, N.S.; Sugathan, S. (Eds.). *Bioresources and Bioprocess in Biotechnology*, 1st ed.; Springer: Singapore, 2017; ISBN 978-981-10-3571-5.
22. Jose, P.A.; Hussain, K.H.; Sreekumar, V.B. *Developing an Information System for the Rare Endangered and Threatened (RET) Plants of Southern Western Ghats*; Kerala Forest Research Institute: Peechi, India, 2014; p. 42.
23. Reddy, B.S.; Rao, N.R.; Vijeepallam, K.; Pandey, V. Phytochemical, Pharmacological and Biological Profiles of *Tragia* Species (Family: Euphorbiaceae). *Afr. J. Tradit. Complement. Altern. Med.* **2017**, *14*, 105–112. [[CrossRef](#)] [[PubMed](#)]
24. WHO. Anatomical Therapeutic Chemical (ATC) Classification. Available online: <https://www.who.int/tools/atc-ddd-toolkit/atc-classification> (accessed on 8 June 2021).
25. WFO. *Tragia* L. Available online: <http://www.worldfloraonline.org/taxon/wfo-4000038765> (accessed on 7 November 2021).
26. Trindade, M.T. Espécies úteis da família Euphorbiaceae no Brasil. *Rev. Cuba. Plantas Med.* **2015**, *19*. Available online: <http://revplantasmedicinales.sld.cu/index.php/pla/article/view/113/105> (accessed on 8 June 2021).
27. Oladosu, I.A.; Balogun, S.O.; Ademowo, G.O. Phytochemical Screening, Antimalarial and Histopathological Studies of *Allophylus Africanus* and *Tragia benthamii*. *Chin. J. Nat. Med.* **2013**, *11*, 371–376. [[CrossRef](#)] [[PubMed](#)]
28. Cho-Ngwa, F.; Monya, E.; Azantsa, B.K.; Manfo, F.P.T.; Babiaka, S.B.; Mbah, J.A.; Samje, M. Filaricidal Activities on *Onchocerca* Ochengi and Loa Loa, Toxicity and Phytochemical Screening of Extracts of *Tragia Benthami* and *Piper Umbellatum*. *BMC Complement. Altern. Med.* **2016**, *16*, 326. [[CrossRef](#)] [[PubMed](#)]
29. Ajao, A.A.; Sibiya, N.P.; Moteetee, A.N. Sexual Prowess from Nature: A Systematic Review of Medicinal Plants Used as Aphrodisiacs and Sexual Dysfunction in Sub-Saharan Africa. *S. Afr. J. Bot.* **2019**, *122*, 342–359. [[CrossRef](#)]
30. Raimi, I.O.; Kopaopa, B.G.; Mugivhisa, L.L.; Lewu, F.B.; Amoo, S.O.; Olowoyo, J.O. An Appraisal of Documented Medicinal Plants Used for the Treatment of Cancer in Africa over a Twenty-Year Period (1998–2018). *J. Herb. Med.* **2020**, *23*, 100371. [[CrossRef](#)]
31. Ochwang'i, D.O.; Kimwele, C.N.; Oduma, J.A.; Gathumbi, P.K.; Kiama, S.G.; Efferth, T. Cytotoxic Activity of Medicinal Plants of the Kakamega County (Kenya) against Drug-Sensitive and Multidrug-Resistant Cancer Cells. *J. Ethnopharmacol.* **2018**, *215*, 233–240. [[CrossRef](#)] [[PubMed](#)]
32. Chekole, G. Ethnobotanical Study of Medicinal Plants Used against Human Ailments in Gubalafto District, Northern Ethiopia. *J. Ethnobiol. Ethnomed.* **2017**, *13*, 55. [[CrossRef](#)]
33. Jeruto, P.; Too, E.; Mwamburi, L.; Omari, A. An Inventory of Medicinal Plants Used to Treat Gynaecological- Obstetric-Urino-Genital Disorders in South Nandi Sub County in Kenya. *J. Nat. Sci. Res.* **2015**, *5*, 136–152.
34. Asmerom, D.; Kalay, T.H.; Araya, T.Y.; Desta, D.M.; Wondafrash, D.Z.; Tafere, G.G. Medicinal Plants Used for the Treatment of Erectile Dysfunction in Ethiopia: A Systematic Review. *BioMed Res. Int.* **2021**, *2021*, 1–12. [[CrossRef](#)]
35. Pascaline, J.; Charles, M.; George, O.; Lukhoba, C. Ethnobotanical Survey and Propagation of Some Endangered Medicinal Plants from South Nandi District of Kenya. *J. Anim. Plant Sci.* **2012**, *8*, 28.
36. Aumeeruddy, M.Z.; Mahomoodally, M.F. Global Documentation of Traditionally Used Medicinal Plants in Cancer Management: A Systematic Review. *S. Afr. J. Bot.* **2021**, *138*, 424–494. [[CrossRef](#)]
37. Amare, F.; Getachew, G. An Ethnobotanical Study of Medicinal Plants in Chiro District, West Hararghe, Ethiopia. *Afr. J. Plant Sci.* **2019**, *13*, 309–323. [[CrossRef](#)]
38. Bekele, G.; Reddy, P.R. Ethnobotanical Study of Medicinal Plants Used to Treat Human Ailments by Guji Oromo Tribes in Abaya District, Borana, Oromia, Ethiopia. *Univers. J. Plant Sci.* **2015**, *3*, 1–8. [[CrossRef](#)]
39. Moges, A.; Moges, Y. Ethiopian Common Medicinal Plants: Their Parts and Uses in Traditional Medicine—Ecology and Quality Control. In *Plant Science—Structure, Anatomy and Physiology in Plants Cultured in Vivo and in Vitro*; Gonzalez, A., Rodriguez, M., Gören Sağlam, N., Eds.; IntechOpen: London, UK, 2020; ISBN 978-1-78984-746-8.
40. Semenya, S.S.; Maroyi, A. Ethnobotanical Survey of Plants Used by Bapedi Traditional Healers to Treat Tuberculosis and Its Opportunistic Infections in the Limpopo Province, South Africa. *S. Afr. J. Bot.* **2019**, *122*, 401–421. [[CrossRef](#)]
41. Tolossa, K.; Debela, E.; Athanasiadou, S.; Tolera, A.; Ganga, G.; Houdijk, J.G. Ethno-Medicinal Study of Plants Used for Treatment of Human and Livestock Ailments by Traditional Healers in South Omo, Southern Ethiopia. *J. Ethnobiol. Ethnomed.* **2013**, *9*, 32. [[CrossRef](#)] [[PubMed](#)]
42. Ndhlovu, P.T.; Omotayo, A.O.; Otang-Mbeng, W.; Aremu, A.O. Ethnobotanical Review of Plants Used for the Management and Treatment of Childhood Diseases and Well-Being in South Africa. *S. Afr. J. Bot.* **2021**, *137*, 197–215. [[CrossRef](#)]
43. Innocent, E.; Moshi, M.; Masimba, P.; Mbwambo, Z.; Kapingu, M.; Kamuhabwa, A. Screening of Traditionally Used Plants for in Vivo Antimalarial Activity in Mice. *Afr. J. Tradit. Complement. Altern. Med.* **2010**, *6*, 163–167. [[CrossRef](#)]
44. Moshi, M.J.; Otieno, D.F.; Mbabazi, P.K.; Weisheit, A. Ethnomedicine of the Kagera Region, North Western Tanzania. Part 2: The Medicinal Plants Used in Katoro Ward, Bukoba District. *J. Ethnobiol. Ethnomed.* **2010**, *6*, 19. [[CrossRef](#)]
45. Tida, M.M.A.; Nanjarisoa, O.; Rabearivony, J.; Ranarijaona, H.L.T.; Fenoradosoa, T.A. Ethnobotanical Survey Of Plant Species Used In Traditional Medicine In Bekaraoka Region, Northeastern Madagascar. *Int. J. Adv. Res. Publ.* **2020**, *4*, 107–114.

46. Barboza, G.E.; Cantero, J.J.; Núñez, C.; Arisa Espinar, L.; Pacciaroni, A. del V. Medicinal Plants: A General Review and a Phytochemical and Ethnopharmacological Screening of the Native Argentine Flora. *Kurtziana* **2009**, *34*, 7–365.
47. Goleniowski, M.E.; Bongiovanni, G.A.; Palacio, L.; Nuñez, C.O.; Cantero, J.J. Medicinal Plants from the “Sierra de Comechingones”, Argentina. *J. Ethnopharmacol.* **2006**, *107*, 324–341. [[CrossRef](#)]
48. Carlomagno, A.; Pardini, A.; Contino Esquijerosa, Y. Medicinal Plants in Ethnobotanical and Religious Traditions in Cuba: A First Review and Updating. 2015. Available online: https://www.researchgate.net/profile/Anna-Carlomagno/publication/276886636_Medicinal_plants_in_ethnobotanical_and_religious_traditions_in_Cuba_a_first_review_and_updating/links/555b088a08aeaaff3bfaefa7/Medicinal-plants-in-ethnobotanical-and-religious-traditions-in-Cuba-a-first-review-and-updating.pdf (accessed on 15 April 2021).
49. Shiddamallayya, N.; Rao, R.; Doddamani, S.; Venkateshwarlu, G. A Glimpse on Forest Flora and Indian System of Medicine Plants of Chitradurga District, Karnataka. *Int. J. Herb. Med.* **2016**, *4*, 25–33.
50. Hmhl, K.; Nwngnd, G. Medical Formulas for Krimidanta (Dental Caries) in Indigenous Medicine in Sri Lanka—A Literary Review. *Int. J. Ayurveda Pharma Res.* **2016**, *4*, 52–56.
51. Velma, W.N.; Isabel, N.W.; Meshack, A.O.; Josphat, C.M. Isolation, Identification and Bioactivity of Fungal Endophytes from Selected Kenyan Medicinal Plants. *Afr. J. Microbiol. Res.* **2018**, *12*, 405–412. [[CrossRef](#)]
52. Pallie, M.; Perera, P.; Goonasekara, C.; Kumarasinghe, N.; Arawwawala, M. Efficacy and Safety of Freeze-Dried Form of *Tragia involucrata* L. Decoction in Treating Diabetes: A Randomized Controlled Clinical Trial. *Clin. Trials Degener. Dis.* **2020**, *5*, 31–36. [[CrossRef](#)]
53. Panda, D.; Dash, S.K. Phytochemical Examination and Antimicrobial Activity of Various Solvent Extracts and the Selected Isolated Compounds from Roots of *Tragia involucrata* Linn. *Int. J. Pharm. Sci. Drug Res.* **2012**, *4*, 44–48.
54. Hosahally, R.V.; Seru, G.; Sutar, P.S.; Joshi, V.G.; Sutar, K.P.; Karigar, A.A. Phytochemical and Pharmacological Evaluation of *Tragia Cannabina* for Anti-Inflammatory Activity. *Int. Curr. Pharm. J.* **2012**, *1*, 213–216. [[CrossRef](#)]
55. Alimuzzaman, M.; Ahmed, M. Analgesic Activity of *Tragia involucrata*. *Dhaka Univ. J. Pharm. Sci.* **2007**, *4*, 35–38. [[CrossRef](#)]
56. Pallie, M.S.; Perera, P.K.; Goonasekara, C.L.; Kumarasinghe, K.M.N.; Arawwawala, L.D.A.M. Evaluation of Diuretic Effect of the Hot Water Extract of Standardized *Tragia involucrata* Linn., in Rats. *Int. J. Pharmacol.* **2018**, *13*, 83–90. [[CrossRef](#)]
57. Alanazi, A.; Anwar, M.J.; Ahmad, M.A. Hepatoprotective and Antioxidant Activity of *Tragia involucrata* Root Extracts against CCl₄ Induced Hepatotoxicity in Rats. *Pharm. Lett.* **2015**, *7*, 146–152.
58. Ediriweera, E.; Ratnasooriya, W. ; A Review on Herbs Used in Treatment of Diabetes Mellitus by Sri Lankan Ayurvedic and Traditional Physicians. *Ayu* **2009**, *30*, 373–391.
59. Drewes, S.E.; Selepe, M.A.; Van Heerden, F.R.; Archer, R.H.; Mitchell, D. Unravelling the Names, Origins and Chemistry of “Muthis” Used for Male Sexual Disorders in KwaZulu-Natal, South Africa. *S. Afr. J. Bot.* **2013**, *88*, 310–316. [[CrossRef](#)]
60. Charlson, A.J. Antineoplastic Constituents of Some Southern African Plants. *J. Ethnopharmacol.* **1980**, *2*, 323–335. [[CrossRef](#)]
61. Ngcobo, M.; Gqaleni, N.; Naidoo, V.; Cele, P. The Immune Effects of an African Traditional Energy Tonic in In Vitro and In Vivo Models. *Evid. Based Complement. Altern. Med.* **2017**, *2017*, 1–14. [[CrossRef](#)]
62. Yineger, H.; Yewhalaw, D.; Teketay, D. Ethnomedicinal Plant Knowledge and Practice of the Oromo Ethnic Group in Southwestern Ethiopia. *J. Ethnobiol. Ethnomed.* **2008**, *4*, 11. [[CrossRef](#)]
63. Hassan-Abdallah, A.; Merito, A.; Hassan, S.; Aboubaker, D.; Djama, M.; Asfaw, Z.; Kelbessa, E. Medicinal Plants and Their Uses by the People in the Region of Randa, Djibouti. *J. Ethnopharmacol.* **2013**, *148*, 701–713. [[CrossRef](#)] [[PubMed](#)]
64. Abdela, G.; Sultan, M. Indigenous Knowledge, Major Threats and Conservation Practices of Medicinal Plants by Local Community in Heban Arsi District, Oromia, South Eastern Ethiopia. *Adv. Life Sci. Technol.* **2018**, *68*, 19.
65. Cheikhoussef, A.; Shapi, M.; Matengu, K.; Mu Ashekele, H. Ethnobotanical Study of Indigenous Knowledge on Medicinal Plant Use by Traditional Healers in Oshikoto Region, Namibia. *J. Ethnobiol. Ethnomed.* **2011**, *7*, 10. [[CrossRef](#)] [[PubMed](#)]
66. Setshogo, M.P.; Mberek, C.M. Floristic Diversity and Uses of Medicinal Plants Sold by Street Vendors in Gaborone, Botswana. *Afr. J. Plant Sci. Biotechnol.* **2011**, *6*, 69–74.
67. Vergiat, A.M. Plantes magiques et médicinales des Féticheurs de l’Oubangui (Région de Bangui). *J. Agric. Tradit. Bot. Appliquée* **1969**, *16*, 335–367.
68. Al-Fatimi, M. Ethnobotanical Survey of Medicinal Plants in Central Abyan Governorate, Yemen. *J. Ethnopharmacol.* **2019**, *241*, 111973. [[CrossRef](#)]
69. Mothana, R.A.A.; Abdo, S.A.A.; Hasson, S.; Althawab, F.M.N.; Alaghbari, S.A.Z.; Lindequist, U. Antimicrobial, Antioxidant and Cytotoxic Activities and Phytochemical Screening of Some Yemeni Medicinal Plants. *Evid. Based Complement. Altern. Med.* **2010**, *7*, 323–330. [[CrossRef](#)]
70. Desta, B. Ethiopian Traditional Herbal Drugs. Part III: Anti-Fertility Activity of 70 Medicinal Plants. *J. Ethnopharmacol.* **1994**, *44*, 199–209. [[CrossRef](#)]
71. Suryawanshi, V. Extraction and Isolation of Clerodane as a Bioactive Molecule from *Tragia Ramosa*. *J. Pharmacogn. Phytochem.* **2019**, *8*, 1135–1138.
72. Welcome, A.K.; Wyk, B.-E.V. An Inventory and Analysis of the Food Plants of Southern Africa. *S. Afr. J. Bot.* **2019**, *122*, 136–179. [[CrossRef](#)]
73. Magwede, K.; van Wyk, B.-E.; van Wyk, A.E. An Inventory of Vhavenda Useful Plants. *S. Afr. J. Bot.* **2019**, *122*, 57–89. [[CrossRef](#)]

74. Agbor, G.A.; Ndjib, R. Systematic Review of Plants Used Against Respiratory Diseases Related to COVID-19 in Africa. *J. Drug Deliv. Ther.* **2021**, *11*, 141–153. [[CrossRef](#)]
75. Amusan, O.O. Some Ethnoremedies Used for HIV/AIDS and Related Diseases in Swaziland. *Afr. J. Plant Sci. Biotechnol.* **2009**, *3*, 20–26.
76. Ogundare, A.O.; Olorunfemi, O.B. Antimicrobial Efficacy of the Leaves of *Dioclea Reflexa*, *Mucuna Pruriens*, *Ficus Asperifolia* and *Tragia spathulata*. *Res. J. Microbiol.* **2007**, *2*, 392–396. [[CrossRef](#)]
77. Tabuti, J.R.S.; Kukunda, C.B.; Waako, P.J. Medicinal Plants Used by Traditional Medicine Practitioners in the Treatment of Tuberculosis and Related Ailments in Uganda. *J. Ethnopharmacol.* **2010**, *127*, 130–136. [[CrossRef](#)] [[PubMed](#)]
78. Silberbauer-Gottberger, I. O Cerrado Como Potencial de Plantas Medicinai s e Tóxicas. *Oréades* **1982**, *8*, 15–30.
79. Ouöba, P.; Lykke, A.; Boussim, J.; Guinko, S. La flore médicina le de la Forêt Classée de Niangoloko (Burkina Faso). *Etudes Flor Vég Burkina Faso* **2006**, *10*, 5–16.
80. Reko, B. La Hierba de Quetzalcoat l. *Bot. Sci.* **1946**, *4*, 13–14. [[CrossRef](#)]
81. Jiofack, T.; Fokunang, C.; Guedje, N.; Kemeuze, V.; Fongnzossie, E.; Nkongmeneck, B.A.; Mapongmetsem, P.M.; Tsabang, N. Ethnobotanical Uses of Medicinal Plants of Two Ethnoecological Regions of Cameroon. *Int. J. Med. Med. Sci.* **2010**, *2*, 60–79.
82. Arnason, T.; Uck, F.; Lambert, J.; Hebda, R. Maya Medicinal Plants of San Jose Succotz, Belize. *J. Ethnopharmacol.* **1980**, *2*, 345–364. [[CrossRef](#)]
83. Seukep, J.A.; Ngadjui, B.; Kuete, V. Antibacterial Activities of *Fagara Macrophylla*, *Canarium Schweinfurthii*, *Myrianthus Arboreus*, *Dischistocalyx Grandifolius* and *Tragia benthamii* against Multi-Drug Resistant Gram-Negative Bacteria. *SpringerPlus* **2015**, *4*, 567. [[CrossRef](#)] [[PubMed](#)]
84. Anthony, S.T.; Ngule, M.C.; Jackie, O.K. In Vitro Antibacterial Activity of Methanolic-Aqua Extract of *Tragia Brevipes* Leaves. *Int. J. Pharm. Life Sci.* **2014**, *5*, 3289–3294.
85. Chepng'etich, J.; Ngule, C.; Jepkorir, M.; Mwangangi, R.; Njuguna, D.; Ndung'u, J.; Kiboi, D.; Mwitari, P. Total Phenolic Content and in Vitro Antiproliferative Activity of *Tragia Brevipes* (Pax) and *Tetradenia Riparia* (Hochst) Leaves Extract. *Eur. J. Med. Plants* **2018**, *22*, 1–10. [[CrossRef](#)]
86. Sivajothi, V.; Dakappa, S.S. In Vitro and in Silico Antidiabetic Activity of Pyran Ester Derivative Isolated from *Tragia Cannabina*. *Asian Pac. J. Trop. Biomed.* **2014**, *4*, S455–S459. [[CrossRef](#)] [[PubMed](#)]
87. Gupta, S.M.; Kumar, K.; Dwivedi, S.K.; Bala, M. Bioactive Potential of Indian Stinging Plants Leaf Extract against Pathogenic Fungi. *J. Complement. Integr. Med.* **2019**, *16*, 20170125. [[CrossRef](#)]
88. Perumal Samy, R.; Manikandan, J.; Al Qahtani, M. Evaluation of Aromatic Plants and Compounds Used to Fight Multidrug Resistant Infections. *Evid. Based Complement. Altern. Med.* **2013**, *2013*, 1–17. [[CrossRef](#)]
89. Thomas, R.; Megha, K.; Surya, P.; Rosalin, T.; Varghese, S.; Elyas, K. Investigation on the Biological Attributes of *Tragia involucrata* Linn. Using in Vitro Methods. *J. Pharmacogn. Phytochem.* **2021**, *10*, 398–404. [[CrossRef](#)]
90. Velu, V.; Das, M.; Raj, N.A.N.; Dua, K.; Malipeddi, H. Evaluation of in Vitro and in Vivo Anti-Urolithiatic Activity of Silver Nanoparticles Containing Aqueous Leaf Extract of *Tragia involucrata*. *Drug Deliv. Transl. Res.* **2017**, *7*, 439–449. [[CrossRef](#)] [[PubMed](#)]
91. Nivya, M.T.; Patil, R.K.; Rao, G.M.A.; Khandagale, A.S.; Somashekarappa, H.; Ananda, D.; Manjunath, H.; Joshi, C.G. Cytotoxicity Based Screening for Radioprotective Properties of Methanolic Extract of *Tragia involucrata* L. on Cultured Human Peripheral Lymphocytes Exposed to Gamma Radiation. *Indian J. Exp. Biol.* **2019**, *57*, 469–477.
92. Leonti, M.; Casu, L. Ethnopharmacology of Love. *Front. Pharmacol.* **2018**, *9*, 567. [[CrossRef](#)] [[PubMed](#)]
93. Attah, A.F.; Omobola, A.I.; Moody, J.O.; Sonibare, M.A.; Adebukola, O.M.; Onasanwo, S.A. Detection of Cysteine-Rich Peptides in *Tragia benthamii* Baker (Euphorbiaceae) and in Vivo Antiinflammatory Effect in a Chick Model. *Phys. Sci. Rev.* **2021**, 10151520200125. [[CrossRef](#)]
94. Bhattacharya, K.; Chandra, G. Phagodeterrence, Larvicidal and Oviposition Deterrence Activity of *Tragia involucrata* L. (Euphorbiaceae) Root Extractives against Vector of Lymphatic Filariasis *Culex Quinquefasciatus* (Diptera: Culicidae). *Asian Pac. J. Trop. Dis.* **2014**, *4*, S226–S232. [[CrossRef](#)]
95. Subramani, P.; Sampathkumar, N.; Ravindiran, G.; Rajalingam, D.; Kumar, B. Evaluation of Nephroprotective and Antioxidant Potential of *Tragia involucrata*. *Drug Invent. Today* **2009**, *1*, 55–60.
96. Joshi, C.G.; Gopal, M.; Kumari, N. Antitumor Activity of Hexane and Ethyl Acetate Extracts of *Tragia involucrata*. *Int. J. Cancer Res.* **2011**, *7*, 267–277. [[CrossRef](#)]
97. Islam, M.S.; Sana, S.; Haque, M.E.; Rahman, S.M.M.; Samad, A.; Noman, A.A.; Alam, R.; Rana, S.; Meem, R.I.; Mondol, D.; et al. Methanol, Ethyl Acetate and n-Hexane Extracts of *Tragia involucrata* L. Leaves Exhibit Anxiolytic, Sedative and Analgesic Activity in Swiss Albino Mice. *Heliyon* **2021**, *7*, e05814. [[CrossRef](#)]
98. Samy, R.P.; Gopalakrishnakone, P.; Houghton, P.; Ignacimuthu, S. Purification of Antibacterial Agents from *Tragia involucrata*—A Popular Tribal Medicine for Wound Healing. *J. Ethnopharmacol.* **2006**, *107*, 99–106. [[CrossRef](#)]
99. Varma, G.G.; Mathai, B.K.; Das, K.; Gowda, G.; Rammohan, S.; Einstein, J.W. Evaluation of Antiepileptic Activity of Methanolic Leaves Extract of *Tragia involucrata* Linn. in Mice. *Int. Lett. Nat. Sci.* **2014**, *12*, 167–179. [[CrossRef](#)]
100. Thimmaiah, N.; Joshi, C.; Patil, R.; Khandagale, A.; Somashekarappa, H.; Ananda, D.; Manjunath, H. Mitigation of Radiation-Induced Oxidative Stress by Methanolic Extract of *Tragia involucrata* in Swiss Albino Mice. *Pharmacogn. Res.* **2019**, *11*, 236. [[CrossRef](#)]

101. Dhara, A.K.; Suba, V.; Sen, T.; Pal, S.; Chaudhuri, A.K.N. Preliminary Studies on the Anti-Inflammatory and Analgesic Activity of the Methanolic Fraction of the Root Extract of *Tragia involucrata* Linn. *J. Ethnopharmacol.* **2000**, *72*, 265–268. [[CrossRef](#)]
102. Samy, R.P.; Gopalakrishnakone, P.; Sarumathi, M.; Ignacimuthu, S. Wound Healing Potential of *Tragia involucrata* Extract in Rats. *Fitoterapia* **2006**, *77*, 300–302. [[CrossRef](#)] [[PubMed](#)]
103. Dhara, A.K.; Pal, S.; Nag Chaudhuri, A.K. Psychopharmacological Studies on *Tragia involucrata* Root Extract. *Phytother. Res.* **2002**, *16*, 326–330. [[CrossRef](#)]
104. Samy, R.P.; Gopalakrishnakone, P.; Houghton, P.; Thwin, M.M.; Ignacimuthu, S. Effect of Aqueous Extract of *Tragia involucrata* Linn. on Acute and Subacute Inflammation. *Phytother. Res.* **2006**, *20*, 310–312. [[CrossRef](#)]
105. Sama, V.; Rajini, T.; Afrooz, H.; Balaraju, P.; Reddy, B.M.; Mullangi, R. Antihyperglycemic Effects of *Tragia Plukenetii* Ethanolic Extract. *Int. J. Pharm. Sci. Nanotechnol.* **2014**, *7*, 2436–2440. [[CrossRef](#)]
106. Kalaivanan, M.; Jesudass, L. Pharmacological Studies on Ethanol Extract of *Tragia Plukenetii* R. Smith. *IOSR J. Pharm.* **2012**, *2*, 1–7.
107. Muthuraman, M.; Dorairaj, S.; Rangarajan, P.; Pemaiah, B. Antitumor and Antioxidant Potential of *Tragia Plukenetii* R. Smith on Ehrlich Ascites Carcinoma in Mice. *Afr. J. Biotechnol.* **2008**, *7*, 3527–3530.
108. Manoharan, S.K. Evaluation of Anticonvulsant Activity of *Tragia Plukenetii* R. Smith Leaf Extracts against Chemoshock Induced by Pentylentetrazole in Mice. *Res. J. Pharm. Biol. Chem. Sci.* **2015**, *6*, 750–753.
109. Altemimi, A.; Lakhssassi, N.; Baharlouei, A.; Watson, D.; Lightfoot, D. Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants* **2017**, *6*, 42. [[CrossRef](#)]
110. Thangiah, A.S.; Mutuku, N.; Ngule, E.; Francis, R. Qualitative Analysis of Phytoconstituents in *Tragia Brevipes* Plant. *Int. J. Pharm. Res. Anal.* **2018**, *3*, 93–98.
111. Antony, C. Phytochemical and Spectral Study of the Medicinal Plant: *Tragia Plukenetii*. *J. Pharm. Res.* **2012**, *5*, 1701–1703.
112. Olaoye, S.B.; Ibrahim, A.O.; Zhiqiang, L. Chemical Compositions and Radical Scavenging Potentials of Essential Oils from *Tragia benthamii* (BAKER) and *Cissus Aralioides* (WELW). *J. Biol. Act. Prod. Nat.* **2016**, *6*, 59–64. [[CrossRef](#)]
113. Gobalakrishnan, R.; Kulandaivelu, M.; Bhuvaneshwari, R.; Kandavel, D.; Kannan, L. Screening of Wild Plant Species for Antibacterial Activity and Phytochemical Analysis of *Tragia involucrata* L. *J. Pharm. Anal.* **2013**, *3*, 460–465. [[CrossRef](#)] [[PubMed](#)]
114. Balogun, O.S.; Oladosu, I.A.; Liu, Z. Isolation of 2, 5-Dithia-3, 6-Diazabicyclo [2.2.1] Heptane and GC-MS Analysis of Silylated Extract from *Tragia benthamii*. *Ife J. Sci.* **2020**, *22*, 75–80. [[CrossRef](#)]
115. Alagar Yadav, S.; Ramalingam, S.; Jabamalai Raj, A.; Subban, R. Antihistamine from *Tragia involucrata* L. Leaves. *J. Complement. Integr. Med.* **2015**, *12*, 217–226. [[CrossRef](#)] [[PubMed](#)]