



AMARANTHACEAE PLANTS OF ISRAEL AND PALESTINE: MEDICINAL ACTIVITIES AND UNIQUE COMPOUNDS

Abdullatif Azab^{[a]*}

Keywords: *Amaranthaceae*, amaranthine, betanin, saponins, flavonoids, nerolidol derivatives, medicinal activities.

Amaranthaceae family is one of the largest and most diverse in the plant kingdom. While some of the plants of this family have important nutritional value, others are considered toxic and/or hazardous weeds and many efforts have been made in controlling them. But both categories were used by humans for traditional medicinal purposes, and some of them were extensively studied by modern science. In this review article, we will present some ethnomedicinal uses of the plants of this family, along with comprehensive literature survey of medicinal, biological and other activities. Information will be presented in tables for the convenience of readers, and many structures of natural products are presented. The extensive discussion section will focus mainly (but not only) on studies of active compounds production. This review article is about plants of this family in general and in particular about *Amaranthaceae* plants of our region. Conclusions and future research recommendations are also presented.

* Corresponding Authors

Fax: +972-4-6205906

Phone: +972-50-5650025

E-Mail: eastern.plants@gmail.com

[a] Eastern Plants Research Institute, Box 868, Arara, Israel
30026

INTRODUCTION

Amaranthaceae plant family is one of the largest in the plant kingdom. This family is botanically considered part of the *Amaranthaceae-Chenopodiaceae* alliance, where the *Amaranthaceae* family includes ca. 69 genera and 1,000 species and the *Chenopodiaceae* includes ca. 100 genera 1,400 species.¹

Archeological studies carried out in different locations suggest that *Amaranthaceae* plants were used by humans for various purposes. Las Canopas inhabitants used what the report authors refer to as *Amaranth* grains as food.² Despite the fact that specific species is not indicated, it is reported that these plants were domesticated and cultivated by these humans. They conclude that the plant species might be *Amaranthus hypochondriacus* or *A. cruentus*, with higher possibility of the second species. Rastogi and Shukla indicate in their comprehensive review article about human use of Amaranth plants that early archeological evidence shows that plants of this family were domesticated for nutritional uses, as early as 8000 years ago by Mayan civilizations.³ Pre-Hispanic (1000-1550 AD) of Northwest Argentina, domesticated *Amaranthus caudatus* for food.⁴ This conclusion was made based on isotopic (carbon and nitrogen) analysis of human remains found in that area. *Amaranthus retroflexus* grains were used as food, by humans who lived on the Eastern shores of the Mediterranean, in late Bronze age, 1550-1180 BC.⁵ Another early evidence of consumption of Amaranth by humans was found in archeological studies in Mexico, where these plants were eaten around 4000 BC.⁶

Some of the domesticated plants of the *Amaranthaceae* family were modified and selected mutations proved to be very successful and widely cultivated by humans. For example, Quinoa (*Chenopodium quinoa*) is one of the

healthiest grains, and its nutritional and economical importance is rising rapidly.⁷ Spinach (*Spinacia oleracea*) has been extensively studied, and its very high health benefits are well known.⁸ Another domesticated species of this family, which was thoroughly studied, and has very high nutritional value, is Beet (*Beta vulgaris*).⁹ Finally, some species on *Amaranthus* were domesticated for gardening and beauty purposes. A good example of these species is *Amaranthus tricolor*. Both domesticated and wild varieties of this species, are very beautiful and has notable medicinal and other biological activities.^{10,11} But this review article, will focus only on wild species of the *Amaranthaceae* family that grow in Israel and Palestine. These are *Achyranthes aspera*, *Aerva javanica*, *Alternanthera pungens*, *Alternanthera sessilis*, *Amaranthus albus*, *A. blitoides*, *A. blitum*, *A. cruentus*, *A. graecizans*, *A. muricatus*, *A. palmeri*, *A. retroflexus*, *A. rudis*, *A. spinosus* and *A. viridis*.



Figure 1. *Amaranthus retroflexus*

Table 1. Ethnomedicinal and ethnobotanical uses of *Amaranthaceae* plants.

Species, region	Region, uses, methods and references
<i>Achyranthes aspera</i>	India. Detailed review article about the many ethnomedicinal uses of this plant, with methods and plant parts. The publication presents some modern research findings. ¹² This publication is titled “research article” despite being a typical review article. Less detailed than ref. 12, but it provides detailed presentation of folk medicine uses of the plant, according to rural areas people. ¹³ Zambia. To treat HIV, cancer, pneumonia, cough, diarrhea, fungal infections of the skin and genital warts. Root infusion or whole plant decoction is administered orally, paste of the plant is applied to skin. ¹⁴ Pakistan. Whole plant decoction, extract or powder are used to treat kidney stone, pneumonia, chest pain, puncture wounds, ulcer, dysmenorrhea, aerodontalgia and asthma. ²⁷
<i>Aerva javanica</i>	Skin diseases, headaches, wounds, injuries, snake bite, insect stings, malaria, kidney stones and bone fractures. Leaf paste is applied on the skin, or infusions are taken orally. ¹⁵ Egypt. Roots and leaves are used to treat acne-like diseases and against “evil eye”. ¹⁶
<i>Alternanthera pungens</i>	Ghana. Leaves are ground and used for treatment of constipation and headaches. ¹⁷
<i>Alternanthera sessilis</i>	Pakistan. Leaves juice or whole plant decoction, or cooked, are used for eye pain, galactagogue, leucorrhoea, snake bite, diarrhea. ²⁷
<i>Amaranthus cruentus</i>	Pakistan. Eating cooked leaves has laxative effect. ¹⁸
<i>Amaranthus graecizans</i>	Egypt. Leaves are used against constipation. ¹⁶
<i>Amaranthus retroflexus</i>	Turkey. Leaves infusion is used for obtaining sterility. ²¹ Leaves are used as food. ^{22,23}
<i>Amaranthus spinosus</i>	Ghana. Whole plant is supplied as a decoction for treating malaria and fever. If crushed and used as a bath, it is used for baby refreshing. ¹⁷ Pakistan. Dried seeds powder is taken orally with water to treat eye vision weakness. ¹⁹ Aerial parts extract to treat diarrhea, antidote, fever. ²⁵ Whole plant decoction is used to treat gynecological disorders. ²⁶ Iran. This publication indicates the use of this plant in folk medicine but does not outline this use. It briefly presents some modern research results. ²⁰ Leaves are cooked, squeezed for juice, extracted or prepared as decoction, to use for vermifuge, dyspepsia, diuretic, odontalgia, cataract, constipation. ²⁷ Tanzania. Leaves are cooked with milk for ulcer treatment. ²⁴
<i>Amaranthus viridis</i>	Pakistan. Dried seeds powder is taken orally with water to treat eye vision weakness. ¹⁹ Leaves Extract, decoction, paste, are used to treat constipation, skin tonic, antidote, diuretic, cooking. ²⁵ Whole plant decoction is used to treat gynecological disorders. ²⁶ Leaves are extracted, powdered or prepared as a decoction, to treat painful urination, eye pain, constipation, piles, snakebite, cough and asthma. ²⁷ Leaves or roots juice or decoctions are used to treat inflammations. ²⁸ Leaves are crushed with sugar, mixed with tea to treat constipation. Extract is mixed with mustard oil to treat skin infections. ²⁹ Iran. This publication indicates the use of this plant in folk medicine but does not outline this use. It briefly presents some modern research results. ²⁰ Italy. Young leaves are used for pies. ³⁰

Ethnobotanical uses of *Amaranthaceae* family

Many human societies have discovered the nutritional and medicinal potential of these plants since many millennia. But compared with other plant families, the published articles about the ethnomedicinal and ethnobotanical uses of *Amaranthaceae*, are few. Moreover, some species were not mentioned at all (6 *Amaranthus* species). In Table 1, these published reports are summarized.

While studying the published review articles about the plants of *Amaranthaceae* plants, we noticed that unlike most other plant families that were reviewed, in the case of this family, there are more review articles about single species, than about the entire family. This is one of the main reasons to write the present review article.

In addition to several review articles about ethnobotany and ethnomedicine of these plants (see previous section), a large number of articles have been published, presenting

various properties, especially medicinal activities and chemical composition. The most important among them have been selected and their summaries are presented here.

Achyranthes aspera

This is the most studied and most reviewed of all plants of this family. One of the earliest review articles about this plant, is by Goyal and her colleagues.³¹ It presents the known medicinal activities of the plant in folk medicine as well as modern research. It provides general chemical composition, and its main importance is a table that presents the activities of different parts of the plant, their medicinal activities and active components (no structures), if known. Unlike that, the mini review article of Varuna *et al.*, introduces a very limited scope of the medicinal activities of this plant.³² It also provides very limited chemical composition, with one structure of Betaine (see Figure 2), with a minor error. There are several photos of the plant, but some of them are of very low quality.




Figure 2 Structure of Betaine, isolated from *Achyranthes aspera*.

Dey in his review article summarized enormous amount of scientific research and reports.³³ The article cites 265 publications, and it provides detailed chemical composition (no structures), and detailed presentations of botanical properties, traditional uses and reported medicinal activities of the plant. Compared with that, the review article of Shukla *et al* is highly valuable.³⁴ It contains detailed chemical composition, with structures of some important natural products, folk medicine uses and modern research reports, including highlighted antivenom activity. Among these natural products, there are few long chain alcohols (see Chemical Composition section). Tables are provided in the last pages of the article. There is one point that needs additional attention. Practically, there is no difference between sections 3 and 4. Both discuss chemical composition, despite having two different titles.

An excellent review article about *A. aspera* was published in 2011 by Srivastav and his colleagues.³⁵ It is comprehensive and presents all important aspects of this plant research, especially detailed chemical composition, with structures. The only weakness of this article is the detailed presentation of natural products names that occupied large space in the article, and were provided, anyhow, in the original research articles. On the contrary, the minireview of Hasan is very brief, does not present chemical composition and even the presentation of medicinal activities is very limited.³⁶ Photos of the plant are of low quality. In the review article of Srivastava, photos have high quality, chemical composition is extensively presented, medicinal activities are introduced, and traditional uses are mentioned.³⁷ There are two weakness points of this good review. First, many of the presented natural products have been isolated from other plants long time before detecting them in *A. aspera*, thus there was no need to show their structures. Secondly there is a mistake in the structure of Betaine, which is presented as an aldehyde, but it is an amino acid (see Figure 2).

In 2015, many review articles about this plant were published. Sharma and Chaudhary³⁸ presented the three major aspects of this plant (chemical composition, but no structures provided, ethnobotanical uses and modern research findings. They also presented a good comparison between traditional medicine and recent scientific medicinal activities. The minireview of Sureshkumar has the same structure.³⁹ The short minireview of Parmar and D. Sharma lacks ethnobotanical uses.⁴⁰ Ghimire and her colleagues published a large, good review article, in terms of chemical composition (some structures are presented) and modern research results of pharmacological activities.⁴¹ It does not indicate ethnomedicinal uses. The review of Sanjay is disappointing. It looks like a copy of ref. 36 with exactly the same low quality photos.⁴²

The review article of Vijayaraj and Vidhya, lists most of the active chemical components of the plant, presents many medicinal activities, but does not refer to folk medicine.⁴³

Anyhow, in our humble opinion, this article is of poor quality. It has two major weaknesses. First, the photos, which evidently were copied from other documents, have very low quality, including those that present chemical structures.

Second, the journal classifies the article as “review”, and the title of the publication is “Biological Activity of *Achyranthes Aspera* Linn. - A Review”, but in the introduction, there is a section titled “Collection and Authentication of Plant”, and a procedure for that is described. It is not clear why authors collected samples of the plant for writing review article. In the same year Saini published a very small review article about *A. aspera* that lists most of the active natural products, some of the published medicinal research articles, but did not relate to traditional medicine.⁴⁴

Later in 2016, Verma published her review article.⁴⁵ This is a very short publication, that if informational part is extracted, it will not be more that two pages. But as far as we understand the term “review article”, even though its size is very small, this publication should not be named this way. It lists very partial chemical composition, does not mention traditional uses, cites only two publications (treatment of blindness and anticancer activity), and concludes that *Achyranthes aspera* L. (Latjeera) is a very important medicinal herb.

The size of the review article of Lakshmi and his colleagues is the same as ref. 45, three pages, but its quality is much higher.⁴⁶ After a short introduction, high quality photo, there is a brief presentation of major medicinal activity. Chemical compositions as well as traditional uses are indicated in brief. Another excellent review article was published in the same year by Rehman and her colleagues.⁴⁷ It presents all major aspects of interest about this plant. The two notes that should be given here are, first, there are many unnecessary common names, and, despite presenting detailed chemical composition, no structures were provided. Finally, the publication of Singh *et al.* is comprehensive and detailed for all aspects, but it lacks structures of important natural products that this plant contains.⁴⁸

Aerva javanica

As far as our literature scan could reach, we found only two relevant review articles about this plant. First, the article of Vinit and Zaveri.⁴⁹ It focuses more on botanical studies, and lacks structures of important natural products, but the overall value of this article is high. But the review article of Payal and colleagues is low in quality.⁵⁰

First, it is a 4 pages minireview titled “A Review on Phytochemistry and Biological Activities of *Aerva*”, but most of the article presents photos and botany of the plant. Second, the presented structure on Narcissin includes R group, without explaining what R stands for (L-rhamnosyl-β(1-6)-D-glucosyl-β-1- group). Third, the presented structures are of natural products contained in many other plants, while one of these natural products that were mentioned in the article, and contained only in the genus of *Aerva*, aervitin, was not presented (see Figure 3). Fourth, the presented structure of feruloyltyramine is wrong (see Figure 3).




Figure 3. Structures of avertin and feruloyltyramine

Alternanthera sessilis

Except for deserts and other arid landscapes, this plant is relatively widespread in the reviewed region, Israel and Palestine, but it is a typical fresh water-neighboring plant. The review article of Walter and his colleagues provides a limited view of this plant.⁵¹ It presents botanical introduction, traditional uses, with a notable table that lists all Indian traditional medications (Siddha) that contain this plant or parts of it. But the presentation of modern research results is very partial. And the major weakness of this article is the presentation of the phytochemistry. The natural products that are introduced (β -Carotene and three natural phytosterols) are present in other plants in much higher concentrations and are not typical of this plant.

The review article of Laxmi *et al.*, is misleading, the least to say but it can be classified as scientific fraud.⁵² It provides short botanical introduction, mentions chemical composition and some traditional uses, including for nutritional purposes. In this introduction, it cites 11 articles, including ref. 51 here. This is followed by the main part of that article, titled “activities reported of *Alternanthera sessilis*”. Here the reporting is exactly as in research articles, including very detailed experimental procedures, and even plant collection and identification. No references cited in this section, and readers could be misled to think that this is an original article. The presentation of each activity includes all research steps, discussion and conclusions. It is not easy to believe that all these investigations were done in a single work. But the detection of misleading is quite easy: there are many mistakes of copying and pasting from original articles. For example, on page 3 of the article, page 2847 of the journal issue, there is a table titled “table 2”. But the table was copied as an image from the original article of Kumar and Sanjib.⁵³ There, it appears as “table 1” (page 4), and Laxmi *et al.* did not pay attention to that numbering. Ref. 52 includes many other faults.

Rao focused in his review article on the botanical and ethnobotanical aspects of this plant.⁵⁴ The article mentions very briefly partial chemical composition (no structures) and few pharmacological activities. Contrary to that, the review article of Shehzad and his colleagues is of a high quality, comprehensiveness and photos.⁵⁵ Its only weakness is that it does not present unique natural products of this plant, and it does not present any chemical structures. The minireview of Chandrashekhar is satisfactory for this size article, but it

lacks chemical structures of important active natural products.⁵⁶

Amaranthus cruentus

This impressive plant annual, 70 cm high on an average, is very widespread in reviewed region (see Figure 1) and yet only one review article was found about it. The article of Wolosik and Markowska is of very high quality, despite having two weaknesses.⁵⁷ Its major strength is its comprehensiveness in terms of presenting the topics mentioned in the title. Its first weakness is that it focuses on big and macromolecules (fats, saccharides, amino acids and peptides), but almost ignores small molecules that are contained in the seeds and have notable medicinal activities. Secondly, the structures and the activities of the mentioned small molecules are not presented.

Amaranthus palmeri

Except for very dry areas, this plant is very widespread all over the reviewed region. It is immensely consumed by all plant eating animals. But despite that, only one review article was published about it by Ward and her colleagues.⁵⁸ This article focuses on the biological and agricultural aspects of this plant, and does not present traditional uses, medicinal activities or chemical composition. The only relation that it has to this section, is its high resistance to synthetic herbicides, such as the compounds shown in figure 4.




Figure 4. Herbicides that *Amaranthus palmeri* showed resistance to (ref. 58).

Amaranthus spinosus

It is considered invasive species in the reviewed region,⁵⁹ like all plants of *Amaranthus* genus, but *A. spinosus* is one of the most recent invaders, unlike some others (*A. rudis*), that invaded this region many centuries ago. It can be easily distinguished from other *Amaranthus* species, that all thrive in cultivated areas, since *A. spinosus* is spiny and painful if mistakenly touched.

The first review article that we present here about this plant, was published by Jhade and his colleagues.⁶⁰ It is comprehensive with wide medicinal activities presentation

but limited in review of traditional uses and chemical composition. The review by Kawade and his colleagues, that was published 4 years later, has basically the same structure and size.⁶¹ The traditional uses presentation is even smaller than in ref. 60, and the phytochemical section is slightly larger, but despite mentioning unique natural products to this plant (spinoside), no structures are presented. A very good review was published by Kumar and his colleagues.⁶² It includes all expected and needed informational parts, with high quality photos and structure of unique natural products to this plant (Betanin and Amaranthine, Figure 5). It lacks a discussion section. And in the same year, Tanmoy and her colleagues published their review article.⁶³ It is comprehensive with structures of many natural products, but the images are vague and unclear. In addition, the traditional uses section is very small, but all other parts are satisfactory.




Figure 5. Betanin and Amaranthine isolated from *A. spinosus* (ref. 62).

Despite being properly written and arranged, we found the review article of K. Chandrashekhar, having low usefulness for readers that do not know Hindi.⁶⁴ The article have the requisite components needed for such a review, but tables 1 and 2 are meaningless for English readers. Since the journal is titled “International Journal of Ayurvedic Medicine”, and since the article is in English, translation of words from Hindi is totally inadequate. We are fully aware that such international articles can include some local common names, but not whole tables, which are supposed to be important parts of the text. Finally, the review article of Raut is superb. It is comprehensive, includes all needed components, well arranged, very useful tables and high-quality photos.⁶⁵

Amaranthus viridis

This plant is unmistakable, it grows around and in human habitats, flowering almost all year except for few cold weeks in the winter, has serrated dark green leaves with white coloring that has the leaf shape.

The minireview of Sowjanya and her colleagues about this plant is informational, satisfactorily written, but has three weaknesses.⁶⁶ First, the traditional uses part is very partial. Second, even though, the article is supposed to present “phytochemical and pharmacological potential” of this plant, the phytochemistry section is notably limited (one paragraph) and does not present structures of natural products. Third, the citation of literature is not appropriate. For example, in the paragraph of phytochemistry, the components of methanolic extract of leaves are presented and an article is cited. Directly after that, in the same paragraph, there is a brief description of the content of the roots of the plant and no reference is cited.

A year later, Ferdous and his colleagues published another minireview (that they titled as “comprehensive review”) about this plant.⁶⁷ It includes useful botanical information, relatively enough information about medicinal activities, high-quality photos, but lacks the parts of traditional uses and phytochemistry. Finally, the minireview of Singh and her colleagues provides very limited information about this plant, probably due to its small size (3 pages), and literature citation is poor, especially in the traditional uses part.⁶⁸

Amaranthaceae Family

M. Graca Miguel published an outstanding review about the family of natural compounds, Betalains, in the *Amaranthaceae* plant family.⁶⁹ But since this family includes many genera that some of them are not represented in the reviewed area, some valuable information presented in this superb review, is not relevant to our review. Anyhow, it presents this natural products family from different and comprehensive points of view. From this information, we present here the classification of Betalains, which is shown in Figure 6. So, Betanin and Amaranthine shown in Figure 5, are part of the Betalains subfamily of Betacyanins.

There are no published review articles for *Alternanthera pungens*, *Amaranthus albus*, *Amaranthus blitoides*, *Amaranthus blitum*, *Amaranthus graecizans*, *Amaranthus muricatus*, *Amaranthus retroflexus* and *Amaranthus rudis*.




Figure 6. General structures of Betalains (ref. 69).

Amaranthus genus

The review article of Alegbejo is informative regarding the species that the author chose to review.⁷⁰ But it is also confusing. In the abstract she states that “Amaranth consists of 60-70 species, 40 of which are considered native to the Americas”. Practically, this information is inaccurate: there are about 60 species native to America.⁷¹ Using any count, the author did not review most of these species. In addition, the author is from Nigeria, and she relate to traditional Nigerian medicinal uses of *Amaranthus*, that are not known to the traditional medicine of American peoples. The article lacks presentation of active ingredients and the traditional medicine part is very limited. Comparing with that, the review article of K. Peter & P. Gandhi is an excellent one.⁷² It is very clear, highly informative with excellent figures. The only weakness of it is that traditional medicine and phytochemistry sections are very limited.

Biological, chemical and other properties of *Amaranthaceae* plants

When studying the published research literature about the plants of *Amaranthaceae* family that grow wild in the reviewed region, the result is quite interesting and not easy to understand. While some of these plants were extensively studied for almost every possible property, some others are completely missing in this type of scientific literature. These “missing” species are not particularly rare or highly toxic. We will refer to this fact in the discussion section.

So, to make it easier for scholars and interested readers, we summarized the published properties of each species in a separate table.

The data related to *Achyranthes aspera*, *Aerva javanica*, *Alternanthera pungens*, *Alternanthera sessilis*, *Amaranthus cruentus*, *Amaranthus graecizans*, *Amaranthus retroflexus*, *Amaranthus spinosus* and *Amaranthus viridis* are given in tables 2-10, respectively.

Amaranthus blitoides, in this interesting report, the allelopathic effect of this plant on common bean (*Phaseolus vulgaris*) was studied by three methods. First, the two plants were grown together. Second, shoots aqueous extract of *A. blitoides* was supplied to common bean seedlings, in greenhouse conditions. Third, active allelochemicals were extracted with ethanol from the aqueous extract, and they were administered to common bean seedlings. In all cases, the allelopathic effect was clearly observed.²⁵⁵

The general chemical composition of the plant with focus on antioxidants and minerals of *Amaranthus blitum* has been reported. Antioxidant capacity of ethanolic leaves extract was determined (DPPH) and its high value confirmed the high values of the antioxidant compounds concentrations.²⁵⁶

Amaranthus muricatus

This plant was very scarcely studied. Nutrients and antinutrients were determined in aerial parts before flowering.²⁸⁰ General chemical composition was studied twice by the same research group.^{281,282}

Amaranthus palmeri

This species was also very limitedly investigated. Water soluble organic compounds were isolated by solid-phase separation techniques, and they had slow, but clear allelopathic effect on onions and carrots.²⁸³ A follow up study showed that volatiles (detailed GC analysis) that were released from aerial parts, had allelopathic effect on onion.²⁸⁴ Since this plant has allergic effect on humans, a study revealed that the allergen in the pollen is profilin (14 kDa protein).²⁸⁵ Finally, to the best of our knowledge, there are no publications that present novel natural products that were isolated from this plant, though the general chemical composition has been published.²⁸⁶

No research articles relevant to this review have been published regarding *Amaranthus albus* and *Amaranthus rudis*.

DISCUSSION

After scanning the published literature about the *Amaranthaceae* plants of Israel and Palestine, the general but the very clear fact that arises is that some of these plants were extensively studied, while others were completely “ignored”. We have indicated this explicitly in the previous and main section of this review article.

The plant species of the family that was and still being most investigated is undeniably *Achyranthes aspera*. Evidently, the major mass of research and publication about this plant originate from India. Among the numerous medicinal properties of this plant, using its products as snakebite treatment is very notable. This use, as we have indicated earlier, is known in traditional Indian medicine and in modern research in this country. Based on the shocking, annual number of snakebite cases in this country, and even the more astonishing annual number of fatalities, 58000 in average,⁴⁰¹ it is very understandable why the use of this plant to treat snakebites is very common in India.

A. aspera is mentioned quite frequently in ethnomedicine literature of snakebite treatment,⁴⁰² along with modern research. But based on the availability of this plant (very widespread) and the number of deaths, this treatment needs very extensive research in order to make more effective.

The chemical composition of this plant was extensively studied, and in addition to know natural products, some very interesting compounds were isolated and characterized from its different parts (see Figure 7). But some of the reports about this chemical composition are misleading.

Generally, for the identification of a known compound in plant extract or essential oil, HPLC device and standards preparations are all what is needed. It is almost obvious that when detailed spectroscopic characterization is reported, they refer to a novel compound, which is not the case in some reports.⁴⁰³

In the previous section we have cited some reports of general chemical composition, and many more were published but we chose not to cite them. Here is another one.⁴⁰⁴

The *Achyranthes* genus is represented by a single species (among 16 in the world), *A. aspera*, and it is the same for the genus of *Aerva*, it is also represented by a single species (11 globally), *A. javanica*. This plant was moderately studied. But despite this, novel, interesting natural products were isolated from this plant (Figure 8). For this reason, there were many attempts to grow and cultivate it. In our region, anyhow, it's a desert and arid regions plant, so it is not widespread but it is very well adapted to this habitat.⁴⁰⁵

Table 2. Biological, medicinal and other properties of *Achyranthes aspera*.

Activity/Property	Major Findings/Reference
Analgesic and pain related activities	Aerial parts were extracted with methanol. The extract has analgesic activity in acetic acid-induced writhing test. ⁷³ Leaves were extracted with 95% aqueous ethanol, and it proved antinociceptive in rat. Three pain-inducing methods were used: tail flick response, hot plate and formalin. ¹¹¹ Roots were extracted with 95% aqueous methanol and extract (oral administration) was found antinociceptive in mice (acetic acid induced writhing). ¹¹²
Antiallergy, skin protection, wound healing	Allergy was induced in mice by potassium dichromate and treated with methanolic whole plant extract. Positive results were recorded. ⁷⁴ Seeds were extracted successively with petroleum ether, chloroform and methanol. Extracts found irritant to the inner surface of rabbit ear. ¹²⁶ An ointment was prepared by using 5% leaves methanolic extract. This formulation was used to treat wounds (burn, wound) in healthy and diabetic (STZ-induced) mice. In both cases, the ointment showed clear wound healing activity. ¹²⁸ Seeds were defatted with petroleum ether and extracted with 90% aqueous ethanol. Extract had wound healing activity (rats) in three tests: excision, incision, and burn wound. ¹²⁹
Antibacterial, antifungal, antiviral and related activities	Stem and leaves were separately extracted with chloroform, methanol, ethanol and ethyl acetate, to obtain 8 extracts. None of them had any antibacterial activity against 5 bacteria species. ⁷⁸ Leaves and callus were extracted with ethanol, methanol, chloroform, benzene and petroleum ether. Contrary to the previous report, ethanol and chloroform extracts were active against all five species of tested bacteria, including <i>E. coli</i> , which ref. 78 reported no activity against it. All callus extracts except petroleum ether, were active against all bacteria species. ⁷⁹ 70% Aqueous ethanol was used to extract leaves and shoots. This extract showed activity against 3 types of bacteria. ⁸⁰ This interesting report, tested three variables. First, the yield of extraction of roots, stem, leaves and seeds, with hexane, ethyl acetate, methanol, ethanol and water; in three different months: January, May and September. Highest yields were in January. Second, antibacterial activity of these extracts against several bacteria species. Third, hexane stem extract was most active. ⁸¹ Trypsin inhibitor was isolated from seeds by standard protocol. It was tested and found active against 6 bacteria species and 4 fungi. ⁸² Stems and leaves were extracted separately with petroleum ether, chloroform, ethyl acetate, ethanol and methanol. All eight extracts were tested and found active against 3 bacteria species, except stem extracts that showed no activity against <i>Enterococcus faecalis</i> . ⁸³ Used part of the plant was not indicated by authors, but in this important report, plant material was extracted with water, ethanol, methanol and chloroform. None of these extracts was successful antibacterial against all 6 species of multi-drug resistant clinical bacteria isolates, but none of them was completely inactive. ⁸⁴ Leaves, roots, stem and flowers were extracted separately with petroleum ether, benzene, chloroform, ethyl acetate, ethanol and water. All extracts were partially active against cultured bacteria, but none was active against all species. ⁸⁵ Aqueous extract of stems and roots was active against <i>Streptococcus mutans</i> . ⁸⁶ Leaves were extracted with diethyl ether, ethyl acetate and acetone, separately. All extracts showed very mild activity against 2 bacteria species, but none was active against <i>Bacillus subtilis</i> . ⁸⁷ Leaves and stems were extracted separately with petroleum ether, methanol, ethanol, ethyl acetate and chloroform. All extracts were active against 3 bacteria species, except stem extracts that were inactive against <i>E. faecalis</i> . ⁸⁸ Methanol, chloroform and petroleum ether extracts of leaves were prepared, and tested against several species of bacteria and fungi. It is reported that (contrary to previous reports), only methanolic extract was active. In addition, extraction yields are notably low. ⁸⁹ Acetone and aqueous extracts of leaves that were collected from different areas and mixed, were prepared. These extracts were active against 5 bacteria species. Extracts were analyzed for phenolic acids. ⁹⁰ Seeds were extracted sequentially with chloroform, petroleum ether, acetone, methanol and water. All extracts had antibacterial activity against 8 bacteria species, except aqueous extract that had no activity against <i>Klebsiella pneumoniae</i> . ⁹¹ This report is very interesting, since part of its result contradict most of previous cited reports. Leaves were extracted separately with water, ethanol and methanol and tested against 12 species of bacteria and fungi. All aqueous extracts were completely inactive, while other extracts had notable activity. ⁹² Whole plant methanolic extract was active against 4 fungi species. ⁹³ Ethanolic leaves extract had notable activity against 3 fungi species. ⁹⁴ 17-Pentatriacontanol (see Figure 7) was isolated from shoots, and it had activity against <i>Aspergillus carneus</i> . ⁹⁵ Methanolic extract (plant part not indicated) was active against HIV viruses. ⁹⁶
Anticancer and related activities	Whole plant was extracted separately with petroleum ether, chloroform, ethanol and water. These extracts had cytotoxic activity (brine shrimp lethality), where aqueous extract was most active. ⁹⁷ Roots were extracted separately with water and ethanol. Both extracts had anticancer activity against human colon cancer cells (COLO-2015). Aqueous extract had higher activity. ⁹⁸

Antidiabetic, anti-obesity and related activities	<p>Whole plant and methanolic extract were orally administered to normal and alloxan-induced diabetic rabbits. In both methods, significant lowering of blood glucose was recorded.⁹⁹ Leaves ethanolic extract had significant effect on STZ-induced diabetes in rats.¹⁰⁰ Plant commercial extract (part/s not indicated) had advanced inhibition of formation of glycation end products.¹⁰¹ Leaves were extracted with 95% aqueous ethanol, and extract was successively fractionized with <i>n</i>-hexane, chloroform and <i>n</i>-butanol. All extracts were active against STZ-induced diabetes in rats, where positive controls were metformin and glybenclamide. In addition, four pure, known compounds that were isolated from extracts were tested and found active antidiabetic: oleolenic acid, ursolic acid, β-sitosterol and 17-triacontanol.¹⁰² Herbal tea of leaves stems and flowers (combined) was prepared and administered to diabetic (alloxan-induced) rats. It had clear antidiabetic and hypolipidemic activities.¹⁰³ Ethanolic seed was supplied to diabetic rats (STZ-induced) and significant increase in blood glucose, as well as lower levels of blood lipids, were observed.¹⁰⁴ Saponin-rich fraction was obtained by extraction with 70% aqueous ethanol and 1-butanol, successively. This fraction has anti-obesity activity in high fat diet fed rats.¹¹³ Powdered seeds were orally administered to high doses of fructose-fed female rats. Hypolipidemic effect was recorded.^{137,138} Triton-induced hyperlipidemia in rats was treated with separately with aqueous and ethanolic leaves extracts. The alcoholic extract had significant effect, while the aqueous extract had none.¹³⁹</p>
Antioxidant, anti-inflammatory and related activities	<p>Whole plant was extracted with methanol and extract was fractionized with several organic solvents. Saponin-rich fraction was found active against FCA-induced arthritis. FCA is mycobacteria.⁷⁶ Leaves were defatted with petroleum ether and extracted with water. The aqueous extract was active against formaldehyde-induced arthritis.⁷⁷ Seeds were extracted sequentially with chloroform, petroleum ether, acetone, methanol and water. All extracts showed antioxidant activity (DPPH), where methanolic extract had highest capacity.⁹¹ Ethanolic seed extract was supplied to diabetic rats (STZ-induced) and significant decrease in the levels of superoxide dismutase and glutathione (GSH) peroxidase and increase in the levels of vitamin E, catalase, and reduced GSH were observed.¹⁰⁴ Ethanolic leaves extract was tested on carrageenan-induced hind paw edema and cotton pellet granuloma models in rats. In both cases, notable anti-inflammatory activity was recorded.¹⁰⁶ Roots ethanolic extract had anti-inflammatory activity in rats, that was observed in two tests: carrageenan-induced paw edema and cotton pellet granuloma.¹⁰⁷ Leaves and whole plants were separately extracted with water, and both extracts had anti-inflammatory activity in carrageenan-induced paw edema in rats.¹⁰⁸ Leaves were extracted with 80% aqueous ethanol, and extract had activity against inflammatory cytokines TNF-α and IL-5 in the lungs of mice.¹⁰⁹ In this important report, gel was prepared (procedure is provided) using aqueous roots extract. The gel was used to treat patients with chronic periodontitis and proved strong activity.¹¹⁰ Methanolic extracts of leaves and roots were prepared and had high antioxidant activity (DPPH, ascorbic acid standard).¹¹⁴ Stems were successively extracted with <i>n</i>-hexane, chloroform, methanol and water. Antioxidant was tested (DPPH) and methanolic extract had highest capacity. Oddly enough, authors report that extracts were tested against 4 bacteria species, and only <i>n</i>-hexane fraction was active against one species.¹¹⁵ Leaves were extracted with 50% aqueous ethanol, and extract had high antioxidant activity (DPPH, NO radical inhibition).¹¹⁶ Roots and flowers were extracted separately and sequentially with petroleum ether, benzene, chloroform, ethyl acetate, ethanol/water. General chemical composition of extracts was determined as well as antioxidant capacity (3 methods).¹¹⁷ Leaves and stems were separately extracted with water, and extracts showed notable antioxidant (DPPH) capacity.¹¹⁸ Ethanolic leaves extract was prepared, analyzed for general chemical composition, and tested for antioxidant capacity (phosphomolybdenum assay) was determined.¹¹⁹ This study reports a comparison between the antioxidant capacity of <i>A. aspera</i> and <i>Cyathulaprostrata</i> that grow in Sri-Lanka. For both species, aqueous extract of whole plant was prepared, general chemical composition was determined, and antioxidant capacity was tested by four methods.¹²⁰ Whole plant was separately extracted with dichloromethane, ethyl acetate, methanol and water. All extracts were analyzed for detailed chemical composition, yet, no new compounds are reported. All extracts were tested for antioxidant capacity (4 methods).¹²¹ Leaf aqueous extract was orally administered to rats, resulting reduction in liver lipid peroxidation and prothyroidic activity.¹²²</p>
Brain related activities	<p>Aerial parts were extracted with methanol and the extract had CNS depressant activity tested by three methods.⁷³ Leaves were extracted with methanol and the extract showed anxiolytic activity. Authors refer this activity to the presence of alkaloids in the extract.⁷⁵ Roots were extracted with 95% aqueous methanol and extract (oral administration) was found neuroprotective in mice (pentobarbital sleeping test, open field test).¹¹² Extracts and pure natural products from the plant were tested for nootropic activity (radial arm maze, step through latency in passive shock avoidance and increased recognition index). Results indicated modulating brain glutamatergic and cholinergic neurotransmission.¹²⁷</p>

Cardiovascular system related activities Digestive system protection, hepatoprotection, nephroprotection, lung protection	Leaves were extracted with 70 % aqueous methanol and extract was administered to mice. As a result, a significant increase in lymphocyte and platelet counts were recorded. ¹²⁵ Ethanol and aqueous leaves extracts were administered to mice that had diarrhea induced by castor oil. Extracts had laxative as well as bronchodilator effects. ¹⁰⁵ Roots were extracted with 95 % aqueous methanol and extract (oral administration) was found diuretic in mice. ¹¹² Aqueous extract of roots prevented nucleation of calcium oxalate (urolithiasis) and preventing renal epithelial NRK-52E cells injury. ¹³⁰ Leaves were extracted with petroleum ether, methanol, ethanol and water, but only ethanolic and aqueous extracts were used for <i>in vivo</i> studies. These extracts were orally administered to rats, and diuretic effect was recorded. ¹³¹ Whole plant was successively extracted with petroleum ether, chloroform, ethyl acetate, methanol and water. All extracts were tested for diuretic activity in rats, and methanolic extract was most active. ¹³² Aqueous extract of seeds had diuretic effect on mice and rats. ¹³³ Leaves were extracted with 70 % aqueous ethanol and this extract had protective effect against pylorus ligation and chronic ethanol-induced ulcer in rats. ¹³⁴ Whole plant was extracted with 50 % aqueous ethanol and this extract had protective effect against N-nitrosodiethylamine and CCl ₄ -induced hepatocarcinogenesis in rats. ¹³⁵ Seeds were defatted with petroleum ether and extracted with 70 % aqueous ethanol. The extract had hepatoprotective activity against CCl ₄ -induced toxicity in rats. ¹³⁶
Enzyme inhibition, antivenom and related activities	Whole plant was separately extracted with dichloromethane, ethyl acetate, methanol and water. All extracts were tested for as enzyme inhibitors: AChE, BChE, α -amylase, α -glucosidase and tyrosinase. ¹²¹ Aqueous and ethanolic extracts of leaves were prepared and tested against snake (Russell's viper and Saw scaled vipe) in mice. This activity was tested by inhibition of venom phospholipase and procoagulant activity. ¹²³ Aqueous extract of stems inhibited protease and phospholipase A2 from the venom of <i>Bitis arietans</i> . ¹²⁴
Insecticidal, molluscicidal, antiparasitic, herbicidal	Acetone and aqueous extracts of leaves that were collected from different areas and mixed, were prepared. These extracts had anthelmintic activity against <i>Caenorhabditis elegans</i> . ⁹⁰ Various extracts and their combinations were used as herbicidal against 6 weed species. In all cases, positive results were recorded, and highest efficiency was found for fruit extract. ¹⁴⁰ Leaves were extracted separately with acetone, chloroform, ethyl acetate, <i>n</i> -hexane and methanol. All extracts had moderate insecticidal activity against <i>Aedes aegypti</i> and <i>Culex quinquefasciatus</i> . A pure saponin that was isolated from the ethyl acetate extract, had strong activity ¹⁴¹ and it was identified as a compound that was isolated from <i>Achyranthes bidentate</i> , a species that is not included in this review. ¹⁴² Contrary to ref. 141, authors of this report published that leaves <i>n</i> -hexane extract had strong activity against <i>Aedes aegypti</i> . ¹⁴³ Leaves and stems were separately extracted with <i>n</i> -hexane, and both extracts had significant activity against <i>Aedes aegypti</i> . ¹⁴⁴ Aqueous extract of leaves had strong molluscicidal effect against <i>Biomphalaria apfeifferi</i> and <i>Lymnaea natalensis</i> . ¹⁴⁵ Pure natural product (hormone) Ecdysterone (see Figure 7), that was isolated from ethyl acetate leaves extract, has strong activity against <i>Aedes aegypti</i> . ¹⁴⁶
Metals related activities	Methanolic leaves extract inhibited corrosion of mild steel that was exposed to industrial water medium. ¹⁴⁷ Aqueous leaves extract was used to reduce silver ions (AgNO ₃), to produce nanoparticles (AgNPs), which had notable larvicide against <i>Aedes aegypti</i> . ¹⁴⁸ Aqueous leaves extract was used to reduce gold ion (HAuCl ₄), to produce nanoparticles (AuNPs), which proved nontoxic, so authors conclude that they can be used for drug delivery. ¹⁴⁹ Aqueous extract was used to prepare CuO nanoparticles, which had notable inhibition of cellular adhesion (<i>E. lenta</i> and <i>E. aerogenes</i>). ¹⁵⁰
Nutrition, toxicity Reproductive system, sex	Leaves powder and their methanolic extract were fed to mice and no toxicity was recorded. ¹⁵¹ Roots were extracted with 50 % aqueous ethanol and extract was orally administered to male rats. Results indicated reduction of sperm counts, weight of epididymis and reproductive system hormones. ¹⁵² Roots were successively extracted with petroleum ether, chloroform, ethanol and water. Chloroform and ethanolic extracts had antifertility activity in female rats. ¹⁵³ Ethanolic extract of roots that was fed to pregnant female rats had 100 % combined antifertility activity (anti-implantation and abortifacient). ¹⁵⁴ Leaves were extracted with 80 % aqueous methanol. This extract was administered to female rats, and it had combined antifertility effect, but it did not reduce reproductivity hormones. ¹⁵⁵ Aqueous ethanol (50 %) was used to separately extract roots of <i>A. aspera</i> and <i>Stephania hernandifolia</i> (not included in this review). A composite of 1:3, respectively, was prepared and orally administered to male rats. The result was clear antifertility effect. ^{156,157} Roots were extracted with 50 % aqueous ethanol, resulting the isolation of 58-kDa protein. Its composition is not reported. This protein had clear antifertility activity. Gossypol was used as control antifertility agent in male mice. ¹⁵⁸ A follow up study with the same protein, showed that it has antifertility effect on human males. ¹⁵⁹ Similar study to references 156 and 157. ¹⁶⁰

Chemical composition

Selected articles that reported general chemical composition, which means, natural products families (phenolics, carbohydrates ... etc.), minerals or detailed chemical composition that included known compounds, without reporting novel natural products and their characterization.¹⁶¹⁻¹⁶⁸ First report of the isolation and the characterization of Ecdysterone from roots methanolic extract (Figure 7) that we mentioned earlier in ref. 146.¹⁶⁹ 36,47-Dihydroxyhenpentacontan-4-one (Figure 7) was isolated and characterized, from methanolic shoot extract.¹⁷⁰ A follow up study by the same research group resulted the isolation of two long chain compounds from the methanolic shoot extract, 27-cyclohexylheptacosan-7-ol and 16-hydroxy-26-methylheptacosan-2-one.¹⁷¹ Two new saponins were isolated and characterized from the methanolic extract of aerial parts (Figure 7).¹⁷² Further analysis of the same extract, in a follow up study by the same group, resulted the isolation and characterization of three new saponins and two known compounds that were reported as new, but actually they are not (for example, Ecdysterone).¹⁷³ New cyclic chain fatty ether (interestingly, authors classified it as cyclic chain fatty acid) was isolated from the seed petroleum ether extract (Figure 7).¹⁷⁴ Even though new compounds were not reported in this study, an interesting and new method was developed for the determination of oleanolic acid in this plant.¹⁷⁵ New derivative (glucoside) of oleanolic acid was isolated and characterized.¹⁷⁶ Even though new compounds were not reported in this publication, it is very detailed, presents structures of active natural products and their GCMS chromatograms. This publication can be very useful.¹⁷⁷ Leave methanolic extract was analyzed by column liquid chromatography, and a new compound, aromadendrene-15-*o*-yl ferulate was isolated and characterized (Figure 7).¹⁷⁸ Leaves were extracted and fractionized with several solvents. Fractions were analyzed resulting a new sesquiterpene (Figure 7) and other known compounds.¹⁷⁹ New 6-prenyl apigenin (Figure 7) was isolated and characterized from methanolic extract of leaves. This compound is candidate for developing novel monoamine oxidase-A inhibitor.¹⁸⁰




Figure 7. Active natural products isolated from *Aerva javanica*.

Table 3. Biological, medicinal and other properties of *Aerva javanica*.

Activity/Property	Major Findings/Reference
Analgesic and pain related activities	Aqueous extract of leaves had antinociceptive (acetic acid induced writhing test, hot plate injury and tail immersion method) and antipyretic (Brewer's yeast induced test) activities. ¹⁸¹
Antibacterial, antifungal, antiviral and related activities	Methanolic, ethyl acetated and petroleum ether leaf extracts were prepared and tested for activity against several species of bacteria and fungi. Methanolic extract was highly active, ethyl acetate extract had moderate activity and the non-polar extract was not active. Plant name is mistakenly written <i>Earvajavanica</i> instead of <i>Aerva</i> . ¹⁸² Whole plant was separately extracted with <i>n</i> -hexane, chloroform, ethyl acetate, methanol and water. Extracts were chromatographed and known active natural products were isolated. All extracts and natural products had weak to moderate antibacterial activity. Some of these compounds were isolated from this plant for the first

Anticancer and related activities	<p>time.^{183,184} Aerial parts were separately extracted with <i>n</i>-hexane, ethyl acetate, dichloromethane and methanol. All extracts were active against 5 bacteria species, where DCM extract was most active.¹⁸⁵ Whole plant was extracted with chloroform, ethanol and methanol. Extracts were tested against several bacteria species, and methanolic extract was most active.¹⁸⁶ Dried and fresh leaves were separately extracted with methanol, chloroform, petroleum ether, acetone or water. Each extract was tested against 4 bacteria species, and methanolic extract was most active.¹⁸⁷ Whole plant was extracted with acetone, ethanol, methanol and water, and essential oil was also prepared. All products were tested against several species of bacteria and fungi. Essential oil was most active.¹⁸⁸ Whole plant was extracted with 80 % aqueous ethanol, and extract was active against 6 bacteria species.¹⁸⁹</p>
Antidiabetic Antioxidant	<p>Dried and fresh leaves were separately extracted with methanol, chloroform, petroleum ether, acetone or water. Each extract was tested against prostate and breast cancer cells, and methanolic extract was most active.¹⁸⁷ Whole plant was extracted with acetone, ethanol, methanol and water and essential oil was also prepared. All products were tested against human red blood cells. Ethanolic extract was the most active one.¹⁸⁸ Whole plant was extracted sequentially with <i>n</i>-hexane, chloroform and 80 % aqueous methanol. Extracts were tested against two human breast cancer cell lines (MCF7 and MDA-MB-231). Hydro-methanolic extract was most active.¹⁹⁰ Leaves were successively extracted with <i>n</i>-hexane, chloroform, ethylacetate, acetone, and methanol. Extracts were attested against human breast cancer cell line MCF7. All extracts had significant activities, where methanolic extract was most active.¹⁹¹</p>
Digestive system protection, hepatoprotection, nephroprotection, lung protection	<p>Leaf ethanolic extract had antihyperglycemic effect on alloxan-induced diabetes in mice.¹⁹² Whole plant was extracted with acetone, ethanol, methanol and water, also, essential oil was prepared. All products were tested for antioxidant capacity (DPPH). Methanolic extract was most active.¹⁸⁸ Whole plant was extracted with 80% aqueous ethanol, and extract had high antioxidant capacity (DPPH).¹⁸⁹ Flowers were extracted with 50% aqueous ethanol, and extract was successively fractionized with petroleum ether, diethyl ether, ethyl acetate, benzene, acetone, and ethanol. Ethanolic and ethyl acetate fractions were used to test antioxidant (lipid peroxidation inhibition) in rats. Ethyl acetate fraction was more active.¹⁹³ Aerial parts were extracted with 70% aqueous ethanol, and antioxidant capacity (DPPH) of this extract was determined.¹⁹⁸</p>
Enzyme inhibition	<p>Whole plant was extracted with 80% aqueous methanol and extract was defatted with <i>n</i>-hexane. The resulting crude solid was fractionized with water, dichloromethane and ethyl acetate. Further fractionation and chromatography resulted the isolation of three known compounds. Fractions were tested for urease inhibition and found active, indicating anti-ulcer potential.¹⁹⁴ Aerial parts were extracted with 95% aqueous ethanol, and extract had activity against ethanol-induced gastric lesion in rats.¹⁹⁵ Whole plants was extracted sequentially with ethyl acetate and methanol. Extract had antilithiatic activity against ethylene glycol-induced hyperoxaluria in rats.¹⁹⁶ Whole plant aqueous and ethanolic extracts were active against castor oil induced diarrhea in rats.¹⁹⁷ Aerial parts were extracted with 70% aqueous ethanol, and this extract showed hepatoprotective activity against CCl₄-induced toxicity in rats.¹⁹⁸ The information in this publication is confusing. The title is “research paper”, which describes the hepatoprotective activity of whole plant extracts. Practically, it is written as review article.¹⁹⁹</p>
Herbicidal Toxicity	<p>Flowers methanolic extract showed clear activity of the following enzymes inhibition: acetylcholinesterase (AChE), butyrylcholinesterase (BChE) and lipoxygenase (LOX).²⁰⁰ Aerial parts aqueous extract had herbicidal activity against <i>Cyperus rotundus</i>.²⁰¹ Aqueous infusions were prepared from aerial parts of female and male plants. These infusions were orally administered to female and male rats. At the dose of 500 mg kg⁻¹ for several weeks, female rats suffered hypochromic anemia.²⁰²</p>
Chemical composition	<p>Flowers methanolic extract was analyzed and four new compounds (Ecdysteroids, Figure 8) were isolated and characterized.²⁰⁰ Chromatographic purification of ethyl acetate soluble fraction of the methanolic extract of the flower yielded three new acylated flavone glycosides (Figure 8).²⁰³ A follow up study of the same extract by the same group obtained Aeryfuraniside (Figure 8).²⁰⁴ The following publications reported general chemical composition, and did not report new compounds: essential oil (stems and leaves) composition,²⁰⁵ essential oil (seeds) composition, with comparison between methods of extraction,²⁰⁶ composition of whole plant ethanolic extract,²⁰⁷ very detailed and very useful analysis of various extracts of all parts, including mineral composition,²⁰⁸ analysis of essential oils compositions resulting from transmission of phytoplasmata.²⁰⁹</p>




Figure 8. Active natural products isolated from *Aerva javanica*.

Table 4. Biological, medicinal and other properties of *Alternanthera pungens*.

Activity/Property	Major Findings/Reference
Analgesic and pain related activities	Whole plant ethanolic extract was prepared and tested for analgesic activity in mice by hot plate test and acetic acid-induced writhing test. ²¹⁰
Antibacterial, antifungal, antiviral and related activities	Aerial parts were separately extracted with water, acetone, ethanol and petroleum ether. All extracts were tested for antibacterial activity and ethanolic extract was most active. ²¹¹
Antioxidant, anti-inflammatory and related activities	Aerial parts were separately extracted with water, acetone, ethanol and petroleum ether. Antioxidant capacity of all extracts were determined by DPPH method. ²¹¹ Whole plant ethanolic extract was prepared and tested for antioxidant (DPPH) and anti-inflammatory (albumin denaturation technique) activities in mice. ²¹⁰ Even though a detailed NMR work is reported, but the analyzed compounds are previously known phenolics. These were tested for antioxidant capacity (Trolox test). The report includes a confusing part (Extraction and isolation), where it states that aerial parts were extracted with “boiling ethanol”, and when extract was concentrated, aqueous concentrate was obtained. ²¹²
Diuretic, gastroprotective	Whole plant ethanolic extract have diuretic effect in rats, like the effect of Furosemide. ²¹³ Whole plant aqueous extract acted as gastro-stimulant in rats. ²¹⁴
Antiparasitic	Whole plant was successively extracted with n-hexane, chloroform, ethyl acetate and methanol. All extracts were tested for antiparasitic (<i>Pheritima posthuma</i>) activity, and all extracts were active except methanolic. ²¹⁵

Table 5. Biological, medicinal and other properties of *Alternanthera sessilis*.

Activity/Property	Major Findings/Reference
Allelopathy, growth promotion	Aqueous extract of leaves was tested for allelopathic activity against seed germination of Rice (<i>Oryza sativa</i>). The allelopathic effect increased with concentration and when this reached 5 %, the germination of seeds was zero. Active phenolics in this extract are chlorogenic acid, gallic acid and vanillic acid. ²¹⁶ Leaves were successively extracted with petroleum ether, acetone and ethanol. The ethanolic extract was fed to <i>Macrobrachium rosenbergii</i> (Prawn) resulting growth promotion in these animals. An excellent presentation of active natural products is provided. ²¹⁷
Analgesic and pain related activities	Aerial parts methanolic extract had analgesic activity in abnormal writhing test in mice. ²¹⁸ Aerial parts were extracted with 90 % aqueous ethanol. Extract had analgesic activity in mice, as prove by two tests: acetic acid induced writhing test and hot-plate test. ²¹⁹ Whole plant was defatted with n-hexane and extracted with ethanol, and this extract had antipyretic (yeast-induced) activity. ²²⁷
Antiallergy, skin protection, wound healing	Aerial parts were extracted with 95 % aqueous ethanol and the extract had inhibitory effect in IgE-mediated allergic response in RBL-2H3 cells. ²²⁰ Leaves were successively extracted with petroleum ether, chloroform, acetone, methanol and water. All extracts (methanolic highest) had wound healing activity (excision, incision) in rats. ²²⁴ Stems were extracted with 90 % aqueous ethanol and extract had wound (scratch) activity. ²²⁵

Antibacterial, antifungal	Aerial parts were extracted with 95 % aqueous ethanol and extract was active against 6 bacteria and 3 fungi species. ²²² Leaves were separately extracted with water, ethanol and acetone. Extracts and leaves juice had activity against 3 bacteria species. ²²³ Leaves were successively extracted with petroleum ether, chloroform, acetone, methanol and water. All extracts (chloroform highest) had weak to moderate activity against 4 bacteria species. ²²⁴ Leaves were extracted with methanol and petroleum ether, and both extracts were active against 3 bacteria and 2 fungi species. ²²⁶
Anticancer	Aerial parts were extracted with ethanol, and extract had activity against human colon cancer cells (HT-29, 3T3). ²²⁸
Antidiabetic, anti-obesity and related activities	Aerial parts methanolic extract had antidiabetic activity in glucose loaded mice. ²¹⁸ Leaves were extracted with methanol and petroleum ether, and both extracts had antidiabetic activity (α -amylase inhibition). ²²⁶ Aerial parts were extracted with 95% aqueous ethanol, and extract was fractionized by <i>n</i> -hexane. This fraction was extracted with ethyl acetate, and the obtained extract antidiabetic activity in obese rats that had STZ-induced diabetes. ²³⁰ Leaves and callus were extracted with methanol, and extract was fractionized with several polar solvents. Extracts and fractions were tested for α -glucosidase inhibition. All showed notable activity, but callus extract was more active. ²³¹
Antioxidant, anti-inflammatory and related activities	Stems were extracted with 90 % aqueous ethanol, and its antioxidant capacity (DPPH) was determined. ²²⁵ Leaves were extracted with methanol and petroleum ether, and antioxidant capacity (DPPH) of both extracts was determined. Anti-inflammatory activity was proved by egg albumin test. ²²⁶ Leaves and callus were extracted with methanol, and extract was fractionized with several polar solvents. Extracts and fractions were tested antioxidant (DPPH) capacities. ²³¹ Antioxidant capacities of ethanolic extracts of two cultivars of this plant, by three methods: DPPH, Trolox and FRAP. ²³² Essential oil was extracted from aerial parts and its antioxidant capacity was determined by DPPH method. ²³³ Leaves were extracted with methanol and extract was fractionized with dichloromethane and <i>n</i> -hexane. Extract and fractions were tested for antioxidant capacity (DPPH). ²³⁴
Brain related activities	Aerial parts were extracted with 90 % aqueous ethanol. Extract had stimulant activity in mice, as prove by three tests: Pentobarbitone-induced sleeping time test, open field test and hole cross test. ²¹⁹ Methanolic extract of leaves had antidepressant (tail suspension and forced swimming tests) activity in mice. ²²⁹
Cardiovascular system	Whole plant was extracted with 70 % aqueous ethanol and the extract was fractionized with water and dichloromethane. Extract and fractions had hypotensive effect in rabbits. Extract had higher activity than fractions ²²¹
Digestive system protection, hepatoprotection, nephroprotection, lung protection	Whole plant was extracted with 70 % aqueous ethanol and the extract was fractionized with water and dichloromethane. Extract and fractions had anti-asthmatic and anti-diarrheal effect in rabbits. Extract had higher activity than fractions ²²¹ Aqueous extract of leaves had strong activity against castor oil-induced diarrhea in mice. ⁵³ Ethanolic extract of aerial parts was active against damages in mice caused by three hepatotoxines: carbon tetrachloride acetaminophen and galactosamine. ²³⁵ Whole plant methanolic extract was active against damages in mice caused by carbon tetrachloride. ²³⁶
Metals related activities	Aqueous extract of leaves was used as a reductant in the production of silver nanoparticles (AgNP's) from AgNO ₃ solution. These AgNPs have antibacterial and antioxidant activities. ²³⁷ Young plants found active chromium (Cr ₂ O ₇ ⁻²) phytoextract in, ²³⁸ domestic sewage, ²⁴⁰ PCB-contaminated soil (with an addition of Bentonite), ²⁴¹ ions (Cd ⁺² >Cr ⁺³ >Pb ⁺²). Antioxidant enzymes concentrations increased in different parts of the plant after absorbing the metal ions. ²⁴²
Nutrition, toxicity	Leaves were analyzed for nutrients and antinutrients. Authors concluded that its safe and recommended to eat it. ²⁴³
Chemical composition	Leaves were extracted with petroleum ether, resulting the isolation and characterization of a new natural product that authors symbolized with PDHC (Figure 9). Detailed structures of known compounds are also presented. ²²⁶ New saponin from methanolic extract of leaves(Figure 9). ²⁴⁴ Analysis of leaves ethanolic extract afforded new ionone (Figure 9). ²⁴⁵ Novel (plant defense) peptides were isolated from leaves aqueous extract. ²⁴⁶ The following references reported chemical compositions (some are very detailed) of known natural products. ^{233,234,247-250} Special method for detection and quantification of gallic acid in the plant. ²⁵¹ Very extensive work was invested in this research, especially in spectroscopic characterization of the reported compound. Authors did not claim novelty, but the presentation of the work ("isolation and characterization") can mislead readers. Anyhow, this compound was isolated and characterized on year earlier. ^{252,253} Detailed work, including some useful chromatograms. ²⁵⁴




Figure 9. Active natural products isolated from *Alternanthera sessilis*.

Table 6. Biological, medicinal and other properties of *Amaranthus cruentus*.

Activity/Property	Major Findings/Reference
Allelopathy	Tomato (<i>Lycopersicon esculentum</i>) was planted in soil that was brought from a field where <i>A. cruentus</i> was growing, showed weak germination and growing compared to control. ²⁵⁷ Allelopathy of <i>A. cruentus</i> was clearly temperature-dependent, when its influence was tested by growing factors of several vegetables that were planted with <i>A. cruentus</i> . Influence was not uniform for all tested vegetables. ²⁵⁸ Aqueous extract of seeds had allelopathic effect on seedlings of garden cress (<i>Lepidium sativum</i>). ²⁶⁹
Antihyperlipidemic	Seeds were defatted with <i>n</i> -hexane and hydrolyzed with pH=11 solution. Proteins contained in this hydrolysate, had inhibition activity of HMG-CoA reductase, suggesting possible hypocholesterolemic effect. ²⁶⁵
Antioxidant	Seeds were extracted with methanol and the antioxidant capacity of this extract was determined by three methods (FRAP, DPPH and ABTS). ²⁵⁹ High fructose diet resulted in oxidative stress in rats. When co-administered with seeds, indications of decreasing malondialdehyde in plasma and increasing activities of antioxidant enzymes were recorded. ²⁶⁰ Aerial parts were extracted with 20 % aqueous acetone and the extract was fractionized with water. Antioxidant capacities of extract and fraction were determined by FRAP and DPPH methods. ²⁶¹
Blood related activities	Seed were separately extracted with <i>n</i> -hexane, ethanol and 70 % aqueous ethanol. All extracts had strong haemagglutinating properties. ²⁶² Ethanolic extract of leaves had protective activity (aided in restoring the levels of red blood cells, white blood cells and hemoglobin) against phenylhydrazine-induced toxicity in rats. ²⁶³ Ethanolic leaves extract had hematopoietic effect on cyclophosphamide-induced toxicity in rats. ²⁶⁴
Enzyme inhibition	Aerial parts were extracted with 20 % aqueous acetone, and extract was fractionized with water. Extract and fraction had xanthine oxidase inhibitory activity. ²⁶¹
Metals related activities	Analysis of seed samples that were collected from wild growing plants (USA), indicated high concentrations (accumulation) of toxic metal ions: As^{+3} , Cr^{+3} and Pb^{+2} . ²⁶⁶
Nutrition, toxicity	Major nutritional components of seeds flour and its protein concentrate were determined as phenolics, mineral, fatty acids and carbohydrates. ²⁶⁷ Dichloromethane extracts of seeds was found rich in saponin and toxic for hamsters. ²⁶⁸
Chemical composition	Young methanolic extract of leaves was analyzed resulting isolation and characterization of (<i>E</i>)-caffeoylisocitric acid (Figure 10). Its enzymatic synthesis is presented. ²⁷⁰ Novel pectinic polysaccharides were isolated and characterized from the methanolic extract of aerial parts. ²⁷¹ The following publications presented interested information about chemical composition of the plant but did not report new natural products, HPLC method for isolation of betalains (Figure 6), ²⁷² proteins and fibers, ²⁷³ HPLC method for analysis of phenolic acids and flavonoids in the seeds ²⁷⁴ and isolation of betacyanins (Figure 6) from the flowers. ²⁷⁵




Figure 10. Active natural products isolated from *Amaranthus cruentus*.

Table 7. Biological, medicinal and other properties of *Amaranthus graecizans*.

Activity/Property	Major Findings/Reference
Analgesic	Whole plant was extracted with 90 % aqueous methanol. This extract had analgesic activity tested by three methods (hot plate, tail immersion, acetic acid induced writhing) in mice. ²⁷⁶
Antibacterial	Whole plant was extracted with chloroform, ethanol and methanol. Extracts were tested against several bacteria species and the methanolic extract was most active. ¹⁸⁶
Antioxidant, anti-inflammatory	Whole plant was extracted with 90 % aqueous methanol. This extract had anti-inflammatory activity tested by carrageenan-induced paw edema method. ²⁷⁶ Whole plant was extracted with methanol and extract was fractionized with water, <i>n</i> -hexane, chloroform, ethyl acetate and <i>n</i> -butanol. The antioxidant capacities of extract and fractions were determined by four methods: DPPH, FRAP, FTC and phosphomolybdenum. General chemical composition was determined in this study. ²⁷⁷
Hepatoprotection	Whole plant was extracted with 90 % aqueous methanol, and extract was active against isoniazid and rifampicin-induced hepatotoxicity. ²⁷⁸
Enzyme inhibition	Whole plant was extracted with 90 % aqueous methanol, and extract had cholinesterase inhibition activity. ²⁷⁶
Chemical composition	Aerial parts were extracted with 10 % aqueous ethanol and the general chemical composition of extract was determined. ²⁷⁹

Table 8. Biological, medicinal and other properties of *Amaranthus retroflexus*.

Activity/Property	Major Findings/Reference
Allelopathy	Aqueous extract of seeds had allelopathic effect on seedlings of garden cress (<i>Lepidium sativum</i>). ²⁶⁹ Leaves, stems and roots were collectively extracted with water, and this extract had allelopathic effect on Barley. ²⁸⁷ Whole plant aqueous extract had allelopathic effect on cucumber, alfalfa, common bean and bread wheat. ²⁸⁸ A follow up study by the same research group that tested the allelopathic effects of aqueous extracts on cucumber and wheat, but in this case, plant organs were extracted separately. ²⁸⁹ Plant powder had allelopathic effect on germination of maize. ²⁹⁰
Allergy	Allergens were isolated from aqueous extract of pollen, and they were tested in human patients. ²⁹¹ An allergen named Polcalcin was isolated from the aqueous extract of pollen. It can be produced in <i>E. coli</i> . ²⁹²
Antibacterial	Leaves and flowers were combinedly extracted with 70 % aqueous ethanol and extract was active against 7 bacteria species. ²⁹³
Antioxidant, anti-inflammatory and related activities	Leaves and flowers were combinedly extracted with 70 % aqueous ethanol and the antioxidant capacity of this extract was tested by DPPH method. ²⁹³ Leaves were extracted with 70 % aqueous ethanol, and this extract had high antioxidant capacity (DPPH). ²⁹⁴ Leaves were extracted with 10% aqueous methanol and 10 known nerolidol derivatives (structures are presented) were isolated and tested for their antioxidant capacities (DPPH). ²⁹⁵
Metals related activities	Plant was found good phytoremediator of soil contaminated with Cr ⁺³ , Cd ⁺² , Cu ⁺² and Ni ⁺² . ²⁹⁶ Lead and zinc ions were successfully removed from wastewater, especially after treating the plants with growth promoter, citric acid. ²⁹⁷ Aqueous extract of leaves was used to reduce silver ions (AgNO ₃ solution), to prepare silver nanoparticle, AgNPs, which had antifungal activity. ²⁹⁸
Toxicity	Leaves were extracted with methanol-acetone-water (ratio is not indicated) and this extract was toxic only to renal cell culture. ²⁹⁹ Following cases of poisoning cases (renal) of livestock in Australia, aerial parts methanolic extract was prepared and analyzed. The major toxic ingredients were mainly quaternary ammonium and polyhydroxylated amines. New compounds were not reported. ³⁰⁰
Chemical composition	Leaves were extracted with 10 % aqueous methanol and the extract was chromatographed, resulting the isolation and characterization of 7 new compounds that were identified as nerolidol derivatives (Figure 11). ³⁰¹ A follow up study by the same research group afforded 4 new compounds from the same family, nerolidol glucoside derivatives. ³⁰² An outstanding and comprehensive analysis of the chemical composition of the plant including a new natural product, a sphingolipid (Figure 11), has been reported. ³⁰³ Detailed composition of volatile compounds was analyzed from gases released by the plant itself, or from whole plant aqueous extract, that was subjected to vacuum distillation. New compounds were not reported. ³⁰⁴ Comprehensive analysis of water-soluble compounds, with special focus on carbohydrates, organic acids and phenolics was reported. ³⁰⁵

Brain related activities	essential oil. Both products were analyzed for chemical composition, and their antioxidant capacities were determined (DPPH, H ₂ O ₂ and enzymatic assay). Detailed presentation of chemical composition is provided. ³²² Methanolic extract of seeds had high antioxidant capacity (DPPH, H ₂ O ₂). ³³³ Antioxidant capacity of leaves was determined by two methods (DPPH, ABTS ⁺). General chemical composition has been presented. ³³⁵
Cardiovascular system related activities	Leaves were defatted with petroleum ether and extracted with 95% aqueous ethanol. Extract had antianxiety activity which was induced in rats by several methods. ³²³ An earlier study by the same group, where whole plant was extracted with methanol and extract was tested for antidepressant activity, that was induced in rats by several methods. In both studies, acute toxicity and general chemical composition were presented. ³²⁴
Digestive system protection, hepatoprotection, nephroprotection, lung protection	Leaves were extracted with 70 % aqueous methanol. When the extract was administered to rats, adverse effects of fat increasing in the plasma of rats were recorded. But when administered with ascorbic acid, a synergetic effect was observed. ³²⁵ Whole plant aqueous extract had toxicity only in high doses in rats, but its cardioprotective effect was weak and temporary. ³²⁶
Insecticidal	Leaves were extracted with methanol and extract had gastroprotective (gastric erosion assay) and antidiarrheal (castor oil-induced diarrhea) activities in mice. ³¹⁸ Whole plant was extracted with 50 % aqueous ethanol. This extract had activity against castor oil-induced diarrhea and ethanol-induced gastric ulcer, in mice. ³²⁷ Leaves were extracted with petroleum ether, chloroform, ethanol and water. All extracts had anti-ulcer (pylorus ligation) activity in rats, where ethanolic extract was more active than others. ³²⁸ Whole plant was extracted with 70 % aqueous methanol and extract was fractionized by ethyl acetate and water. Three animal species were administered with carbachol and treated with extract and fractions. These had bronchodilator and gut modulatory activities, where extract was more active than fractions. ³²⁹ Whole plant aqueous extract had diuretic activity in rats. ^{330,331} Whole plants were extracted with 50 % aqueous ethanol and this extract had activity against d-galactosamine/LPS-induced hepatic failure. ³³² Seeds methanolic extract had activity against hepatotoxicity associated with deltamethrin in rats. ³³³
Metals related activities	Whole plant was extracted with 70% aqueous ethanol. This extract combined with the same extract of <i>Andrographis paniculate</i> were active against <i>Plasmodium berghei</i> -infected mice. ³³⁶
Nutrition, toxicity	Leaves ethanolic extract was used to reduce Au ⁺³ ions (H ₄ AuCl ₃) to prepare gold nanoparticles, AuNP's. ³³⁷ Aqueous leaves extract was used to prepare FeO-NPs from FeCl ₃ solution. ³³⁸
Reproductive system, sex	Aerial parts were successively extracted with <i>n</i> -hexane, chloroform, ethanol and water. Ethanolic extract showed toxicity in brine shrimp lethality test. Partial general chemical composition is presented. Interestingly, it is reported that none of these extracts had antibacterial activity. ³¹³ Leaves were fed to <i>Clarias gariepinus</i> (African catfish species) as a potential protein source. ³³⁹ Consumption of aqueous extract of leaves by postpartum mothers, increased prolactin and breast milk. ³⁴⁰ Single report of poisoning as a result of consumption of the plant by sheep in Brazil. Various adverse physical damages were observed, but the major adverse effect was renal insufficiency. ³⁴¹
Chemical composition	Roots were extracted with acetone, and extract was co-administered to male rats with acetone seeds extract of <i>Dolichos biflorus</i> . Various adverse effects of on sex organ and enzymes were recorded. ³⁴² Methanolic extract of leaves was co-administered with <i>Abrus precatorius</i> extract (70 % aqueous methanol) to male rats, resulting in increasing of infertility. ³⁴³ Whole plant ethanolic extract was orally administered to female rats, resulting in contraceptive and abortifacient activities. ³⁴⁴
	Analysis of whole plant methanolic extract (<i>n</i> -butanol fraction) afforded new natural product, Spinoside (Figure 12). ³⁴⁵ The following reports presented general chemical composition or known compounds, but they also presented uniqueness or special interest. Betacyanins and phenolics, ³⁴⁶ Very detailed composition with (or without) tables, ³⁴⁷⁻³⁵⁴ isolation of α -spinasterol from stem extract, ³⁵⁵ identification and quantification of quercetin in leaves extract, ³⁵⁶ isolation of mucilages from leaves ³⁵⁷ have been reported




Figure 12. Active natural products isolated from *Amaranthus spinosus*.

Table 10. Biological, medicinal and other properties of *Amaranthus viridis*.

Activity/Property	Major Findings/Reference
Analgesic and pain related activities	Methanolic whole plant extract had analgesic activity in three tests (mentioned above) in mice. ³³⁴ Whole plant methanolic extract of whole plant was had antinociceptive activity tested by acetic acid-induced writhing test, hot plate test and tail immersion test in mice. Antipyretic activity was tested by yeast-induced pyrexia method in rats. ³⁶⁵ Whole plant methanolic extract had antinociceptive effect against acetic acid-induced gastric pain in mice. ³⁶⁷
Wound healing	Methanolic extract of leaves had wound healing activity (excision, incision, dead space) in diabetic rats. ³⁶⁶
Antibacterial, antifungal	Whole plant was extracted with chloroform, ethanol and methanol. Extracts were tested against several bacteria species, and methanolic extract was most active. ¹⁸⁶ Aerial parts ethanolic extract had activity against 5 bacteria and 5 fungi species. ³⁵⁸ Seeds (in the title it is indicated as leaves) methanolic extract was prepared and tested against 5 bacteria species. A GC-MS analysis is presented. ³⁵⁹ Extract of leaves alkaloids was found active against several bacteria species. ³⁶⁰ Seeds, leaves or aerial parts, were separately extracted with organic solvents, and extracts were active against some bacteria species. ³⁶¹⁻³⁶³ Leaves were separately extracted with dichloromethane, ethyl acetate and ethanol. All extracts were active against two fungi species. ³⁶⁴ Analysis of seed extract (70 % aqueous methanol) afforded a new lectin that was active against two fungi species. ³⁶⁸
Anticancer and related activities	Analysis of seed s extract (70 % aqueous methanol) afforded a new lectin that was active against HB98 and P388D1 murine cancer cell lines. ³⁶⁸ Leaves and stems were combinedly extracted with 50 % aqueous ethanol. Extract had activity against 3 human leukemic cell lines. General chemical composition is presented. ³⁶⁹ Leaves were extracted with <i>n</i> -hexane, chloroform, methanol and water. Methanolic extract had antigenotoxic activity. ³⁷⁷
Antidiabetic and related activities	Whole plant methanolic extract had antihyperglycemic effect in glucose-loaded mice. ³⁶⁷ Whole plant was extracted with 75 % aqueous ethanol and the extract had activity against STZ-induced diabetes in rats. ³⁷⁰ Whole plant or methanolic extract of leaves had activity against alloxan/STZ-induced diabetes in rats, and antihyperlipidemic activity. ^{371,372}
Antioxidant, anti-inflammatory and related activities	Antioxidant capacity of methanolic extract of leaves was determined by two methods (DPPH and ABTS ⁺). General chemical composition is presented. ³³⁵ Various parts, were separately extracted with organic solvents, and extracts were tested for antioxidant capacities (DPPH). ^{361-363,374,375} Leaves and stems were combinedly extracted with 50 % aqueous ethanol, and the antioxidant (DPPH) capacity of extract was determined. ³⁶⁹ Whole plant methanolic extract had activity against lipid peroxidation in rats. ³⁷¹ Leaves were analyzed for nutrient and mineral contents, extracted with methanol, and the antioxidant capacity of this extract was determined (DPPH). ³⁷³ Alkaloid-rich leaves extract had activity against H ₂ O ₂ damage in human erythrocytes. ³⁷⁶ Leaves were extracted with <i>n</i> -hexane, chloroform, methanol and water. Methanolic extract had highest antioxidant capacity (5 methods). ³⁷⁷ Leaves, stems and seeds were extracted with methanol and the antioxidant capacity of each extract was determined by 5 methods. Anti-inflammatory activity was tested by hyaluronidase and lipoxygenase inhibition. ³⁷⁸
Neuroprotection	Aqueous extract of leaves had activity against cyclophosphamide-induced neuro-endocrine dysfunction toxicity in rats. ³⁸⁴
Cardiovascular system related activities	Methanolic extract of leaves had hypocholesterolemic and anti-atherosclerotic effects in rabbits. Detailed HPLC analysis of extract is presented. ³⁷⁹ Leaves or whole plant methanolic extract had protective effect against isoproterenol toxicity in rats. ^{380,381}
Hepatoprotection, nephroprotection	Aqueous extract of roots had activity against ethylene glycol-induced urolithiasis in rats. ³⁸² Whole plant methanolic extract had activity against paracetamol liver toxicity in rats. Antioxidant biomarkers concentrations are reported. ³⁸³
Enzyme inhibition	Leaves were extracted with <i>n</i> -hexane, chloroform, methanol and water. Methanolic extract had tyrosinase inhibition activity. ³⁷⁷ Leaves, stems and seeds were extracted with methanol and each extract had xanthine oxidase inhibition activity. ³⁷⁸ Phenolics rich aqueous extract of leaves, from cultivated plants, had trypsin inhibition activity. ³⁸⁵
Metals related activities	Supplementation of FYM (farmyard manure) to cultivated plants, enhanced their ability as Cd ⁺² accumulator. ³⁸⁹ Uptake of Ni ⁺² inhibits growth of cultivated plants. ³⁹⁰ Plants have high tolerance to metal (Cd ⁺² , Cr ⁺³) soil contaminations. ³⁹¹ Risk assessment of metals (Cd ⁺² , Ni ⁺² , Pb ⁺²) accumulation in plants, that grow in contaminated soil and can be consumed by humans as food, been reported. ³⁹² Aqueous extract of leaves/twigs reduced Ag ⁺ (AgNO ₃) for the preparation of silver nanoparticles, AgNPs, which had antibacterial activity. ^{393,394}
Nutrition, toxicity	Starch from this plant was modified (functional groups) for the purpose of using it as fat replacement. ^{395,396} Aqueous extract of leaves was not toxic to rats. ³⁹⁷

Reproductive system	Aqueous extract of leaves had activity against cyclophosphamide-induced reproductive system toxicity in rats. ³⁸⁴ Roots powder had notable anti-menorrhagia in human females aged between 16 - 45 years suffering from excessive and or irregular vaginal bleeding. ³⁸⁶ Aqueous extract of roots had abortifacient effect in female rats. ³⁸⁷ Methanolic extract of leaves had estrogenic activity in female rats. ³⁸⁸
Chemical composition	In addition to general chemical composition that was presented in previous cited publications, two more interesting articles did the same. ^{398,399} First isolation and characterization of Amasterol, an ecdysone precursor and a growth inhibitor (Figure 13). ⁴⁰⁰




Figure 13. Active natural products isolated from *Amaranthus viridis*.

One of these notable attempts used the combination of Murashige and Skoog medium and indole acetic acid, as growth promoters. This research was performed for the aim of enhancement of active compounds production.⁴⁰⁶ Another interesting cultivation used this plant to protect Cowpea (*Vigna unguiculata*) and Mung bean (*Vigna radiata*) crops from root rot fungi.⁴⁰⁷

Alternanthera pungens was very limitedly studied and most of its chemical and medicinal properties were not published (Table 4). Even its general chemical composition is still unknown and to the best of our knowledge, isolation of novel natural products from this plant has never been published.

On the contrary, the second representative of the same genus, *A. sessilis* has been well investigated and most of its properties are known and so is its chemical composition (Table 5, Figure 9). Some interesting natural products were isolated from that plant. But some of these cited reports seem confusing when the information that they include are compared. Lalitha Sree *et al.* state that leaves of this plant are highly nutritional, and it is safe to consume them.²⁴³ Singh and her colleagues go even beyond that: they consider it a famine food plant and report its cultivation under different growth promoting conditions.⁴⁰⁸ But Gayathri *et al.* reported three years earlier that aqueous extract of aerial parts was found hepatotoxic in mice.⁴⁰⁹ Finally, it is important to include into toxicity considerations not only the location of the plant harvest and soil contamination levels, but seasonal variations in the possibly toxic contents of the plant also.⁴¹⁰

All other 11 plants of the *Amaranthaceae* family, that grow wild in Israel and Palestine, are of the genus *Amaranthus*. Many published articles present various aspects and properties of this genus, and we presented some of these publications earlier in this article. But some of these reports may need further examination.

One of these outstanding reports was published by Noori and her colleagues.⁴¹¹ They analyzed aerial parts (excluding stem) of 7 *Amaranthus* (5 of them are included in this review article) species in Iran, for flavonoid content. Even though they did not report novel compounds, this work is very detailed, comprehensive and can be very useful for interested researchers and scholars. An early review article by Suma *et al.* focuses on the allelopathy of some *Amaranthus* species.⁴¹² It's a useful publication, but it needs updating and despite including a section of allelochemicals, no structures are presented, and it is very brief. Finally, an important work was published by Srivastava, about the nutritional potential of some cultivated wild *Amaranthus* plants.⁴¹³ The importance of this work emerges from the reported ability to cultivate these plants and to partially control their nutritional values.

The cultivation of *Amaranthus cruentus* drew notable attention from researchers and many works were published about this issue. Marin and his colleagues studied the influence of various cultivation parameters on seeds production due to their high nutritional value.⁴¹⁴ Kornarzynski and his colleagues reported growth promotion of this plant by electromagnetic stimulation to enhance the production of pigments (chlorophylls and carotenoids) in leaves.⁴¹⁵ Hlinkova and her colleagues grew this plant for two consecutive years and discovered high content of fatty acids (seeds) with notable unsaturation index (UI).⁴¹⁶

As we mentioned earlier, *Amaranthus palmeri* also has not been investigated much. Its general chemical composition and its allelopathic effect on some domestic plants are partially known. Since in general it is considered a weed, its remarkable resistance to herbicides (see Figure 4 and ref. 58), like all *Amaranthus* plants, was studied.⁴¹⁷ Authors report that the plant has multiple resistance for 4 herbicides.

Amaranthus retroflexus is one of the most widespread *Amaranthus* plants in the reviewed region (Figure 1). It has been sufficiently studied (Table 8, Figure 11) and a remarkable number of very interesting, new natural products were isolated from this plant. In addition to the cited literature above, a special report about its chemical composition, was published by Woo.⁴¹⁸ This report indicated the very early scientific interest in this plant and its usefulness lies in the good graphical presentations of the chemical contents. The plant is mostly considered a weed and its resistance to herbicides is well known. Francischini *et al.* studied the resistance of this plant to acetolactate-synthase inhibitors, such as trifloxysulfuron-sodium and pyriathiobac-sodium (Figure 14),⁴¹⁹ and Wang *et al.* studied its resistance to thifensulfuronmethyl and fomesafen (Figure 14).⁴²⁰




Figure 14. Herbicides that *A. retroflexus* showed resistance to (ref. 419, 420).

The toxicity of this plant goes way beyond allelopathy. An interesting study that was published by Dinu and her colleagues showed that the 50 % aqueous ethanol extract of leaves had toxic activity against mice, *Daphnia magna* (plankton) and *Triticum aestivum* (common wheat).⁴²¹ So, in addition to attempts of combating this plant with herbicides, some controlled cultivation trials were published, that aim to find the best method of preventing seeds germination,⁴²² including using growth promoters.⁴²³

The case of *Amaranthus rudis* is not quite understandable in terms of the lack of published studies about it. Its relatively not very common in the reviewed region as it grows only in the Northern parts of it but worldwide and especially in North America it is very widespread weed. This is probably the reason that the only publication, we found about it, is a study of its control by herbicides, in maize fields.⁴²⁴

Amaranthus spinosus is the most studied plant of the *Amaranthus* genus so far. Its medicinal properties and chemical composition (Table 9, Figure 12) are well known, especially the biologically active long chain alcohols. Its allelopathic effects are also known,^{306,417} and so like other plants of this genus, several studies were performed about its control as a weed. One of these studies was carried out by Syahri and his colleagues where they tried a biocontrol method of controlling *A. spinosus* using Mango leaf extract.⁴²⁵ It should be noticed that the title of this publication stated that a “bioactive compound” was used, but actually its an extract not a pure compound.

Amaranthus viridis was also extensively (properties but very limited chemical composition) studied, but oddly enough, there are no significant reports about is allelopathy (Table 10, Figure 13). In many countries it is an invasive harmful plant to the local domestic and wild vegetation.⁴²⁶ And like other plants of this genus, it has remarkable resistance to herbicides.⁴²⁷ But despite that, in some regions in the world, it is cultivated mainly as a protein source.⁴²⁸

To conclude this part, we will cite two interesting reports related to *Amaranthus viridis*. First is the publication of Ratnasooriya and his colleagues about the lack of diuretic activity of hot water extract of whole plant of *Amaranthus viridis* L. in rats.⁴²⁹ This work does not look obvious since generally such investigations are done to clarify, opposing, previous reports or to contradict unreliable reports. To the best of our knowledge, there are no publications that report diuretic activity of this plant, save in traditional medicine.²⁵

Second, two new interesting natural products have been isolated and characterized, from *Chaetomium globosum*, a fungal endophyte isolated from *A. viridis* leaves (Figure 15).⁴³⁰




Figure 15. Novel compounds from *Chaetomium globosum*, a fungal endophyte isolated from *A. viridis* leaves.

CONCLUSIONS AND FUTURE VISION

(1) Plants of the *Amaranthaceae* plants of Israel and Palestine, possess great properties. They contain unique natural products with special structural sub-units.

(2) Medicinal potential and properties of the plants is far from being fully explored. Most of the unique natural products isolated from these plants have not been studied yet for medicinal activities and their synthetic analogues have not been prepared and studied. Some of the species of this family have either limited studies or not at all.

(3) A very thorough study for the medicinal potential of these plants is needed. Efforts are needed to isolate new natural products from all the plants of this family. It is important to test the medicinal potential of these natural products. It is highly important to prepare synthetic analogues of these natural products and intensify the research of biocontrol of some of the species.

REFERENCES

- ¹Borsch, T., Clemants, S., Mosyakin, S., Biology of the *Amaranthaceae-Chenopodiaceae* alliance, *J. Torrey Bot. Soc.*, **2001**, 128, 234-235. DOI:10.2307/3088715
- ²Fritz, G. J., Adams, K. R., Rice, G. E., Czarzasty, J. L., Evidence for Domesticated Amaranth from a Sedentary Period Hohokam House Floor at Las Canopas, *Kiva*, **2009**, 74, 393-419. DOI: 10.1179/kiv.2009.74.4.002
- ³Rastogi, A., Shukla, S., Amaranth: A New Millennium Crop of Nutraceutical Values, *Crit. Rev. Food Sci. Nutr.*, **2013**, 53, 109-125. DOI: 10.1080/10408398.2010.517876
- ⁴Gheggi, M. S., Williams, V. I., New Data on Food Consumption in Pre-Hispanic Populations from Northwest Argentina (ca. 1000-1550 A.D.):The Contribution of Carbon and Nitrogen Isotopic Composition of Human Bones, *J. Anthropol.*, **2013**, Article 258190. <http://dx.doi.org/10.1155/2013/258190>

- ⁵Olsvig-Whittaker, L., Maeir, A. M., Weiss, E., Frumin, S., Ackermann, O., Kolska Horwitz, L., Ecology of the Past - Late Bronze and Iron Age Landscapes, People and Climate Change in Philistia (the Southern Coastal Plain and Shephelah), Israel, *J. Mediter. Ecol.*, **2015**, *5*, 57-75. <https://www.researchgate.net/publication/283122930>
- ⁶Jimoh, M. O., Afolayan, A. J., Lewu, F. B., Suitability of *Amaranthus* species for alleviating human dietary deficiencies, *S. Afr. J. Bot.*, **2018**, *115*, 65-73. <https://doi.org/10.1016/j.sajb.2018.01.004>
- ⁷Graf, B. L., Rojas-Silva, P., Rojo, L. E., Delatorre-Herrera, J., Baldeon, M. E., Raskin, I., Innovations in Health Value and Functional Food Development of Quinoa (*Chenopodium quinoa* Willd.), *Compr. Rev. Food Sci. Food Saf.*, **2015**, *14*, 431-445. DOI:10.1111/1541-4337.12135
- ⁸Roberts, J. L., Moreau, R., Functional properties of spinach (*Spinacia oleracea* L.) phytochemicals and bioactives, *Food Funct.*, **2016**, *7*, 3337-3353. DOI: 10.1039/c6fo00051g
- ⁹Hamedi, S., Honarvar, M., *Beta vulgaris* - A Mini Review of Traditional Uses in Iran, *Phytochemistry and Pharmacology, Curr. Drug Discov. Technol.*, **2019**, *16*, 74-81. DOI: 10.2174/1570163815666180308142912
- ¹⁰Ramanarayanan, R., Nijisha, P., Niveditha, C.V., Sindhu S., Natural dyes from red amaranth leaves as light-harvesting pigments for dye-sensitized solar cells, *Mater. Res. Bull.*, **2017**, *90*, 156-161. <http://dx.doi.org/10.1016/j.materresbull.2017.02.037>
- ¹¹Sarker, U., Oba, S., Antioxidant constituents of three selected red and green color Amaranthus leafy vegetable, *Sci. Rep.*, **2019**, *9*, Article 18233. <https://doi.org/10.1038/s41598-019-52033-8>
- ¹²Sumeet, D., Raghvendra, D., Kushagra, M., *Achyranthes aspera* Linn. (Chirchira): A Magic Herb in Folk Medicine, *Ethnobot. Leafl.*, **2008**, *12*, 670-676. <https://www.researchgate.net/publication/41115502>
- ¹³Verma, R. K., Ethnomedicinal Potential of *AchyranthesAspera* Linn. (Amaranthaceae), *Int. J. Sci. Res.*, **2014**, *3*, 55-60. DOI: 10.36106/ijsr
- ¹⁴Chinsebu, K. C., Ethnobotanical Study of Plants Used in the Management of HIV/AIDS-Related Diseases in Livingstone, Southern Province, Zambia, *Evid. Based Complement. Alternat. Med.*, **2016**, Article 4238625. <http://dx.doi.org/10.1155/2016/4238625>
- ¹⁵Tounekti, T., Mahdhi, M., Khemira, K., Ethnobotanical Study of Indigenous Medicinal Plants of Jazan Region, Saudi Arabia, *Evid. Based Complement. Alternat. Med.*, **2019**, Article 3190670. <https://doi.org/10.1155/2019/3190670>
- ¹⁶Mahmoud, T., Sanjay Gairola, S., Traditional knowledge and use of medicinal plants in the Eastern Desert of Egypt: a case study from Wadi El-Gemal National Park, *J. Med. Plants Stud.*, **2013**, *1*, 10-17. <https://www.researchgate.net/publication/270884815>
- ¹⁷Appiah, K. S., Oppong, C. P., Mardani, H. K., Omari, R. A., Kpabitey, S., Amoatey, C. A., Onwona-Agyeman, S., Oikawa, Y., Katsura, K., Fujii, Y., Medicinal Plants Used in the Ejisu-Juaben Municipality, Southern Ghana: An Ethnobotanical Study, *Medicines*, **2019**, *6*, 1-27. DOI:10.3390/medicines6010001
- ¹⁸Wali, S., Jan, H. A., Bussmann, R. W., Quantitative ethnomedicinal study of indigenous medicinal plants used for digestive disorders of Laspur Valley, Chitral, Northern Pakistan, *Ethnobot. Res. Appl.*, **2019**, *18*, 1-18. <http://dx.doi.org/10.32859/era.18.32.1-18>
- ¹⁹Abbasi, A., M., Khan, M. A., Shah, M. H., Shah, M. M., Pervez, A., Ahmad, M., Ethnobotanical appraisal and cultural values of medicinally important wild edible vegetables of Lesser Himalayas-Pakistan, *J. Ethnobiol. Ethnomed.*, **2013**, *9*, Article 66, 13 pages. DOI:10.1186/1746-4269-9-66
- ²⁰Miraj, S., A Medicinal plant with antioxidant activity in Iranian folk medicine: *Amaranthus*, *Der Pharm. Lett.*, **2016**, *8*, 27-30. <https://www.researchgate.net/publication/308658809>
- ²¹Altundag, E., Ozturk, M., Ethnomedicinal studies on the plant resources of east Anatolia, Turkey, *Procedia Soc. Behav. Sci.*, **2011**, *19*, 756-777. DOI:10.1016/j.sbspro.2011.05.195
- ²²Ari, S., Temel, M., Kargioglu, M., Konuk, M., Ethnobotanical survey of plants used in Afyonkarahisar-Turkey, *J. Ethnobiol. Ethnomed.*, **2015**, *11*, Article 84, 15 pages. DOI: 10.1186/s13002-015-0067-6
- ²³Yesil, Y., Celik, M., Yilmaz, B., Wild edible plants in Yesilli (Mardin-Turkey), a multicultural area, *J. Ethnobiol. Ethnomed.*, **2019**, *15*, Article 52, 19 pages. <https://doi.org/10.1186/s13002-019-0327-y>
- ²⁴Moshi, M. J., Otieno, D. F., Weisheit, A., Ethnomedicine of the Kagera Region, north western Tanzania. Part 3: plants used in traditional medicine in Kikuku village, Muleba District, *J. Ethnobiol. Ethnomed.*, **2012**, *8*, Article 14, 25 pages. DOI:10.1186/1746-4269-8-14
- ²⁵Zahoor, M., Yousaf, Z., Aqsa, T., Haroon, M., Saleh, N., Aftab, A., Javed, S., Qadeer, M., Ramazan, H., An ethnopharmacological evaluation of Navapind and Shahpur Virkanin district Sheikupura, Pakistan for their herbal Medicines, *J. Ethnobiol. Ethnomed.*, **2017**, *13*, Article 27, 26 pages. DOI: 10.1186/s13002-017-0151-1
- ²⁶Abdul Aziz, M., Khan, H. A., Ullah, H., Adnan, M., Hashem, A., Abd Allah, E. F., Traditional phytomedicines for gynecological problems used by tribal communities of Mohmand Agency near the Pak-Afghan border area, *Braz. J. Pharmacog.*, **2018**, *28*, 503-511. <https://doi.org/10.1016/j.bjp.2018.05.003>
- ²⁷Umair, M., Altaf, M., Bussmann, R. W., Abbasi, A. M., Ethnomedicinal uses of the local flora in Chenab riverine area, Punjab province Pakistan, *J. Ethnobiol. Ethnomed.*, **2019**, *15*, Article 7, 31 pages. <https://doi.org/10.1186/s13002-019-0285-4>
- ²⁸Adnan, M., Ullah, I., Tariq, A., Murad, W., Azizullah, A., Khan, A., Ali, N., Ethnomedicine use in the war affected region of northwest Pakistan, *J. Ethnobiol. Ethnomed.*, **2014**, *10*, Article 16, 16 pages. DOI:10.1186/1746-4269-10-16
- ²⁹Khan, I., AbdElsalam, N. M., Fouad, H., Tariq, A., Ullah, R., Adnan, M., Application of Ethnobotanical Indices on the Use of Traditional Medicines against Common Diseases, *Evid. Based Complement. Alternat. Med.*, **2014**, Article 635371, 21 pages. <http://dx.doi.org/10.1155/2014/635371>
- ³⁰Pieroni, A., Cattero, V., Wild vegetables do not lie: Comparative gastronomic ethnobotany and ethnolinguistics on the Greek traces of the Mediterranean, Diet of southeastern Italy, *Acta Bot. Bras.*, **2019**, *33*, 198-211. DOI: 10.1590/0102-33062018abb0323
- ³¹Goyal, B. R., Goyal, R. K., Mehta, A. A., Phyto-pharmacology of *Achyranthes aspera*: A Review, *Pharmacog. Rev.*, **2007**, *1*, 143-150. <http://www.phcogrev.com/article/2007/1/1-15>
- ³²Varuna, K. M., Khan, M. U., Sharma, P. K., Review on *Achyranthes Aspera*, *J. Pharm. Res.*, **2010**, *3*, 714-717. <https://www.researchgate.net/publication/44035930>
- ³³Dey, A., *Achyranthes aspera* L: Phytochemical and Pharmacological Aspects, *Int. J. Pharmaceut. Sci. Rev. Res.*, **2011**, *9*, 72-82. <https://www.researchgate.net/publication/286592553>
- ³⁴Shukla, A., Bukhariya, V., Mehta, J., Bajaj, J., Charde, R., *Achyranthes aspera* Linn. (Chirchira): A Magic Herb in Folk Medicine, *Int. J. Biomed. Adv. Res.*, **2011**, *2*, 228-240. <https://www.researchgate.net/publication/41115502>
- ³⁵Srivastav, S., Singh, P., Mishra, G., Jha, K. K., Khosa, R. L., *Achyranthes aspera*-An important medicinal plant: A review, *J. Nat. Prod. Plant Resour.*, **2011**, *1*, 1-14. <https://www.researchgate.net/publication/216410155>

- ³⁶Hasan, S., Pharmacological and Medicinal Uses of *Achyranthes aspera*, *Int. J. Sci. Environ. Technol.*, **2014**, *3*, 123-129. <https://www.researchgate.net/publication/260983682>
- ³⁷Srivastava, P. K., *Achyranthes aspera*: A Potent Immunostimulating Plant for Traditional Medicine, *Int. J. Pharmaceut. Sci. Res.*, **2014**, *5*, 1601-1611. DOI:10.13040/IJPSR.0975-8232.5(5).1601-11
- ³⁸Sharma, V., Chaudhary, U., An Overview on Indigenous Knowledge of *Achyranthes aspera*, *J. Crit. Rev.*, **2015**, *2*, 7-19. <http://www.jcreview.com/fulltext/197-1572431201.pdf>
- ³⁹Sureshkumar, S., Roopan, D. J., Sivaraj, A., Devi, K., Senthilkumar, B., Pharmacognosy of *Achyranthes aspera* Linn: A Review, *Adv. Res. J. Life Sci.*, **2015**, *1*, 1-5. <http://arjls.info/index.php/arjls/article/view/1>
- ⁴⁰Parmar, H. K., Sharma, D., Pharmacological and Medicinal importance of *Achyranthes aspera*: A review, *Int. J. Pharm. Life Sci.*, **2015**, *6*, 4627-4630. <https://www.researchgate.net/publication/311886486>
- ⁴¹Ghimire, K., Banerjee, J., Gupta, A. K., Dahal, P., Phytochemical Constituents and Pharmacological Uses of Medicinal Plant *Achyranthes aspera*: A Review, *World J. Pharmaceut. Res.*, **2015**, *4*, 470-489. <https://www.researchgate.net/publication/277248219>
- ⁴²Sanjay, P., A Review Article on Phytochemical and Pharmacological Profiles of Apamarga (*Achyranthes aspera* Linn), *Int. Ayurv. Med. J.*, **2015**, *3*, 2901-2909. http://www.iamj.in/posts/images/upload/2901_2909.pdf
- ⁴³Vijayaraj, R., Vidhya, R., Biological Activity of *Achyranthes Aspera* Linn. - A Review, *Asian J. Biochem. Pharmaceut. Res.*, **2016**, *6*, 86-93. <https://www.researchgate.net/publication/303700963>
- ⁴⁴Saini, S., Review on *Achyranthes Aspera* L., *Int. Edu. Res. J.*, **2016**, *2*, 84-86. <http://ierj.in/journal/index.php/ierj/article/view/156>
- ⁴⁵Verma, S., A review study on *Achyranthes aspera* (Amaranthaceae)-A valuable medicinal herb, *J. Med. Plants Stud.*, **2016**, *4*, 6-7. <http://www.plantsjournal.com/archives/2016/vol4issue3/PartA/4-2-7.pdf>
- ⁴⁶Lakshmi, V., Mahdi, A. A., Sharma, D., Agarwal, S. K., An Overview of *Achyranthes aspera* Linn., *J. Sci. Innov. Res.*, **2018**, *7*, 27-29. http://www.isirjournal.com/Vol7_Issue1_07.pdf
- ⁴⁷Rehman, R., Melki, D., Shehzad, A., Nadeem, F., Khalid, T., Commercial Importance, Medicinal Value and Therapeutic Potentials of Chaff Flower (*Achyranthes aspera*) – A Review, *Int. J. Chem. Biochem. Sci.*, **2018**, *14*, 62-70. <http://www.iscientific.org/wp-content/uploads/2019/09/7-IJCBS-18-14-7.pdf>
- ⁴⁸Singh, N., Sharma, P. M., Gupta, V. K., A Review on Pharmacological Aspects of *Achyranthes Aspera*, *Int. J. Pharmacog. Chin. Med.*, **2019**, *3*, Article 000188, 10 pages. DOI:10.23880/ipcm-16000188
- ⁴⁹Movaliya, V., Zaveri, M., A Review on the Pashanbheda Plant “*Aerva javanica*”, *Int. J. Pharm. Sci. Rev. Res.*, **2014**, *25*, 268-275. <https://www.researchgate.net/publication/261878577>
- ⁵⁰Payal, C., Gurlaganjeet, K., Davinder, K., Gagan, S., Amit, C., Dhawan, R. K., A Review on Phytochemistry and Biological Activities of *Aerva.Med.Aromat.Plants*, **2015**, *4*, Article 187, 4 pages. DOI:10.4172/2167-0412.1000187
- ⁵¹Walter, T. M., Merish, S., Tamizhamuthu, M., Review of *Alternanthera Sessilis* with Reference to Traditional Siddha Medicine, *Int. J. Pharmacog. Phytochem. Res.*, **2014**, *6*, 249-254. <https://www.researchgate.net/publication/263085590>
- ⁵²Sravani, V. L., Abbas, Z., Surya, P., Review on *Alternanthera Sessilis*, *Indo Am. J. P. Sci.*, **2017**, *4*, 2845-2852. <http://www.iajps.com/pdf/september2017/20.IAJPS20092017.pdf>
- ⁵³Kumar, Y. S., Sanjib, D., Evaluation of Anti-Diarrhoeal Property of Crude Aqueous Extract of *Alternanthera Sessilis* Linn., *Int. J. Pharmaceut. Innov.*, **2013**, *3*, 110-115. <https://pdfs.semanticscholar.org/b4ea/d625fc4e4e085e700911ecca5d196be0f6df.pdf>
- ⁵⁴Rao, P., Ayurvedic View of *Alternanthera sessilis* Linn. with Special Reference to Mathsyakshi: A Brief Review, *SFJ. Herb Med.*, **2018**, *2*, Article 1000013, 6 pages. <https://www.researchgate.net/publication/335391966>
- ⁵⁵Shehzad, A., Qayyum, A., Rehman, R., Nadeem, F., Shehzad, M. R., A Review of Bioactivity Guided Medicinal Uses and Therapeutic Potentials of Noxious Weed (*Alternanthera sessilis*), *Int. J. Chem. Biochem. Sci.*, **2018**, *14*, 95-103. <https://www.researchgate.net/publication/336825620>
- ⁵⁶Chandrashekhar, K., Ethnobotanical and Phyto pharmacological Overview of Matsyakshi (*Alternanthera sessilis* R Br.ex DC), *J. Ayu. Herb. Med.*, **2019**, *5*, 152-155. http://www.ayurvedjournal.com/JAHM_201954_07.pdf
- ⁵⁷Wolosik, K., Markowska, A., *Amaranthus Cruentus*- Taxonomy, Botanical Description and Review of its Seed Chemical Composition, *Nat. Prod. Commun.*, **2019**, *14*, 1–10. DOI: 10.1177/1934578X19844141
- ⁵⁸Ward, S. M., Webster, T. M., Steckel, L. E., Palmer Amaranth (*Amaranthus palmeri*): A Review, *Weed Technol.*, **2013**, *27*, 12–27. DOI: 10.2307/23358302
- ⁵⁹Dufour-Dror, J. M., Fragan-Sapir, O., (Eds)“Alien plants species in Natural and Disturbed areas in Israel”, in *Alien Invasive plants in Israel*, 2nd edition, **2018**, 2-9, Cana publishing, Jerusalem, Israel. DOI: 10.13140/RG.2.2.16505.36965
- ⁶⁰Jhade, D., Ahirwar, D., Jain, R., Sharma, N. K., Gupta, S., A Pharmacological Review: *Amaranthus spinosus*, *Res. J. Pharmacog. Phytochem.*, **2009**, *1*, 169-172. <https://www.researchgate.net/publication/309076182>
- ⁶¹Kawade, R. M., Ghiware, N. B., Sarje, S. K., Vadvalkar, M. S., Pharmacognostic and Pharmacological Review: *Amaranthus spinosus*, *World J. Pharmaceut. Res.*, **2013**, *2*, 2099-2110. <https://www.researchgate.net/publication/281857872>
- ⁶²Kumar, R. P., Shammy, J., Nitin, G., Rinu, R., An Inside Review of *Amaranthus spinosus* Linn: A Potential Medicinal Plant of India, *Int. J. Res. Pharm. Chem.*, **2014**, *4*, 643-653. <http://www.ijrpc.com/files/20-4130.pdf>
- ⁶³Tanmoy, G., Arijit, M., Tanushree, S., Jagadish, S., Kumar, M. T., Pharmacological Actions and Phytoconstituents of *Amaranthus spinosus* Linn: A Review, *Int. J. Pharmacog. Phytochem. Res.*, **2014**, *6*, 405-413. <http://impactfactor.org/PDF/IJPPR/6/IJPPR.Vol6.Issue2.Article46.pdf>
- ⁶⁴Chandrashekhar, K., A Review on Tanduliyaka (*Amaranthus spinosus* L) - A Weed, A Vegetable and A Medicinal Plant, *Int. J. Ayu. Med.*, **2018**, *9*, 231-238. <https://ijam.co.in/index.php/ijam/article/view/1169>
- ⁶⁵Ganjare, A., Raut, N., Nutritional and medicinal potential of *Amaranthus spinosus*, *J. Pharmacog. Phytochem.*, **2019**, *8*, 3149-3156. <http://www.phytojournal.com/archives/2019/vol8issue3/PartAS/8-3-128-378.pdf>
- ⁶⁶Sowjanya, P., Babu, P. S., Narasu, M. L., Phytochemical and Pharmacological Potential of *Amaranthus viridis* L.: A Review, *Int. J. Phytomed.*, **2014**, *6*, 322-326. <https://www.arjournals.org/index.php/ijpm/article/view/1444/784>
- ⁶⁷Ferdous, R., Shahjahan, S., Tanvir, S., Mukti, M., Present Biological Status of Potential Medicinal Plant of *Amaranthus viridis*: A Comprehensive Review, *Am. J. Clin. Exper. Med.*, **2015**, *3*, 12-17. DOI:10.11648/j.ajcem.s.2015030501.13

- ⁶⁸Singh, R., Chauhan, D. C., Tomar, P., Tanduliyaka (*Amaranthus viridis*): A Highly Nutritional Herb. *Int. J. Ayu. Pharm. Res.*, **2016**, *4*, 53-55. <https://www.ijapr.in/index.php/ijapr/article/view/350>
- ⁶⁹Graca Miguel, M., Betalains in Some Species of the *Amaranthaceae* Family: A Review, *Antioxidants*, **2018**, *7*, 53-85. DOI:10.3390/antiox7040053
- ⁷⁰Alegbejo, J. O., Nutritional Value and Utilization of *Amaranthus* (*Amaranthus* spp.) - A Review, *Bayero J. Pure Appl. Sci.*, **2013**, *6*, 136-143. <http://dx.doi.org/10.4314/bajopas.v6i1.27>
- ⁷¹Thapa, R., W. Blair, M. W., Morphological Assessment of Cultivated and Wild *Amaranth* Species Diversity, *Agronomy*, **2018**, *8*, 272-284. DOI:10.3390/agronomy8110272
- ⁷²Peter, K., Gandhi, P., Rediscovering the therapeutic potential of *Amaranthus* species: A review, *Egypt. J. Basic Appl. Sci.*, **2017**, *4*, 196-205. <http://dx.doi.org/10.1016/j.ejbas.2017.05.001>
- ⁷³Alam, A., Slahin, N., Uddin, R., Hasan, R., Akter, R., Kamaluddin, M., Faroque, A., Ghani, A., Analgesic and Neuropharmacological Investigations of the Aerial Part of *Achyranthes aspera* Linn., *Stamford J. Pharm. Sci.*, **2008**, *1*, 44-50. DOI: 10.3329/sips.vlil.1807
- ⁷⁴Kamble, A., Antiallergy activity of *Achyranthes aspera*, *World Sci. News*, **2018**, *106*, 24-45. <http://www.worldscientificnews.com/wp-content/uploads/2018/07/WSN-106-2018-24-45-2.pdf>
- ⁷⁵Barua, C. C., Talukdar, A., Begum, S. A., Borah, P., Lahkar, M., Anxiolytic activity of methanol leaf extract of *Achyranthes aspera* Linn in mice using experimental models of anxiety, *Indian J. Pharmacol.*, **2012**, *44*, 63-67. DOI: 10.4103/0253-7613.91869
- ⁷⁶Kothavade, P. S., Bulani, V. D., Nagmoti, D. M., Deshpande, P. S., Gawali, N. B., Juvekar, A. R., Therapeutic Effect of Saponin Rich Fraction of *Achyranthes aspera* Linn. on Adjuvant-Induced Arthritis in Sprague-Dawley Rats, *Autoimmune Dis.*, **2015**, Article 943645, 8 pages. <http://dx.doi.org/10.1155/2015/943645>
- ⁷⁷Chinnasamy, V., Subramaniyan, V., Chandiran, S., Kayarohanam, S., Kannian, D. C., Velaga, V. S., Muhammad, S., Antiarthritic Activity of *Achyranthes Aspera* on Formaldehyde-Induced Arthritis in Rats, *Maced. J. Med. Sci.*, **2019**, *7*, 2709-2714. <https://doi.org/10.3889/oamjms.2019.559>
- ⁷⁸Alam, M. T., Karim, M. M., Khan, S. N., Antibacterial Activity of Different Organic Extracts of *Achyranthes Aspera* and *Cassia Alata*, *J. Sci. Res.*, **2009**, *1*, 393-398. DOI: 10.3329/jsr.v1i2.2298
- ⁷⁹Prasad, S. H., Swapna, N. L., Anthonamma, K., Madanprasad, A., Rajasekhar, D., Antimicrobial activity of *Achyranthes aspera* and *Aerva lanata* leaf and callus extracts, *Biosci., Biotech. Res. Asia*, **2009**, *6*, 887-891. <https://www.researchgate.net/publication/286220770>
- ⁸⁰Narayan, G. R., Viswas, K., Pathak, M., Singh, P. S., Gupta, A., Antibacterial activities of ethanolic extracts of plants used in folk medicine. *Int. J. Res. Ayu. Pharm.*, **2010**, *1*, 529-535. <https://www.cabdirect.org/cabdirect/abstract/20113195417>
- ⁸¹Ranwan, S., Yadav, J. P., Seasonal variation in antibacterial activity of different parts of *Achyranthes aspera* against some bacteria, *Int. J. Med. Arom. Plants*, **2012**, *2*, 369-375. <https://www.researchgate.net/publication/249991047>
- ⁸²Konala, G., Seva, L., Davuluri, S. P., Antimicrobial Activity of Prickly Chaff (*Achyranthes aspera*) Seed Trypsin Inhibitor. *Int. J. Pharm. Sci. Res.*, **2012**, *3*, 3241-3248. [http://dx.doi.org/10.13040/IJPSR.0975-8232.3\(9\).3241-48](http://dx.doi.org/10.13040/IJPSR.0975-8232.3(9).3241-48)
- ⁸³Pandey, R., Sambasivarao, Y., Gurumurthy, Y., Antibacterial Activity of Medicinal Plants against Pathogens from Extracts of *Achyranthes Aspera*, *Med. Aromat. Plants*, **2013**, *2*, 135-137. DOI: 10.4172/2167-0412.1000135
- ⁸⁴Patil, U., Sharma, M. C., Studies on antibacterial effect of apamarga (*Achyranthes aspera*) on multi-drug resistant clinical isolates, *Int. J. Res. Ayurv. Pharm.*, **2013**, *4*, 262-265. DOI: 10.7897/2277-4343.04236
- ⁸⁵Sharma, V., Singh, R., Chaudhary, U., Curative efficacy of *Achyranthes asper* against various Pathogenic Microbial Strains, *Int. J. Drug Dev. Res.*, **2013**, *5*, 392-401. <https://www.researchgate.net/publication/261684238>
- ⁸⁶Yadav, R., Rai, R., Yadav, A., Pahuja, M., Solanki, S., Yadav, H., Evaluation of antibacterial activity of *Achyranthes aspera* extract against *Streptococcus mutans*: An invitro study, *J. Adv. Pharm. Technol. Res.*, **2016**, *7*, 149-152. DOI: 10.4103/2231-4040.191426
- ⁸⁷Lavanya, R., Antibacterial Activity of *Achyranthes aspera* Linn., *Indian Am. J. Pharmaceut. Res.*, **2017**, *7*, 7656-7660. <https://zenodo.org/record/2383287#Xxkbsp4zblU>
- ⁸⁸Parveen, P., Phytochemical Screening and Antimicrobial Activity of Herbal Plant Extracts- *Achyranthes aspera*, *Int. J. Pharmaceut. Clin. Res.*, **2018**, *10*, 201-209. <https://www.researchgate.net/publication/312630817>
- ⁸⁹Londonkar, R., Reddy, V. C., Kumar, K. A., Potential antibacterial and antifungal activity of *Achyranthes aspera* L., *Recent Res. Sci. Technol.*, **2011**, *3*, 53-57. <https://www.researchgate.net/publication/285310689>
- ⁹⁰Ndhala, A. R., Ghebrehiwot, H. M., Ncube, B., Aremu, A. O., Gruz, J., Subrtova, M., Dolezal K., du Plooy C. P., Abdelgadir, H. A., Van Staden, J., Antimicrobial, Anthelmintic Activities and Characterisation of Functional Phenolic Acids of *Achyranthes aspera* Linn.: A Medicinal Plant Used for the Treatment of Wounds and Ringworm in East Africa, *Front. Pharmacol.*, **2015**, *6*, Article 274. 8 pages. DOI:10.3389/fphar.2015.00274
- ⁹¹Kaveti, B., Xian, P. X., Fern, R. C., Lee, M. T., Baig, M., Antimicrobial and Antioxidant Activities of the Seeds of *Achyranthes Aspera*, *Int. J. Pharm. Teach. Pract.*, **2013**, *4*, 747-751. <https://www.iomcworld.org/articles/antimicrobial-and-antioxidant-activities-of-the-seeds-of-achyranthes.pdf>
- ⁹²Elumalai, E. K., Chandrasekaran, N., Thisrumalai, T., Sivakumar, C., Therasa, S. V., David, E., *Achyranthes aspera* Leaf Extracts Inhibited Fungal Growth, *Int. J. PharmTech Res.*, **2009**, *1*, 1576-1579. [http://sphinxsai.com/PTVOL4/pdf_vol4/PT-98%20\(1576-1579\).pdf](http://sphinxsai.com/PTVOL4/pdf_vol4/PT-98%20(1576-1579).pdf)
- ⁹³Baraik, B., Jain, P., Sharma, H. P., *Achyranthes aspera* L.: As a Source of Biofungicide, *Am. J. Adv. Drug Deliv.*, **2014**, *2*, 686-696. <https://www.researchgate.net/publication/272106354>
- ⁹⁴Waghmare, V., Khandagle, A. J., Morey, R., Antifungal Potential of *Parthenium hysterophorus*, *Achyranthes aspera* and *Catharanthus roseus* against *Aspergillus sp.*, *Candida sp.* and *Penicillium sp.* of Aquarium Fish, *Egypt. Acad. J. Biol. Sci.*, **2020**, *12*, 1-8. https://ejbsz.journals.ekb.eg/article_69337.html
- ⁹⁵Misra, T. N., Singh, R. S., Pandey, H. S., Singh, B. P., Antifungal essential oil and long chain alcohol from *Achyranthes aspera*, *Phytochemistry*, **1992**, *31*, 1811-1812. [https://doi.org/10.1016/0031-9422\(92\)83153-P](https://doi.org/10.1016/0031-9422(92)83153-P)
- ⁹⁶Palshetkar, A., Pathare, N., Jadhav, N., Pawar, M., Wadhvani, A., Kulkarni, S., Singh, K. K., *In vitro* anti-HIV activity of some Indian medicinal plant extracts, *BMC Complement. Med. Therap.*, **2020**, *20*, Article 69, 11 pages. <https://doi.org/10.1186/s12906-020-2816-x>
- ⁹⁷Hullatti, K. K., Murthy, U. D., Activity Guided Isolation of Cytotoxic Compounds from Indian Medicinal Plants Using BSL Bioassay, *J. Curr. Pharmaceut. Res.*, **2010**, *1*, 16-18. <https://www.researchgate.net/publication/44900174>
- ⁹⁸Arora, S., Tandon, S., *Achyranthes aspera* Root Extracts Induce Human Colon Cancer Cell (COLO-205) Death by Triggering the Mitochondrial Apoptosis Pathway and S Phase Cell Cycle Arrest, *Sci. World J.*, **2014**, Article129697, 15 pages. <http://dx.doi.org/10.1155/2014/129697>

- ⁹⁹Akhtar, M. S., Iqbal, J., Evaluation of the hypoglycaemic effect of *Achyranthes aspera* in normal and alloxan-diabetic rabbits, *J. Ethnopharmacol.*, **1991**, *31*, 49-57. DOI: 10.1016/0378-8741(91)90143-2
- ¹⁰⁰Kumar A. S., Gnananath, K., Gande, S., Goud E. R., Rajesh, P., Nagarjuna, S., *J. Pharm. Res.*, **2011**, *4*, 3124-3125. <https://www.researchgate.net/publication/230793855>
- ¹⁰¹Mahendran, S., Thangavelu, L., Roy, A., Advanced glycation endproduct inhibitory assay- 'Achyranthes Aspera leaf', *J. Adv. Pharm. Educ. Res.*, **2017**, *7*, 479-481. <https://japer.in/en/article/advanced-glycation-end-product-inhibitory-assay-achyranthes-aspera-leaf>
- ¹⁰²Lakshmi, V., Mahdi, A. A., Mishra, V., Agarwal, S. K., Antidiabetic Potential of *Achyranthes Aspera* Leaves in Rats, *J. Biopharmaceut. Therap. Chal.*, **2018**, *2*, 1-3. <http://sciaeon.org/biopharmaceutics-and-therapeutic-challenges/archive/abstract/112>
- ¹⁰³Njideka, B. E., Theophilus, A. E., Ugochukwu, N. T., Use of *Achyranthes aspera* Linn tea as antidiabetic and hypolipidemic herbal tea, *Int. J. Health Sci. Res.*, **2019**, *9*, 32-38. https://ijhsr.org/IJHSR_Vol.9_Issue.2_Feb2019/6.pdf
- ¹⁰⁴Vijayaraj, R., Kumaran, N. S., Evaluation of *in vivo* antidiabetic and antioxidant activity of *Achyranthes aspera* Linn. seeds by streptozotocin induced diabetic rats, *Int. J. Green Pharm.*, **2018**, *12*, 29-37. <https://www.greenpharmacy.info/index.php/ijgp/article/view/1520/0>
- ¹⁰⁵Rahman, H.M., Bashir, S., Mandukhail, S. R., Huda, S., Gilani, A. H., Pharmacological Evaluation of Gut Modulatory and Bronchodilator Activities of *Achyranthes aspera* Linn., *Phytother. Res.*, **2017**, *31*, 1776-1785. DOI: 10.1002/ptr.5907
- ¹⁰⁶Vetrichelvan, T., Jegadeesan, M., Effect of alcohol extract of *Achyranthes aspera* Linn. on acute and subacute inflammation, *Phytother. Res.*, **2003**, *17*, 77-79. DOI: 10.1002/ptr.1070
- ¹⁰⁷Kumar, S.V., Sankar, P., Varatharajan, R., Anti-inflammatory activity of roots of *Achyranthes aspera*, *Pharmaceut. Biol.*, **2009**, *47*, 973-975. DOI: 10.1080/13880200902967979
- ¹⁰⁸Bhosale, U. A., Yegnanarayan, R., Pophale, P., Somani, R., Effect of aqueous extracts of *Achyranthes aspera* Linn. on experimental animal model for inflammation, *Anc. Sci. Life*, **2012**, *31*, 202-206. DOI: 10.4103/0257-7941.107362: 10.4103/0257-7941.107362
- ¹⁰⁹Kandasamy, R., Hellermann, G. R., Mohapatra, S. S., Lockey, R. F., Flavonoid-Rich Alcoholic Extract of Leaves of *Achyranthes aspera* Reduces Inflammation in a Murine Model of Ova-Induced Asthma, *J. Allergy Clin. Immunol.*, **2012**, *129*, AB78. <https://doi.org/10.1016/j.jaci.2011.12.614>
- ¹¹⁰Boyapati, R., Gojja, P., Chintalapani, S., Nagubandi, K., Ramiseti, A., Salavathi, S. S., Efficacy of local drug delivery of *Achyranthes aspera* gel in the management of chronic periodontitis: A clinical study, *J. Indian Soc. Periodontol.*, **2017**, *21*, 46-49. DOI:10.4103/jisp.jisp_130_17
- ¹¹¹Uma, B., Yegnanarayan, R., Prachi, P., Zambare, M., Somani, R. S., Antinociceptive evaluation of an ethanol extract of *Achyranthes aspera* (agadha) in animal models of Nociception, *Int. J. Phytomed.*, **2010**, *2*, 440-445. DOI:10.5138/ijpm.2010.0975.0185.02063
- ¹¹²Shahid, I. Z., Moniruzzaman, M., Das, A. K., Choudhuri, M. S., Firoj, A., Pharmacological studies on roots of *Achyranthes aspera* Linn, *Orient. Pharm. Exp. Med.*, **2006**, *6*, 202-207. <https://doi.org/10.3742/OPEM.2006.6.3.202>
- ¹¹³Latha, B. P., Reddy, I. R., Vijaya, T., Rao, S. D., Ismail, S. M., Girish, B. P., Effect of saponin rich extract of *Achyranthes aspera* on high fat diet fed male wistar rats, *J. Pharm. Res.*, **2011**, *4*, 3190-3193. <https://www.researchgate.net/publication/269942967>
- ¹¹⁴Nehete, J. Y., Deshmukh, V. N., Shewale, V. V., Narkhede, M. R., Aurangabadkar, V. M., *In-vitro* antioxidant activity of *Achyranthes aspera* L., *J. Pharm. Res.*, **2009**, *2*, 1402-1403. <https://www.researchgate.net/publication/244988081>
- ¹¹⁵Priya, C. L., Kumar, G., Karthik, L., Rao, K. V., Antioxidant Activity of *Achyranthes aspera* Linn. Stem Extracts, *Pharmacologyonline*, **2010**, *2*, 228-237. <https://www.researchgate.net/publication/213173733>
- ¹¹⁶Babu, M. N., Elango, K., Pharmacognostical, Phytochemical and Antioxidant studies of *Achyranthes aspera* Linn. and *Achyranthes bidentata* Blume, *J. Pharm. Res.*, **2011**, *4*, 1050-1055. <http://jprsolutions.info/newfiles/journal-file-56d1bae16b1e50.17082187.pdf>
- ¹¹⁷Sharma, V., Chaudhary, U., Singh, R., Janmeda, P., Evaluation of quantitative and antioxidant activity of *Achyranthes aspera* roots and inflorescences, *Asian J. Pharmaceut.*, **2014**, *8*, 54-60. DOI:10.4103/0973-8398.134085
- ¹¹⁸Bahule, B. B., Mahajan, R. P., More, D. H., Antioxidant Activity of Wood and Leaf Extracts of *Achyranthes aspera* Using Radical Scavenging Method, *Int. J. Chem. Sci.*, **2017**, *15*, Article 217, 5. <https://www.tsijournals.com/13503.html>
- ¹¹⁹Sivasankari, S., Banu, B. J., Sadiq, A. M., Kanagavalli, U., Phytochemical evaluation and Antioxidant potential of ethanolic leaf extract of *Achyranthes aspera*, *Int. Res. J. Pharmaceut. Biosci.*, **2017**, *4*, 15-24. <https://irjpbs.com/volumes/vol4/issue5/IRJPBS-4502.pdf>
- ¹²⁰Dissanayake D. P., Sivaganesh, S., Tissera M. H., Handunnetti, S. M., Arawawala, L. D., Comparison of antioxidant properties of *Cyathula prostrata* Linn and *Achyranthes aspera* Linn. grown in Sri Lanka, *Res. Rev. Insights*, **2018**, *2*, 1-3. DOI: 10.15761/RRI.1000138
- ¹²¹Sinan, K. I., Zengin, G., Zheleva-Dimitrova, D., Etienne, O. K., Mahomoodally, M. F., Bouyahya, A., Lobine, D., Chiavaroli, A., Ferrante, C., Menghini, L., Recinella, L., Brunetti, L., Leone, S., Orlando, G., Qualitative Phytochemical Fingerprint and NetworkPharmacology Investigation of *Achyranthes aspera* Linn. Extracts, *Molecules*, **2020**, *25*, 1973-1991. DOI:10.3390/molecules25081973
- ¹²²Tahiliani, P., Kar, A., *Achyranthes aspera* elevates thyroid hormone levels and decreases hepatic lipid peroxidation in male rats, *J. Ethnopharmacol.*, **2000**, *71*, 527-532. DOI: 10.1016/s0378-8741(00)00170-7
- ¹²³Kothapalli, L., Gite, P. R., Thomas, A., Nanda, R., Salvi, N., Evaluation of Antivenom Potential of *Achyranthes aspera* Leaves Against Snake Venom, *Int. J. Pharm. Drug Anal.*, **2016**, *4*, 505-513. <http://www.ijpda.com/admin/uploads/CG0BDX.pdf>
- ¹²⁴Nwune, H. C., Milala, M. A., Zanna, H., Effects of Aqueous Stem Extract of *Achyranthes aspera* on *Bitis arietans* Venom Protease and Phospholipase A2 Activity, *Am. J. BioSci.*, **2017**, *5*, 54-58. DOI: 10.11648/j.ajbio.20170503.13
- ¹²⁵Etim, E. A., Adebayo, Y. A., Ifeanyi, O. E., Effect of *Achyranthes Aspera* Leaf Extract on Hematological Parameters of Swiss Albino Mice, *World J. Public Health*, **2019**, *4*, 96-101. DOI: 10.11648/j.wjph.20190404.14
- ¹²⁶Hassan, S., Rasool, S., Niaz, U., Rehman, M. K., Ali, M. Y., Abbas, G., Rasool, N., Riaz, M., Abbas, K., Dermatological potential of crude extracts and different fractions of *Achyranthes aspera* Linn., *Afr. J. Pharm. Pharmacol.*, **2012**, *6*, 3096-3101. DOI: 10.5897/AJPP12.1209
- ¹²⁷Gawande, D.Y., Goel, R. K., Pharmacological validation of in-silico guided novel nootropic potential of *Achyranthes aspera* L., *J. Ethnopharmacol.*, **2015**, *175*, 324-334. DOI: 10.1016/j.jep.2015.09.025
- ¹²⁸Barua, C. C., Talukdar, A., Begum, S. A., Handique, A. K., Handique, G. K., Roy, J. D., Buragohain, B., Impact of *Achyranthes aspera* L. on Protein Profile in Impaired Wound Models, *Indo. Glob. J. Pharmaceut. Sci.*, **2011**, *1*, 13-24. <http://www.iglobaljournal.com/wp-content/uploads/2011/02/Barua-et-al-2.pdf>

- ¹²⁹Mondal, S., Ghosh, D., Ganapaty, S., Reddy, M. S., Ramakrishna, K., Evaluation of Healing Potential of *Achyranthes aspera* L. (Amaranthaceae) seeds in excision, incision, dead space and burn wound model-An *in-vivo* Study, *Pharmacog. J.*, **2016**, *8*, 300-306. DOI:10.5530/pj.2016.3.20
- ¹³⁰Aggarwal, A., Tandon, S., Singla, S. K., Tandon, C., Reduction of oxalate-induced renal tubularepithelial (NRK-52E) cell injury and inhibition of calcium oxalate crystallisation *in vitro* by aqueous extract of *Achyranthes aspera*, *Int. J. Green Pharm.*, **2010**, *4*, 159-164. DOI: 10.4103/0973-8258.69173
- ¹³¹Niranjan, S., Alok, D. K., Soumya, M., Priyanka, G., Sangita, M. S., Diuretic Activity of *Achyranthes aspera* Leaves Extract, *Int. Res. J. Pharm.*, **2012**, *3*, 216-218. https://irjponline.com/admin/php/uploads/1009_pdf.pdf
- ¹³²Srivastav, S., Singh, P., Jha, K. K., Mishra, G., Srivastava, S., Karchuli, M. S., Khosa, R. L., Diuretic activity of whole plant extract of *Achyranthes aspera* Linn, *Euro. J. Exp. Bio.*, **2011**, *1*, 97-102. <https://www.researchgate.net/publication/216410620>
- ¹³³Asif, M., Jabeen, Q., Atif, M., Majid, A. M., Zaman, M., Diuretic Activity of *Achyranthes aspera* Linn. Crude Aqueous Extract in Albino Rats, *Trop. J. Pharm. Res.*, **2014**, *13*, 2039-2045. <http://dx.doi.org/10.4314/tjpr.v13i12.14>
- ¹³⁴Das, A. K., Bigoniya, P., Verma, N. K., Rana, A. C., Gastroprotective effect of *Achyranthes aspera* Linn. leaf on rats, *Asian Pac. J. Trop. Med.*, **2012**, *5*, 197-201. DOI: 10.1016/S1995-7645(12)60024-8
- ¹³⁵Kartik, R., Rao, C. V., Trivedi, S. P., Pushpangadan, P., Reddy, G. D., Amelioration effects against N-nitrosodiethylamine and CCl₄-induced hepatocarcinogenesis in Swiss albino rats by whole plant extract of *Achyranthes aspera*, *Ind. J. Pharmacol.*, **2010**, *42*, 370-375. DOI: 10.4103/0253-7613.71921
- ¹³⁶Manjunatha, B. K., Abhilash, N., Hegde, V., Suchitra, M. N., Vidya, S. M., Hepatoprotective Potency of *Achyranthes aspera*: An In-vivo Study, *Int. J. Pharm. Phytopharmacol. Res.*, **2012**, *1*, 387-390. <https://www.researchgate.net/publication/299487743>
- ¹³⁷Malarvili, T., Veerappan, R. M., Begum, V. H., Effect of *Achyranthes aspera* seeds on lipid profiles in selected tissues of rats fed with high doses of fructose, *J. Pharm. Res.*, **2011**, *4*, 1769-1771. <http://jpr solutions.info/newfiles/journal-file-56e15ce206c926.30903797.pdf>
- ¹³⁸Ramanathan, V., Thekkumalai, M., Modulatory effect of *Achyranthes aspera* L., seeds on carbohydrates and protein bound enzymes on high fructose fed diet, *J. Pharmacol. Toxicol.*, **2017**, *12*, 154-160. DOI: 10.3923/jpt.2017.154.160
- ¹³⁹Sarvesh, C. N., Fernandes, J., Evaluation of Antihyperlipidemic Activity of Leaves of *Achyranthes aspera* Linn. Using Hyperlipidemic Rats, *Asian J. Pharm. Clin. Res.*, **2017**, *4*, 211-215. <http://dx.doi.org/10.22159/ajpcr.2017.v10i4.16576>
- ¹⁴⁰Tanveer, A., Safdar, M. E., Farooq, N., Sudozai, M. I., Nadeem, M. A., Abbas, T., Exploring the Herbicidal Potential of *Achyranthes aspera* Against Some Weeds, *Planta Daninha*, **2019**, *37*, Article e019188205, 8 pages. DOI: 10.1590/S0100-83582019370100066
- ¹⁴¹Bagavan, A., Rahuman, A. A., Kamaraj, C., Geetha, K., Larvicidal activity of saponin from *Achyranthes aspera* against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae), *Parasitol. Res.*, **2008**, *103*, 223-229. DOI: 10.1007/s00436-008-0962-z
- ¹⁴²Sun, H. X., Adjuvant effect of *Achyranthes bidentata* saponins on specific antibody and cellular response to ovalbumin in mice, *Vaccine*, **2006**, *24*, 3432-3439. <https://doi.org/10.1016/j.vaccine.2006.02.014>
- ¹⁴³Sharma, A., Kumar, S., Tripathi, P., Impact of *Achyranthes aspera* Leaf and Stem Extracts on the Survival, Morphology and Behaviour of an Indian Strain of Dengue Vector, *Aedes aegypti* L. (Diptera: Culicidae), *J. Mosq. Res.*, **2015**, *5*, 1-9. DOI:10.5376/jmr.2015.05.0007
- ¹⁴⁴Sharma, A., Kumar, S., Tripathi, P., Effects of *Achyranthes aspera* Extracts on the Survival and Midgut Histo-architecture of *Aedes aegypti* L. Early IV Instars, *Open Parasitol. J.*, **2018**, *6*, 41-51. DOI: 10.2174/1874421401806010041
- ¹⁴⁵Mandefro, B., Mereta, S. T., Tariku, Y., Ambelu, A., Molluscicidal effect of *Achyranthes aspera* L. (Amaranthaceae) aqueous extract on adult snails of *Biomphalaria pfeifferi* and *Lymnaea natalensis*, *Infect. Dis. Poverty*, **2017**, *6*, Article 133, 5 pages. DOI: 10.1186/s40249-017-0349-4
- ¹⁴⁶Ikan, R., Ravid, U., Trosset, D., Shulman, E., Ecdysterone: an insect moulting hormone from *Achyranthes aspera* (amaranthaceae), *Experientia*, **1971**, *27*, 504-505. DOI: 10.1007/BF02147560
- ¹⁴⁷Kumar, C. B., Mohana, K. N., The Effect of *Achyranthes aspera* Extracts on Mild Steel Corrosion in Industrial Water Medium, *ISRN Corros.*, **2013**, Article 261847, 9 pages. <http://dx.doi.org/10.1155/2013/261847>
- ¹⁴⁸Sharma, A., Tripathi, P., Kumar, S., One-pot synthesis of silver nanocomposites from *Achyranthes aspera*: An eco-friendly larvicide against *Aedes aegypti* L., *Asian Pac. J. Trop. Biomed.*, **2020**, *10*, 54-64. DOI: 10.4103/2221-1691.275420
- ¹⁴⁹Tripathi, A., Kumari, S., Kumar, A., Toxicity evaluation of pH dependent stable *Achyranthes aspera* herbal gold nanoparticles, *Appl. Nanosci.*, **2016**, *6*, 61-69. DOI: 10.1007/s13204-015-0414-x
- ¹⁵⁰Pavithra, S., Mohana, B., Mani, M., Jayavel, R., Kumaresan, S., Physicochemical and Morphological Properties of *Achyranthes aspera* Mediated CuO Nanoparticles for Inhibiting Cellular Adhesion, *J. Clust. Sci.*, **2020**, no volume number, no paging, 11 pages. <https://doi.org/10.1007/s10876-020-01796-6>
- ¹⁵¹Sadashiv, P. S., Krishna, A. R., Acute toxicity study for *Achyranthes aspera* leaves, *J. Pharm. Res.*, **2011**, *4*, 2221-2222. <https://www.researchgate.net/publication/285188795>
- ¹⁵²Sandhyakumary, K., Bobby, R. G., Indira, M., Impact of feeding ethanolic extracts of *Achyranthes aspera* Linn. on reproductive functions in male rats, *Indian J. Exp. Biol.*, **2002**, *40*, 1307-1309. <http://nopr.niscair.res.in/handle/123456789/23582>
- ¹⁵³Vasudeva, N., Sharma, S. K., Estrogenic and Pregnancy Interceptive Effects of *Achyranthes Aspera* Linn. Root, *Afr. J. Tradit. Complement. Altern. Med.*, **2007**, *4*, 7-11. DOI: 10.4314/ajtcam.v4i1.31185
- ¹⁵⁴Vasudeva, N., Sharma, S. K., Post-coital antifertility activity of *Achyranthes aspera* Linn. root, *J. Ethnopharmacol.*, **2006**, *107*, 179-181. DOI: 10.1016/j.jep.2006.03.009
- ¹⁵⁵Shibeshi, W., Makonnen, E., Zerihun, L., Debella, A., Effect of *achyranthes aspera* L. on fetal abortion, uterine and pituitary weights, serum lipids and hormones, *Afr. Health Sci.*, **2006**, *6*, 108-112. <https://www.ajol.info/index.php/ahs/article/view/6930>
- ¹⁵⁶Paul, D., Mallick, C., Ali, K. M., Nandi, D. K., Duration dependent effect of hydro-ethanolic extract of leaf of *S. hermandifolia* and root of *A. aspera* on testicular androgenic and gametogenic activity: An approach for male herbal contraceptive development, *Int. J. Appl. Res. Nat. Prod.*, **2009**, *2*, 1-8. [http://journaldatabase.info/articles/duration dependent effect.html](http://journaldatabase.info/articles/duration%20dependent%20effect.html)
- ¹⁵⁷Paul, D., Ali, K. M., Ghosh, D., Nandi, D. K., Chenni, H., Trabelsi, M. M., Effect of Hydro-ethanolic (1:1) Extract of *Stephania hermandifolia* and *Achyranthes Aspera* in Composite Manner on Testicular Activity in Male Albino Rat: A Dose Dependent Analysis, *J. Reprod. Contracept.*, **2012**, *23*, 209-221. [https://doi.org/10.1016/S1001-7844\(12\)60030-3](https://doi.org/10.1016/S1001-7844(12)60030-3)

- ¹⁵⁸Anuja, M. N., Nithya, R. N., Rajamanickam, C., Madambath, I., Spermatotoxicity of a protein isolated from the root of *Achyranthes aspera*: a comparative study with gossypol, *Contraception*, **2010**, *82*, 385-390. DOI: 10.1016/j.contraception.2010.04.011
- ¹⁵⁹Anuja, M. M., Nithya R. S., Swathy, S. S., Rajamanickam, C., Indira, M., Spermicidal action of a protein isolated from ethanolic root extracts of *Achyranthes aspera*: an in vitro study, *Phytomedicine*, **2011**, *18*, 776-782. DOI:10.1016/j.phymed.2011.01.007
- ¹⁶⁰Sarwar, A., Khillare, B., Thakur, S. C., Suppression of Fertility in Male Rats by Hydroethanolic Extract of *Achyranthes Aspera*, *J. Appl. Pharm.*, **2014**, *6*, 156-165. DOI: 10.21065/19204159.6.3.135
- ¹⁶¹Dhale, D. A., Bhoi, S., Pharmacognostic Characterization and Phytochemical Screening of *Achyranthes Aspera* Linn, *Curr. Agric. Res. J.*, **2013**, *1*, 51-57. <http://dx.doi.org/10.12944/CARJ.1.1.07>
- ¹⁶²Rishikesh, M., Rahman, M., Goffar, R., Mamun, M. R., Dutta, P. R., Maruf, A., Phytochemical and Pharmacological Investigation of *Achyranthes Aspera* Linn., *Sch. Acad. J. Pharm.*, **2013**, *2*, 74-80. <http://saspublisher.com/wp-content/uploads/2013/03/SAJP2274-80.pdf>
- ¹⁶³Somagari, D. R., Basappa, K., Rolla, S., Jithendar, P., Rajashekar, B., Phytochemical investigation of seeds of *Achyranthes aspera* Linn., *J. Pharmacog. Phytochem.*, **2014**, *3*, 190-193. <http://www.phytojournal.com/archives/2014/vol3issue1/PartD/14.1-842.pdf>
- ¹⁶⁴Talreja, T., Sirohi, P., Sharma, T., Proximate Composition Analysis of Two Medicinally Important Plants *Achyranthes aspera* and *Cissus quadrangularis*, *Int. J. Pharm. Pharmaceut. Sci.*, **2015**, *7*, 416-418. <https://innovareacademics.in/journals/index.php/ijpps/article/view/3420>
- ¹⁶⁵Pawar, S.G., Kamble, V.M., Diversity of Bioactive Components of *Achyranthes aspera* Linn. leaves, *Asian J. Multidisc. Stud.*, **2016**, *4*, 49-52. <https://www.researchgate.net/publication/301559215>
- ¹⁶⁶Tiwari, P., Gond, P., Koshale, S., Phytochemical analysis of different parts of *Achyranthes Aspera*, *J. Pharmacog. Phytochem.*, **2018**, *7*, 60-62. <http://www.phytojournal.com/archives/2018/vol7issue2S/ParTB/SP-7-2-23-263.pdf>
- ¹⁶⁷Kamble, A., Phytochemical studies on *Achyranthes aspera*, *World Sci. News*, **2018**, *100*, 16-34. <http://www.worldscientificnews.com/wp-content/uploads/2018/04/WSN-100-2018-16-34.pdf>
- ¹⁶⁸Kushwaha, U., Evaluation of qualitative phytochemical analysis of water extract of *Achyranthes aspera* and *Achyranthes aspera* 30., *Indian J. Res. Homoeop.*, **2019**, *13*, 37-40. DOI:10.4103/ijrh.ijrh_30_18
- ¹⁶⁹Banerji, A., Chadha, M.S., Insect moulting hormone from *Achyranthes aspera*, *Phytochemistry*, **1970**, *9*, 1671. [https://doi.org/10.1016/S0031-9422\(00\)85295-4](https://doi.org/10.1016/S0031-9422(00)85295-4)
- ¹⁷⁰Misra, T. N., Singh, R. S., Pandey, H. S., Prasad, C., An aliphatic dihydroxyketone from *Achyranthes aspera*, *Phytochemistry*, **1991**, *30*, 2076-2078. [https://doi.org/10.1016/0031-9422\(91\)85071-7](https://doi.org/10.1016/0031-9422(91)85071-7)
- ¹⁷¹Misra, T. N., Singh, R. S., Pandey, H. S., Prasad, C., Singh, B. P., Two long chain compounds from *Achyranthes aspera*, *Phytochemistry*, **1993**, *33*, 221-223. [https://doi.org/10.1016/0031-9422\(93\)85427-S](https://doi.org/10.1016/0031-9422(93)85427-S)
- ¹⁷²Michl, G., Abebe, D., Bucar, F., Debella, A., Kunert, O., Schmid, M., Mulatu, E., Haslinger, E., New Triterpenoid Saponins from *Achyranthes aspera* LINN., *Helvetica*, **2000**, *83*, 359-363. DOI:10.1002/(SICI)15222675(20000216)83:2<359::AIDHLC A359>3.0.CO;2-7
- ¹⁷³Kunert, O., Haslinger, E., Schmid, M., Three Saponins, a Steroid, and a Flavanol Glycoside from *Achyranthes aspera*, *Monatsh. Chem.*, **2000**, *131*, 195-204. <https://doi.org/10.1007/PL00010306>
- ¹⁷⁴Chauhan, A. S., Rawat, G. S., Singh, C. P., Phytochemical study of *Achyranthes aspera* Linn, *Asian J. Chem.*, **2002**, *14*, 1059-1061. <https://www.researchgate.net/publication/295522483>
- ¹⁷⁵Mehta, F. A., Patel, B. G., Pandya, S. S., Ahir, K., Densitometric HPTLC Method for Analysis of Oleanolic Acid in *Achyranthes aspera* L., *J. Planar Chromat.*, **2010**, *23*, 289-292. <https://doi.org/10.1556/JPC.23.2010.4.10>
- ¹⁷⁶Ningwal, M., Chauhan, R., Phytochemical Screening of *Achyranthes aspera* Linn, *Int. J. Sci. Res.*, **2016**, *5*, 385-387. https://www.ijsr.net/get_abstract.php?paper_id=28021601
- ¹⁷⁷Varadharaj, V., Kuppan, M., Identification and determination of bioactive phytochemical constituents from the hydro-alcoholic extract of *Achyranthes aspera* whole plant by gas chromatography-mass spectrometry analysis, *Asian J. Pharmaceut. Clin. Res.*, **2015**, *8*, 125-129. <https://innovareacademics.in/journals/index.php/ajpcr/article/view/7858>
- ¹⁷⁸Sultana, S., Ali, M., Mir, S. R., Mittal, A., Analysis of Spectral Data of the Chemical Constituents from the Leaves of *Jasminum grandiflorum* L., *Achyranthes aspera* L. and *Tinospora cordifolia* (Willd.) Miers, *Euras. J. Anal. Chem.*, **2018**, *13*, Article 43, 9. <https://doi.org/10.29333/ejac/93438>
- ¹⁷⁹Meskel, G., Fiseha, A., Sesquiterpene and Ecdysteroid from *Achyranthes aspera* L., *Civ. Environ. Res.*, **2019**, *11*, 10-17. DOI: 10.7176/CER
- ¹⁸⁰Beula, S. J., Raj, V. B., Mathew, B., Isolation and molecular recognition of 6-prenyl apigenin towards MAO-A as the active principle of seeds of *Achyranthes aspera*, *Biomed. Prev. Nutr.*, **2014**, *4*, 379-382. DOI: 10.1016/j.bionut.2014.03.003
- ¹⁸¹Pandey, A., Kaushik, A., Evaluation of Anti-nociceptive and Anti-pyretic Activity of *Aerva pseudotomentosa* Leaves Aqueous Extract, *Int. J. Pharmaceut. Sci. Res.*, **2016**, *7*, 4686-4692. DOI: 10.13040/IJPSR.0975-8232.7(11).4686-92
- ¹⁸²Srivivan, K. R., Reddy, V. M., Antimicrobial studies on the leaves of *Earva Javanica*, *J. Pharmaceut. Allied Sci.*, **2008**, *5*, 5 pages. DOI: 10.4314/jophas.v5i1.48464
- ¹⁸³Sharif, A., Ahmed, E., Hussain, M. U., Malik, A., Ashraf, M., Antimicrobial constituents from *Aerva javanica*, *J. Chem. Soc. Pak.*, **2011**, *33*, 439-443. <https://www.researchgate.net/publication/285948285>
- ¹⁸⁴Srinivas, P., Reddy, S. R., Screening for antibacterial principle and activity of *Aerva javanica* (Burm.f) Juss. ex Schult, *Asian Pac. J. Trop. Biomed.*, **2012**, *2*, S838-S845. DOI:10.1016/S2221-1691(12)60321-9
- ¹⁸⁵Shad, A. A., Asmat, S., Bakht, J., Jan, S., Khan, M. A., Antimicrobial potentials and phytochemical analysis of desert cotton (*A. Javanica*) and flax (*L. Ustitatissimum*), *Pak. J. Pharm. Sci.*, **2016**, *29*(3), 861-868. <http://www.pjps.pk/wp-content/uploads/pdfs/29/3/Paper-14.pdf>
- ¹⁸⁶El-Shabasy, A. E., El -Gayar, K. E., Comparative studies between six taxa of Amaranthaceae based on their effects on some pathogenic bacterial isolates and morphological characters, *Egypt. J. Exp. Biol. (Bot.)*, **2019**, *15*, 341 - 351. DOI: 10.5455/egyjebb.20190914061823
- ¹⁸⁷Al-Shehri, M., Moustafa, M., Anticancer, Antibacterial, and Phytochemicals Derived from Extract of *Aerva javanica* (Burm.f.) Juss. ex Schult. Grown Naturally in Saudi Arabia, *Trop. Conserv. Sci.*, **2019**, *12*, 1-10. DOI: 10.1177/1940082919864262
- ¹⁸⁸Khalid, M., Bilal, M., Hassani, D., Iqbal, H. M., Huang, D., Antimicrobial, antioxidant, cytotoxicity and LC-MS analyses of *Aerva javanica*: an ethnomedicinally important plant, *J. Biol. Regul. Homeost. Agents*, **2017**, *31*, 963-969. <https://www.researchgate.net/publication/320011043>

- ¹⁸⁹Suleiman, M. H., Ethnobotanical, Phytochemical, and Biological Study of *Tamarix aphylla* and *Aerva javanica* Medicinal Plants Growing in the Asir Region, Saudi Arabia, *Trop. Conserv. Sci.*, **2019**, *12*, 1–14. DOI: 10.1177/1940082919869480
- ¹⁹⁰Eltayeb, N. M., Eltayeb, G. M., Salhimi, S. M., Anti-proliferative Effect of *Aerva javanica* Extracts on MCF7 and MDA-MB-231 Breast Cancer Cell Lines, *Malaysian J. Anal. Sci.*, **2017**, *21*, 1028-1035. <https://doi.org/10.17576/mjas-2017-2105-04>
- ¹⁹¹Kamalanathan, D., Natarajan, D., Anticancer potential of leaf and leaf-derived callus extracts of *Aerva javanica* against MCF-7 breast cancer cell line, *J. Cancer Res. Ther.*, **2018**, *14*, 321-327. DOI: 10.4103/0973-1482.171210
- ¹⁹²Reddy, K. S., Reddy, V. M., Antihyperglycaemic Activity of Ethanol Extract of *Aerva javanica* Leaves in Alloxan-Induced Diabetic Mice, *J. Pharm. Res.*, **2009**, *2*, 1259-1261. http://jprsolutions.info/article_detail.php?article_id=1405
- ¹⁹³Khunteta, A., Swarnkar, S. K., Gupta, M. K., Swarnkar, A., Sharma, S., Paliwal, S., Assessment of *In-Vivo* Antioxidant Potential of Hydro-Alcoholic Extract and Ethyl Acetate Fraction of *Aerva javanica* Linn. Flowering Tops, *Asian J. Pharmaceut. Res. Dev.*, **2019**, *7*, 72-78. DOI: 10.22270/ajprd.v7i6.614
- ¹⁹⁴Khan, A., Jan, S., Parveen, S., Khan, R. A., Saeed, S., Tanveer, A., Shad, A. A., Phytochemical analysis and Enzyme Inhibition Assay of *Aerva javanica* for Ulcer, *Chem. Cent. J.*, **2012**, *6*, Article 76, 7 pages. DOI: 10.1186/1752-153X-6-76
- ¹⁹⁵Ghanem, H. Z., Foda, D. S., Ibrahim, B., M., Evaluation of the anti-ulcer effect of *Aerva javanica* aerial parts against ethanol-induced gastric lesions in albino rats, *J. Innov. Pharmaceut. Biol. Sci.*, **2017**, *4*, 1-8. http://www.jipbs.com/VolumeArticles/FullTextPDF/302_JIPBSV4I301.pdf
- ¹⁹⁶Padala, K., Ragini, V., Anti Urolithiatic activity of Extracts of *Aerva javanica* in Rats, *Int. J. Drug Dev. Res.*, **2014**, *6*, 35-45. <https://www.researchgate.net/publication/287020652>
- ¹⁹⁷Joanofarc, J., Vamsadhara, C., Evaluation of Antidiarrhoeal Activity of *Aerva* species, *Nat. Prod. Sci.*, **2003**, *9*, 177-179. <https://www.researchgate.net/publication/293496301>
- ¹⁹⁸Arbab, A. H., Parvez, M. K., Al-Dosari, M. S., Al-Rehaily, A. J., Ibrahim, K. E., Alam, P., Alsaid, M. S., Rafatullah, S., Therapeutic efficacy of ethanolic extract of *Aerva javanica* aerial parts in the amelioration of CCl₄-induced hepatotoxicity and oxidative damage in rats, *Food Nutr. Res.*, **2016**, *60*, Article 30864, 10 pages. <http://dx.doi.org/10.3402/fnr.v60.30864>
- ¹⁹⁹Asija, R., Sharma, A., Kumar, P., Hepatoprotective activity of *Aerva javanica* Plant against Cyclophosphamide induced Hepatotoxicity, *Int. J. Pharmaceut. Erud.*, **2016**, *6*, 44-52. <http://www.pharmaerudition.org/ContentPaper/2016/may%202016%2044-52.pdf>
- ²⁰⁰Saleem, M., Musaddiq, S., Riaz, N., Zubair, M., Ashraf, M., Nasar, R., Jabbar, A., Ecdysteroids from the flowers of *Aerva javanica*, *Steroids*, **2013**, *78*, 1098-1102. DOI: 10.1016/j.steroids.2013.07.012
- ²⁰¹Iqbal, J., Zahra, S. T., Ahmad, M., Shah, A. N., Hassan, W., Herbicidal Potential of Dryland Plants on Growth and Tuber Sprouting in Purple Nutsedge (*Cyperus rotundus*), *Planta Daninha*, **2018**, *36*, Article e018170606, 8 pages. DOI: 10.1590/S0100-83582018360100001
- ²⁰²Mouhoub, F., Ouafi, S., Chabane, D., Acute and subacute oral toxicity assessment of gender of ahaggar's *Aerva javanica* in animal models, *Int. J. Pharmacol.*, **2018**, *14*, 640-651. DOI: 10.3923/ijp.2018.640.651
- ²⁰³Mussadiq, S., Riaz, N., Saleem, M., Ashraf, M., Ismail, T., Abdul Jabbar, A., New acylated flavonoid glycosides from flowers of *Aerva javanica*, *J. Asian Nat. Prod. Res.*, **2013**, *15*, 708–716. <http://dx.doi.org/10.1080/10286020.2013.795553>
- ²⁰⁴Mussadiq, S., Saeed, M. S., Riaz, N., Nazir, M., Khatoun, T., Abdul Jabbar, A., Aervfuranoside: A New Chlorinated Dibenzofuran Glycoside and other Metabolites from the Flowers of *Aerva javanica*, *J. Chem. Soc. Pak.*, **2017**, *39*, 572. <https://www.researchgate.net/publication/320518911>
- ²⁰⁵Samejo, M. Q., Memon, S., Bhangar, M. I., Khan, K. M., Chemical Compositions of the Essential oil of *Aerva javanica* Leaves and Stems, *Pak. J. Anal. Environ. Chem.*, **2012**, *13*, 48-52. <http://pjaec.pk/index.php/pjaec/article/view/190>
- ²⁰⁶Samejo, M. Q., Memon, S., Bhangar, M. I., Khan, K. M., Comparison of chemical composition of *Aerva javanica* essential oil obtained by different extraction methods, *Pak. J. Pharm. Sci.*, **2013**, *26*, 757-760. <https://pubmed.ncbi.nlm.nih.gov/23811454/>
- ²⁰⁷Abbas, N. M., El Imam, Y. M., Abdelmageed, M. A., The Phytochemical Analysis of the Ethanolic Extract of Sudanese *Aerva javanica* (Burm. f) Juss. ex Schult., *World J. Pharmaceut. Res.*, **2015**, *4*, 2253-2263. https://wjpr.net/dashboard/abstract_id/3043
- ²⁰⁸Karthishwaran, K., Kamalraj, S., Jayabaskaran, C., Kurup, S. S., Sakkir, S., Cheruth, A., GC-MS Assisted Phytoactive Chemical Compounds Identification and Profiling with Mineral Constituents from Biologically Active Extracts of *Aervajavanica* (Burm. f) Juss. ex Schult., *Not. Bot. Horti. Agrobo.*, **2018**, *46*, 517-524. DOI:10.15835/nbha46211037
- ²⁰⁹Hemmati, C., Nikooei, M., Bertaccini, A., Identification and transmission of phytoplasm and their impact on essential oil composition in *Aerva javanica*, *3 Biotech*, **2019**, *9*, 369-378. DOI: 10.1007/s13205-019-1843-0
- ²¹⁰Hossain, S., Rana, S., Islam, R., Islam, M., Phenolic Content Analysis and Evaluation of Antinociceptive, Antioxidant, Anti-Inflammatory Potential of *Alternanthera Pungens* Kunth, *J. Bangladesh Acad. Sci.*, **2018**, *42*, 129-136. DOI: 10.3329/jbas.v42i2.40040
- ²¹¹Jakhar, S., Dahiya, P., Antimicrobial, Antioxidant and Phytochemical Potential of *Alternanthera pungens* HB&K, *J. Pharm. Sci. Res.*, **2017**, *9*, 1305-1311. <https://www.jpsr.pharmainfo.in/Documents/Volumes/vol9Iss ue08/jpsr09081712.pdf>
- ²¹²Petrus, A., Seetharaman, T. R., Antioxidant flavone C-biosides from the aerial parts of *Alternanthera pungens*, *Indian J. Pharmaceut. Sci.*, **2005**, *67*, 187-193. <https://www.researchgate.net/publication/286959694>
- ²¹³Calderon, C. P., Garcia Aseff, S. B., Fuentes, L. B., Evaluation of diuretic activity of *Alternanthera pungens* extract in rats, *Phytother. Res.*, **1997**, *11*, 606-608. DOI:10.1002/(SICI)1099-1573(199712)11:8<606::AID-PTR163>3.0.CO;2-F
- ²¹⁴Garcia, S. B., Calderon, C. P., Fuentes, L.B., Preliminary evaluation of the gastrointestinal activity of *Alternanthera pungens*, *Fitoterapia*, **1995**, *66*, 324-327. <https://eurekamag.com/research/002/928/002928023.php>
- ²¹⁵Ramesh, P., Kumar, B., Tabu, K., Sree, M., Hyder, I., Parimi, U., Anti-helminthic Activity of *Alternanthera pungens* Krunth, *Int. J. ChemTech Res.*, **2019**, *12*, 214-218. <http://dx.doi.org/10.20902/IJCTR.2019.120426>
- ²¹⁶Mahmood, A., Tanveer, A., Nadeem, M. A., Zahir, Z. A., Comparative Allelopathic Potential of Two *Alternanthera* Species Against Germination and Seedling Growth of Rice, *Planta Daninha*, **2014**, *32*, 10 pages. <https://doi.org/10.1590/S0100-83582014000100001>
- ²¹⁷Kanagarasu, R., Bhavan, P. S., Rajkumar, G., Nathiya, V., Satgurunathan, T., Manjula, T., Phytochemical Characterization of *Alternanthera sessilis* and Assessment of its Growth Promoting Potential on the Freshwater Prawn *Macrobrachium rosenbergii*, *Int. J. Res. Stud. Zool.*, **2017**, *3*, 25-38. <http://dx.doi.org/10.20431/2454-941X.0304004>
- ²¹⁸Hossain, A. I., Faisal, M., Rahman, S., Jahan, R., Rahmatullah, M., A preliminary evaluation of antihyperglycemic and analgesic activity of *Alternanthera sessilis* aerial parts, *BMC Complement. Alternat. Med.*, **2014**, *14*, Article 169, 5 pages. DOI:10.1186/1472-6882-14-169

- ²¹⁹Mondal, H., Saha, S., Awang, K., Hossain, H., Ablat, A., Islam, K., Jahan, I. I., Sadhu, S. K., Hossain, G., Shilpi, J. A., Uddin, S. J., Central-stimulating and analgesic activity of the ethanolic extract of *Alternanthera sessilis* in mice. *BMC Complement. Alternat. Med.*, **2014**, *14*, Article 398, 9 pages. DOI:10.1186/1472-6882-14-398
- ²²⁰Rayees, S., Kumar, A., Rasool, S., Kaiser, P., Satti, N. K., Sangwan, P. L., Singh, S., Johri, R. K., Singh, G., Ethanolic Extract of *Alternanthera sessilis* (AS-1) Inhibits IgE-mediated Allergic Response in RBL-2H3 Cells, *Immunol. Invest.*, **2013**, *42*, 470-480. DOI: 10.3109/08820139.2013.789909
- ²²¹Saqib, F., Janbaz, K. H., Rationalizing ethnopharmacological uses of *Alternanthera sessilis*: A folk medicinal plant of Pakistan to manage diarrhea, asthma and hypertension, *J. Ethnopharmacol.*, **2016**, *182*, 110–121. <http://dx.doi.org/10.1016/j.jep.2016.02.017>
- ²²²Kumar, D. A., Das, M., Mohanraj, P., Das, A., Antimicrobial Activity Study of Ethanolic Extract of *Alternanthera sessilis* Linn. Aerial Parts, *J. App. Pharmaceut. Res.*, **2014**, *2*, 1-4. <https://www.semanticscholar.org/9a6b3b794fd481af97ecbf0c625a68a8037a1c7c>
- ²²³Monroy, A. E., Limsiaco, C. L., Phytochemical and Antimicrobial Analysis of “Lupo” (*Alternanthera sessilis* L.R.BR.), *WVSU Res. J.*, **2016**, *5*, 21-34. <https://ejournals.ph/article.php?id=13547>
- ²²⁴Jalalpure, S. S., Agrawal, N., Patil, M. B., Chimkode, R., Tripathi, A., Antimicrobial and wound healing activities of leaves of *Alternanthera sessilis* Linn, *Int. J. Green Pharm.*, **2008**, *2*, 141-144. <https://www.greenpharmacy.info/index.php/ijgp/article/view/18>
- ²²⁵Muniandy, K., Gothai, S., Tan, W. S., Kumar, S. S., Esa, N. M., Chandramohan, G., Al-Numair, K. S., Arulselvan, P., In Vitro Wound Healing Potential of Stem Extract of *Alternanthera sessilis*, *Evid. Based Complement. Alternat. Med.*, **2018**, Article 3142073, 13 pages. <https://doi.org/10.1155/2018/3142073>
- ²²⁶Sundar, R.D., Ravi, L., Mythili, S., Discovery of new anti-fungal phytochemical PDHC (propane-diyl-bis-hexahydroisochromene) isolated from *Alternanthera sessilis* leaves, *Int. J. Pharmaceut. Sci. Res.*, **2019**, *10*, 1148-1159. DOI:10.13040/IJPSR.0975-8232.10(3).1148-59
- ²²⁷Nayak, P., Nayak, S., Kar, D. M., Das, P., Pharmacological evaluation of ethanolic extracts of the plant *Alternanthera sessilis* against temperature regulation., *J. Pharm. Res.*, **2010**, *3*, 1381-1383. http://jpr solutions.info/article_detail.php?article_id=805
- ²²⁸Gothai, S., Muniandy, K., Mohd Esa, M. N., Subbiah, S. K., Arulselvan, P., Anticancer potential of *Alternanthera sessilis* extract on HT-29 human colon cancer cells. *Asian Pac. J. Trop. Biomed.*, **2018**, *8*, 394-402. DOI: 10.4103/2221-1691.239427
- ²²⁹Gupta, R., Singh, H. K., Antidepressant Like Effects of *Alternanthera sessilis* and *Clerodendrum infortunatum* Leaves Extract in Immobility Models, *Nat. Prod. J.*, **2014**, *4*, 33-37. DOI: 10.2174/2210315504666140515004826
- ²³⁰Tan, K. K., Kim, K. H., *Alternanthera sessilis* Red Ethyl Acetate Fraction Exhibits Antidiabetic Potential on Obese Type 2 Diabetic Rats, *Evid. Based Complement. Alternat. Med.*, **2013**, Article 845172, 8 pages. <http://dx.doi.org/10.1155/2013/845172>
- ²³¹Chai, T. T., Khoo, C. S., Tee, C. S., Wong, F. C., Alpha-glucosidase inhibitory and antioxidant potential of antidiabetic herb *Alternanthera sessilis*: Comparative analyses of leaf and callus solvent fractions, *Pharmacogn. Mag.*, **2016**, *12*, 253-258. DOI:10.4103/0973-1296.192202
- ²³²Azizah, O., Amin, I., Fouad, A. R., Antioxidant properties of *Alternanthera sessilis* red and green, *Acta Hort.*, **2015**, *1106*, 131-136. DOI: 10.17660/ActaHortic.2015.1106.20
- ²³³Khan, M. S., Yusufzai, S. K., Kaun, L. P., Shah, M. D., Idris, R., Chemical composition and antioxidant activity of essential oil of leaves and flowers of *Alternanthera sessilis* red from Sabah, *J. Appl. Pharmaceut. Sci.*, **2016**, *6*, 157-161. DOI: 10.7324/JAPS.2016.601222
- ²³⁴Khan, M. S., Yusufzai, S. K., Ying, L. Y., Zulfashriq, W., GC-MS Based Chemical Profiling and Evaluation of Antioxidant Potential of Leaves and Stems of *Alternanthera Sessilis* Red from Sabah, Malaysia, *Int. J. Pharm. Pharmaceut. Sci.*, **2018**, *10*, 4-9. DOI: 10.22159/ijpps.2018v10i7.25204
- ²³⁵Lin, S. C., Lin, Y. H., Shyuu, S. J., Lin, C. C., Hepatoprotective effects of Taiwan folk medicine: *Alternanthera sessilis* on liver damage induced by various hepatotoxins, *Phytother. Res.*, **1994**, *8*, 391-398. DOI:10.1002/ptr.2650080703
- ²³⁶Biman Bhuyan, B., Baishya, K., Rajak, P., Effects of *Alternanthera sessilis* on Liver Function in Carbon Tetrachloride Induced Hepatotoxicity in Wister Rat Model, *Indian J. Clin. Biochem.*, **2018**, *33*, 190–195. DOI: 10.1007/s12291-017-0666-1
- ²³⁷Niraimathi, K. L., Sudha, V., Lavanya, R., Brindha, P., Biosynthesis of silver nanoparticles using *Alternanthera sessilis* (Linn.) extract and their antimicrobial, antioxidant activities, *Colloid. Surf. B*, **2013**, *102*, 288-291. DOI:10.1016/j.colsurfb.2012.08.041
- ²³⁸Subhalakshmi, A., Rajagopal, K., Nithyanandhan, G., Devi, J., Karthikeyan, S., Chromium phytoaccumulation by *alternanthera sessilis* in vitro-a first report, *Int. J. Pharm. Biosci.*, **2013**, *4*, 618–624. <https://www.researchgate.net/publication/250612318>
- ²³⁹Gajbhihiye, S. P., Bhalerao, S. A., Phytoremediation potential of *Alternanthera sessilis* L. growing in industrially contaminated environment, *Int. J. Curr. Res.*, **2017**, *9*, 44550-44556. www.journalcra.com/article/2017.9.1.44550-44556
- ²⁴⁰Abbasi, S. A., Abbasi, T., Ponni, G., Tauseef, S. M., Potential of joyweed *Alternanthera sessilis* for rapid treatment of domestic sewage in SHEFROL® bioreactor, *Int. J. Phytoremed.*, **2019**, *21*, 160-169. DOI:10.1080/15226514.2018.1488814
- ²⁴¹Salimizadeh, M., Shirvani, M., Shariatmadari, H., Mortazavi, M. S., Bentonite addition to a PCB-contaminated sandy soil improved the growth and phytoremediation efficiency of *Zea mays* L. and *Alternanthera sessilis* L., *Int. J. Phytoremed.*, **2020**, *22*, 176-183. DOI: 10.1080/15226514.2019.1652564
- ²⁴²Chinmayee, M. D., Sheeba, A. M., Mini, I., Swapna, T.S., Phytoremediation potential and antioxidant responses in *Alternanthera sessilis* (L.) r.Br., *J. Aquat. Biol. Fish.*, **2014**, *2*, 114-118. <https://pdfs.semanticscholar.org/55aa/73f54f3fed5ce101f993f755777278306b3b.pdf>
- ²⁴³Lalitha Sree, T., Vijayalakshmi, K., Evaluation of vitamins and antinutrients in the leaves of traditional medicinal plant *Alternanthera Sessilis* (L.) R.Br.Ex DC, *Int. J. Health Sci. Res.*, **2018**, *8*, 244-253. https://www.ijhsr.org/IJHSR_Vol.8_Issue.10_Oct2018/36.pdf
- ²⁴⁴Penders, A., Kapundu, M., Delaude, C., Structure du Principal Saponoside de *Alternanthera sessilis*, *Bull. Soc. Chim. Belges*, **1992**, *101*, 227-232. DOI:10.1002/bscb.19921010310
- ²⁴⁵Consolacion, Y. R., Tremor, N., John, A. R., Ionone derivatives from *Alternanthera sessilis*, *J. Asian Nat. Prod. Res.*, **2002**, *4*, 109-115. DOI: 10.1080/10286020290027380
- ²⁴⁶Kini, S. G., Nguyen, P. Q., Weissbach, S., Mallagaray, A., Shin, J., Yoon, H. S., Tam, J. P., Studies on the Chitin Binding Property of Novel Cysteine-Rich Peptides from *Alternanthera sessilis*, *Biochemistry*, **2015**, *54*, 6639-6649. DOI:10.1021/acs.biochem.5b00872
- ²⁴⁷Anitha, R., Kanimozhi, S., Pharmacognostic Evaluation of *Alternanthera Sessilis* (L.) R.Br.ex.DC, *Pharmacog. J.*, **2012**, *4*, 31-34. DOI:10.5530/pj.2012.28.6

- ²⁴⁸Dhanya, V. T., Jollykutty, E., Pharmacognostic and Preliminary Physico-Chemical Evaluation of Matsyakshi [*Alternanthera Sessilis* (Linn.) R.Br], *Int. J. Ayurv. Pharm. Res.*, **2017**, *5*, 16-22. <https://ijapr.in/index.php/ijapr/article/view/742>
- ²⁴⁹Shreshtha, S., Anushi, J., Ambika, J., Nitesh, J., Anupma, H., Study of total phenol, flavonoid contents and phytochemical screening of methanolic crude extracts of two weed plants, *Ann. Plant Sci.*, **2017**, *6*, 1645-1648. DOI: 10.21746/1651-1654
- ²⁵⁰Umate, S. K., Marathe, V. R., Proximate and Phytochemical screening of *Alternanthera sessilis* (L.) R. Br. ex DC. (Amaranthaceae) – An underutilised leafy vegetable, *Int. J. Life Sci.*, **2017**, *5*, 703-705. <http://www.ijlsci.in/abstract-5-4-31>
- ²⁵¹Gupta, R., Singh, H. K., Detection and quantitation of gallic acid in *Alternanthera sessilis* and *Clerodendrum infortunatum* by HPTLC, *Pharm. Pharmacol. Int. J.*, **2016**, *4*, 467-471. DOI: 10.15406/ppij.2016.04.00097
- ²⁵²Das, M., Kumar, D. A., Deb, J., Rao, D. S., Isolation and Characterization of Flavonoid from Ethanolic extract of *Alternanthera sessilis* Linn., *J. Biomed. Pharmaceut. Res.*, **2017**, *6*, 23-31. <https://jbpr.in/index.php/jbpr/article/view/82>
- ²⁵³Kumar, H. K., Raju, M. B., Acharyya, S., Mishra, B., Isolation of Phytoconstituents from the Roots of *Bambusa arundinacea*, *Eur. J. Biomed. Pharmaceut. Sci.*, **2016**, *3*, 343-347. https://www.ejbps.com/ejbps/abstract_id/812
- ²⁵⁴Vani, M., Rahaman, S. K., Rani, A. P., Detection and Quantification of Major Phytochemical Markers for Standardization of Talinum *Portulacifolium*, *Gomphrena serrata*, *Alternanthera sessilis* and *Euphorbia heterophylla* by HPLC, *Pharmacog. J.*, **2018**, *10*, 439-46. DOI: 10.5530/pj.2018.3.72
- ²⁵⁵Amini, R., Namdari, T., Inhibitory effects of prostrate amaranth (*Amaranthus blitoides* S. Wats) on common bean cultivars, *Allelopath. J.*, **2013**, *32*, 63-76. <https://www.researchgate.net/publication/259997108>
- ²⁵⁶Sarker, U., Oba, S., Nutrients, minerals, pigments, phytochemicals, and radical scavenging activity in *Amaranthus blitum* leafy vegetables, *Sci. Rep.*, **2020**, *10*, Article 3836, 9 pages. <https://doi.org/10.1038/s41598-020-59848-w>
- ²⁵⁷Allemann, J., Denner, F. D., Allelopathic influence of soils planted to *Amaranthus cruentus* L. on two tomato (*Lycopersicon esculentum* Mill.) cultivars, *S. Afr. J. Plant Soil*, **2006**, *23*, 142-143. DOI: 10.1080/02571862.2006.10634745
- ²⁵⁸Allemann, I., Cawood, M. E., Allemann, J., Influence of altered temperatures on allelopathic properties of *Amaranthus cruentus* L., *Acta Agric. Slov.*, **2017**, *109*, 465-471. DOI:10.14720/aas.2017.109.2.29
- ²⁵⁹Pasko, P., Barton, H., Fořta, M., Gwizdz, J., Evaluation of antioxidant activity of amaranth (*Amaranthus cruentus*) grain and by-products (flour, popping, cereal), *Rocz. Panstw. Zakl. Hig.*, **2007**, *58*, 35-40. <https://pubmed.ncbi.nlm.nih.gov/17711088/>
- ²⁶⁰Pasko, P., Barton, H., Zagrodzki, P., Chlopicka, J., Izewska, A., Gawlik, M., Gawlik, M., Gorinstein, S., Effect of amaranth seeds in diet on oxidative status in plasma and selected tissues of high fructose-fed rats, *Food Chem.*, **2011**, *126*, 85-90. DOI:10.1016/j.foodchem.2010.10.081
- ²⁶¹Nana, F. W., Hilou, A., Millogo, J. F., Nacoulma, O. G., Phytochemical Composition, Antioxidant and Xanthine Oxidase Inhibitory Activities of *Amaranthus cruentus* L. and *Amaranthus hybridus* L. Extracts, *Pharmaceuticals*, **2012**, *5*, 613-628. DOI: 10.3390/ph5060613
- ²⁶²Dandeu, L. N., Repollo, R., Oyhenart, J. A., Ramirez, M. R., Preliminary Analysis of the Hemagglutinating Activity of Seed Extracts of the Amaranth Genus, *Int. J. Adv. Res.*, **2018**, *6*, 515-519. DOI:10.21474/IJAR01/7240
- ²⁶³Pandey, S., Ganeshpurkar, A., Bansal, D., Dubey, N., Hematopoietic Effect of *Amaranthus cruentus* Extract on Phenylhydrazine-Induced Toxicity in Rats, *J. Diet Suppl.*, **2016**, *13*, 607-615. DOI: 10.3109/19390211.2016.1155685
- ²⁶⁴Pandey, S., Ganeshpurkar, A., Dubey, N., Hematopoietic Effect of *Amaranthus cruentus* Extract on Cyclophosphamide Induced Toxicity in Rats, *Pharmacog. Commn.*, **2019**, *9*, 139-142. DOI:10.5530/pc.2019.4.29
- ²⁶⁵Soares, R. A., Mendonca, S., Castro, L. I., Menezes, A. C., Areas, J. A., Major Peptides from Amaranth (*Amaranthus cruentus*) Protein Inhibit HMG-CoA Reductase Activity, *Int. J. Mol. Sci.*, **2015**, *16*, 4150-4160. DOI:10.3390/ijms16024150
- ²⁶⁶Aguilar, E. G., Cantarelli, M. A., Marchevsky, E. J., Escudero, N. L., Camina, J. M., Study and Determination of As, Cr, and Pb in Amaranth Seeds, *J. Chem.*, **2013**, Article 138391, 4 pages. <http://dx.doi.org/10.1155/2013/138391>
- ²⁶⁷Escudero, N. L., De Arellano, M. L., Luco, J. M., Gimenez, M. S., Mucciarelli, S. I., Comparison of the Chemical Composition and Nutritional Value of *Amaranthus cruentus* Flour and Its Protein Concentrate, *Plant Foods Hum. Nutr.*, **2004**, *59*, 15-21. DOI: 10.1007/s11130-004-0033-3
- ²⁶⁸Oleszek, W., Junkuszew, M., Stochmal, A., Determination and toxicity of saponins from *Amaranthus cruentus* seeds, *J. Agric. Food Chem.*, **1999**, *47*, 3685-3687. DOI:10.1021/jf990182k
- ²⁶⁹Mlakar, S. G., Jakop, M., Bavec, M., Bavec, F., Allelopathic effects of *Amaranthus retroflexus* and *Amaranthus cruentus* extracts on germination of garden cress, *Afr. J. Agric. Res.*, **2012**, *7*, 1492-1497. DOI: 10.5897/AJAR11.1145
- ²⁷⁰Strack, D., Leicht, P., Bokern, M., Wray, V., Grotjahn, L., Hydroxycinnamic acid esters of isocitric acid: Accumulation and enzymatic synthesis in *Amaranthus cruentus*, *Phytochemistry*, **1987**, *26*, 2919-2922. DOI: 10.1016/S0031-9422(00)84563-X
- ²⁷¹Minzanova, S. T., Mironov, V. F., Tsepaveva, O. V., Mironova, L. G., Vyshtakalyuk, A. B., Mindubaev, A. Z., Nizameev, I. R., Kholin, K. V., Pashagin, A. V., Isolation and Structural and Chemical Analysis of Pectinic Polysaccharides from *Amaranthus cruentus*, *Chem. Nat. Compd.*, **2014**, *50*, 54-59. <https://doi.org/10.1007/s10600-014-0865-4>
- ²⁷²Cai, Y., Sun, M., Corke, H., HPLC Characterization of Betalains from Plants in the Amaranthaceae, *J. Chromatogr. Sci.*, **2005**, *43*, 454-460. DOI: 10.1093/chromsci/43.9.454
- ²⁷³Pisarikova, B., Peterka, J., Trckova, M., Moudry, J., Zraly, Z., Herzig, I., The content of insoluble fibre and crude protein value of the above ground biomass of *Amaranthus cruentus* and *A. hypochondriacus*, *Czech J. Anim. Sci.*, **2007**, *52*, 348-353. DOI: 10.17221/2339-CJAS
- ²⁷⁴Pasko, P., Sajewicz, M., Gorinstein, S., Zachwieja, Z., Analysis of Selected Phenolic Acids and Flavonoids in *Amaranthus cruentus* and *Chenopodium quinoa* Seeds and Sprouts by HPLC, *Acta Chromatogr.*, **2008**, *20*, 661-672. DOI: 10.1556/AChrom.20.2008.4.11
- ²⁷⁵Sporna-Kucab, A., Kumorkiewicz, A., Szmyr, N., Szneler, E., Wybraniec, S., Separation of betacyanins from flowers of *Amaranthus cruentus* L. in a polar solvent system by high-speed counter-current chromatography, *J. Sep. Sci.*, **2019**, *42*, 1676-1685. DOI: 10.1002/jssc.201801172
- ²⁷⁶Ishtiaq, S., Ali, T., Ahmad, B., Anwar, F., Afridi, M. S., Humera Shaheen, H., Phytochemical and Biological Evaluations of Methanolic Extract of *Amaranthus graecizans* subsp. *silvestris* (Vill.) Brenan, *Br. J. Pharmaceut. Res.*, **2017**, *15*, Article BJPR.31698, 11 pages. DOI: 10.9734/BJPR/2017/31698
- ²⁷⁷Ishtiaq, S., Ahmad, M., Hanif, U., Akbar, S., Mehjabeen, M., Kamran, S. H., Phytochemical and in vitro antioxidant evaluation of different fractions of *Amaranthus graecizans* subsp. *silvestris* (Vill.) Brenan, *Asian Pac. J. Trop. Med.*, **2014**, *7*, S342-S347. DOI: 10.1016/S1995-7645(14)60256-X

- ²⁷⁸Ishtiaq, S., Afridi, M. S., Masood, N., Amelioration of isoniazid and rifampicin-induced liver toxicity by *Amaranthus graecizans* subsp. *silvestris* in rat, *Bangladesh J. Pharmacol.*, **2017**, *12*, 354-358. DOI: 10.3329/bjp.v12i3.32985
- ²⁷⁹Kamil, M., El Sheikh, M. O., Ahmad, F., Al Katheeri, B. A., Al Rihawi, F., Naji, M. A., Pharmacognostic and Phytochemical Studies on *Amaranthus graecizans* L., *Hamdard Med.*, **2012**, *55*, 62-66.
https://applications.emro.who.int/imemrf/Hamdard_Med/Hamdard_Med_2013_55_3_62_66.pdf
- ²⁸⁰Escudero, N. L., Albarracin, G., Fernández, S., De Arellano, L. M., Mucciarelli, S., Nutrient and antinutrient composition of *Amaranthus muricatus*, *Plant Foods Hum. Nutr.*, **1999**, *54*, 327-336. DOI:10.1023/a:1008149721435
- ²⁸¹De Ruiz, R. E., Fusco, M. D., Sosa, A., Ruiz, S. O., Isolation of flavonoids and anthraquinones of *Amaranthus muricatus* (Moquin) ex Hicken (Amaranthaceae), *Acta Farm. Bonaerense*, **2001**, *20*, 9-12.
<https://www.researchgate.net/publication/294630637>
- ²⁸²De Ruiz, R. E., Silva, R. A., Ruiz, S. O., Isolation of steroids, ammonium bases and saponines of *Amaranthus muricatus* (Moquin) Gillies ex Hicken (Amaranthaceae), *Acta Farm. Bonaerense*, **2003**, *22*, 101-104.
<https://www.researchgate.net/publication/289131908>
- ²⁸³Bradow, J. M., Connick, W. J., Allelochemicals from palmer amaranth, *Amaranthus palmeri* S. Wats., *J. Chem. Ecol.*, **1987**, *13*, 185-202. DOI:10.1007/BF01020362
- ²⁸⁴Connick, W. J., Bradow, J. M., Legendre, M. G., Vail, S. L., Menges, R. M., Identification of volatile allelochemicals from *Amaranthus palmeri* S. Wats., *J. Chem. Ecol.*, **1987**, *13*, 463-472. DOI: 10.1007/BF01880093
- ²⁸⁵Landa-Pineda, C. M., Arroyo-Becerra, A., Rosas-Alvarado, A., Teran, L. M., Garcia-Cruz, M. L., Marchat, L. A., Reyes-Lopez, C. A., Major allergen from *Amaranthus palmeri* pollen is a profilin: Isolation, partial characterisation and IgE recognition., *Allergol. Immunopathol.*, **2015**, *44*, 160-166.
<http://dx.doi.org/10.1016/j.aller.2015.05.002>
- ²⁸⁶Douglas, K., Evaluation of Phytochemicals and Lead Ions Concentration in *Amaranthus palmeri* and *Brassicaoleraceae* *Acephala* Plants Extracts, *Eur. J. Med. Plants*, **2016**, *16*, Article EJMP.28656, 7 pages. DOI: 10.9734/EJMP/2016/28656
- ²⁸⁷Shahrokhi, S., Hejazi, S. N., Khodabandeh, H., Farboodi, M., Faramarzi, A., Allelopathic Effect of Aqueous Extracts of Pigweed, *Amaranthus retroflexus* L. Organs on Germination and Growth of Five Barley Cultivars, *Int. Proceed. Chem. Biol. Environ. Engin.*, **2011**, *20*, 80-84.
<http://www.ipcbee.com/vol20/15-ICBEE2011E10020.pdf>
- ²⁸⁸Bakhshayeshan-Agdam, H., Salehi-Lisar, S. Y., Motafakkerzad, R., Talebpour, A., Farsadi, N., Allelopathic effects of redroot pigweed (*Amaranthus retroflexus* L.) on germination & growth of cucumber, alfalfa, common bean and bread wheat, *Acta Agric. Slov.*, **2015**, *105*, 193-202. DOI: 10.14720/aas.2015.105.2.02
- ²⁸⁹Bakhshayeshan-Agdam, H., Salehi-Lisar, S. Y., Motafakkerzad, R., Talebpour, A., Farsadi, N., Allelopathic Effects of Different Organs of Redroot Pigweed (*Amaranthus retroflexus* L.) on Cucumber and Wheat Plants, *J. Agric. Sci. Technol.*, **2015**, *5*, 678-685. DOI: 10.17265/2161-6256/2015.08.006
- ²⁹⁰Szabo, R., Nadasy, E., Pasztor, G., Study on the allelopathic effect of *Amaranthus retroflexus* L., *Datura stramonium* L. and *Panicum miliaceum* L. on the germination of maize, *Julius-KuhnArchiv.*, **2018**, *458*, 458-467. DOI: 10.5073/jka.2018.458.068
- ²⁹¹Tehrani, M., Sankian, M., Assarehzadegan, M. A., Falak, R., Jabbari, F., Varasteh, A., Immunochemical Characterization of *Amaranthus retroflexus* Pollen Extract: Extensive Cross-reactive Allergenic Components among the Four Species of Amaranthaceae/Chenopodiaceae, *Iran J. Allergy Asthma Immunol.*, **2010**, *9*, 87-95.
<https://pubmed.ncbi.nlm.nih.gov/20683102/>
- ²⁹²Vakili, M. M., Fallahpour, M., Mohammadi, M., Rasi, F. S., Mokhtarian, K., Khoshmirsafa, M., Jafari, R., Shirzad, N., Falak, R., Identification of polcalcin as a novel allergen of *Amaranthus retroflexus* pollen, *Allergol. Immunopathol.*, **2019**, *47*, 357-364. DOI:10.1016/j.aller.2018
- ²⁹³Marinas, I. C., Chifiriuc, C., Oprea, E., Lazar, V., Antimicrobial and antioxidant activities of alcoholic extracts obtained from vegetative organs of *A. retroflexus*, *Roum. Arch. Microbiol. Immunol.*, **2014**, *73*, 35-42.
<https://pubmed.ncbi.nlm.nih.gov/25518569>
- ²⁹⁴Pandit, J., Setty, S. R., Antioxidant activities of *Amaranthus retroflexus* leaves, *Int. J. Pharm. Drug Anal.*, **2015**, *3*, 223-225. <http://www.ijpda.com/admin/uploads/IO9Wsj.pdf>
- ²⁹⁵Pacifico, S., D'Abrosca, B., Golino, A., Mastellone, C., Piccolella, S., Fiorentino, A., Monaco, P., Antioxidant evaluation of polyhydroxylated nerolidols from redroot pigweed (*Amaranthus retroflexus*) leaves, *LWT Food Sci. Technol.*, **2008**, *41*, 1665-1671. DOI:10.1016/j.lwt.2007.10.006
- ²⁹⁶Khoramnejadian, S., Saeb, K., Accumulation and Translocation of Heavy Metals by *Amaranthus Retroflexus*, *J. Earth Environ. Health Sci.*, **2015**, *1*, 58-60. DOI:10.4103/2423-7752.170581
- ²⁹⁷Abdelrahman, H., Chelate Induced Redistribution of Pb and Zn Fractions in Contaminated Soils and Implications on Phytoremediation, *Egypt. J. Soil Sci.*, **2019**, *59*, 145-155. DOI: 10.21608/ejss.2019.11501.1262
- ²⁹⁸Bahrami-Teimoori, B., Nikparast, Y., Hojatianfar, M., Akhlaghi, M., Ghorbani, R., Pourianfar, H. R., Characterisation and antifungal activity of silver nanoparticles biologically synthesised by *Amaranthus retroflexus* leaf extract, *J. Exp. Nanosci.*, **2017**, *12*, 129-139. DOI:10.1080/17458080.2017.1279355
- ²⁹⁹Salar Amoli, J., Sadighara, P., Barin, A., Yazdani, A., Satari, S., Biological screening of *Amaranthus retroflexus* L. (Amaranthaceae), *Braz. J. Pharmacog.*, **2009**, *19*, 617-620. DOI:10.1590/S0102-695X2009000400019
- ³⁰⁰Weston, P. A., Gurusinge, S., Birkhead, E., Skoneczny, D., Quinn, J. C., Weston, L. A., Chemometric analysis of *Amaranthus retroflexus* in relation to livestock toxicity in southern Australia, *Phytochemistry*, **2019**, *161*, 1-10.
<https://doi.org/10.1016/j.phytochem.2019.01.016>
- ³⁰¹D'Abrosca, B., De Maria, P., DellaGreca, M., Fiorentino, A., Golino, A., Izzo, A., Monaco, P., Amarantholidols and amarantholidosides: new nerolidol derivatives from the weed *Amaranthus retroflexus*, *Tetrahedron*, **2006**, *62*, 640-646. DOI:10.1016/j.tet.2005.10.007
- ³⁰²Fiorentino, A., DellaGreca, M., D'Abrosca, B., Golino, A., Pacifico, S., Izzo, A., Monaco, P., Unusual sesquiterpene glucosides from *Amaranthus retroflexus*, *Tetrahedron*, **2006**, *62*, 8952-8958. DOI:10.1016/j.tet.2006.07.017
- ³⁰³Qian, G., Wang, Z., Zhang, L., Xu, F., Wang, B., Li, D., Chen, Y., Chemical Compositions of *Amaranthus retroflexus*, *Chem. Nat. Compd.*, **2016**, *52*, 982-985.
<https://doi.org/10.1007/s10600-016-1841-y>
- ³⁰⁴Flath, R. A., Altieri, M. A., Mon, T. R., Volatile constituents of *Amaranthus retroflexus* L., *J. Agric. Food Chem.*, **1984**, *32*, 92-94. DOI:10.1021/jf00121a024

- ³⁰⁵Soshnikova, O. V., Yatsyuk, V. Y., Yatsuk, V. Y., Soshnikova, O. V., The investigations of the chemical composition of *Amaranthus retroflexus* L., *I. P. Pavlov Russ. Med. Biol. Herald.*, **2010**, *18*, 135-141. DOI: 10.17816/PAVLOVJ20102135-141
- ³⁰⁶Sarkar, E., Chakraborty, P., Allelopathic effect of *Amaranthus spinosus* Linn. on growth of rice and mustard, *J. Trop. Agric.*, **2015**, *53*, 139-148. <https://www.researchgate.net/publication/300053371>
- ³⁰⁷Singh, A. B., Dahiya, P., Antigenic and allergenic properties of *Amaranthus spinosus* pollen-a commonly growing weed in India, *Ann. Agric. Environ. Med.*, **2002**, *9*, 147-151. http://www.aem.pl/72785_0_2.html
- ³⁰⁸Taiab, J., Nazmul, Q., Asif, A. M., Amran, H., Shams, K. M., Apurba, S. A., Analgesic activity of extracts of the whole plant of *amaranthus spinosus* Linn, *Int. J. Drug Dev. Res.*, **2011**, *3*, 189-193. <https://www.researchgate.net/publication/292593959>
- ³⁰⁹Zeashan, H., Amresh, G., Rao, C. V., Singh S., Antinociceptive activity of *Amaranthus spinosus* in experimental animals, *J. Ethnopharmacol.*, **2009**, *122*, 492-496. DOI:10.1016/j.jep.2009.01.031
- ³¹⁰Rjeibi, I., Ben Saad, A., Sdayria, J., Feriani, A., Ncib, S., Allagui, M.S., Hfaiedh, N., Souid, S., HPLC-DAD identification of polyphenols from ethyl acetate extract of *Amaranthus spinosus* leaves and determination of their antioxidant and antinociceptive effects, *Inflammopharmacol.*, **2019**, *27*, 975-984. <https://doi.org/10.1007/s10787-018-0482-0>
- ³¹¹Amabye, T. G., Evaluation of Physicochemical, Phytochemical, Antioxidant and Antimicrobial Screening Parameters of *Amaranthus spinosus* Leaves, *Nat. Prod. Chem. Res.*, **2015**, *3*, Article 199, 5 pages. DOI:10.4172/2329-6836.1000199
- ³¹²Majumdar, J., Boro, S., Bharadwaj, B., Unni, B., Phytochemical Analysis and Isolation of Compounds from Methanolic Extract of *Amaranthus spinosus* and its Antimicrobial Effect Against *Staphylococcus aureus*, *World J. Pharmaceut. Life Sci.*, **2017**, *3*, 100-103. https://www.wjpls.org/home/article_abstract/638
- ³¹³Khanal, D., Raut, B., Dangol, K. S., Phytochemical Screening, Pharmacognostic Evaluation and Biological Activity of *Amaranthus spinosus* L., *J. Manmohan Mem. Inst. Health Sci.*, **2015**, *1*, 29-34. DOI: 10.3126/jmmihs.v1i4.11999
- ³¹⁴Jadhav, V., Biradar, S. D., Evaluation of Antifungal Activity of *Amaranthus spinosus* L. (Amaranthaceae), *Int. J. Curr. Microbiol. Appl. Sci.*, **2016**, *5*, 38-43. <http://dx.doi.org/10.20546/ijemas.2016.509.005>
- ³¹⁵Yusnawan, E., Inayati, A., Antifungal activity of crude extracts of *Ageratum conyzoides*, *Cyperus rotundus*, and *Amaranthus spinosus* against rust disease, *AGRIVITA J. Agric. Sci.*, **2018**, *41*, 403-414. DOI:10.17503/agrivita.v40i0.1889
- ³¹⁶Mondal, A., Guria, T., Maity, T. K., Bishayee, A., A Novel Tetraenoic Fatty Acid isolated from *Amaranthus spinosus* inhibits proliferation and Induces apoptosis of Human Liver Cancer Cells, *Int. J. Mol. Sci.*, **2016**, *17*, 1604-1612. DOI:10.3390/ijms17101604
- ³¹⁷Mondal, A., Guria, T., Maity, T. K., A new ester of fatty acid from a methanol extract of the whole plant of *Amaranthus spinosus* and its α -glucosidase inhibitory activity, *Pharmaceut. Biol.*, **2015**, *53*, 600-604. DOI:10.3109/13880209.2014.935863
- ³¹⁸Olajide, O. A., Ogunleye, B. R., Erinle, T. O., Anti-inflammatory Properties of *Amaranthus spinosus* Leaf Extract, *Pharmaceut. Biol.*, **2004**, *42*, 521-525, DOI: 10.3109/13880200490893285
- ³¹⁹Barku, V. Y., Opoku-Boahen, Y., Owusu-Ansah, E., Mensah, E. F., Antioxidant activity and the estimation of total phenolic and flavonoid contents of the root extract of *Amaranthus spinosus*, *Asian J. Plant Sci. Res.*, **2013**, *3*, 69-74. <https://www.researchgate.net/publication/235339319>
- ³²⁰Cherian, P., Sheela, D., Antioxidant activity of ethyl acetate extract of genus *Amaranthus* Linn., *Int. J. Pharm. Sci. Res.*, **2017**, *8*, 5199-5204. DOI:10.13040/IJPSR.0975-8232.8(12).5199-04
- ³²¹Kumar, B. S., Lakshman, K., Jayaveera, K. N., Shekar, D. S., Kumar, A. A., Manoj, B., Antioxidant and antipyretic properties of methanolic extract of *Amaranthus spinosus* leaves, *Asian Pac. J. Trop. Med.*, **2010**, *3*, 702-706. [https://doi.org/10.1016/S1995-7645\(10\)60169-1](https://doi.org/10.1016/S1995-7645(10)60169-1)
- ³²²Rjeibi, I., Ben Saad, A., Ncib, S., Souid, S., Alimi, H., Characterization of *Amaranthus spinosus* collected from different regions: Phytochemical and biological properties, *J. Food Biochem.*, **2017**, *41*, Article e12397, 10 pages. <https://doi.org/10.1111/jfbc.12397>
- ³²³Abid, M., Kumar Gosh, A., Ali Khan, N., In Vivo Psychopharmacological Investigation of *Delphinium denudatum* and *Amaranthus spinosus* Extracts on Wistar Rats, *Basic Clin. Neurosci.*, **2017**, *8*, 503-512. DOI:10.29252/NIRP.BCN.8.6.503
- ³²⁴Ashok Kumar, B. S., Lakshman, K., Velmurugan, C., Sridhar, S. M., Gopisetty, S., Antidepressant Activity of Methanolic Extract of *Amaranthus spinosus*, *Basic Clin. Neurosci.*, **2014**, *5*, 11-17. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4202599>
- ³²⁵Faponle, A. S., Attunise, A. K., Adegbesan, B. O., Ogunlabi, O. O., Odufuwa, K. T., Ajani, E. O., Separate and co-administration of *Amaranthus spinosus* and vitamin C modulates cardiovascular disease risk in high fat diet-fed experimental rats, *J. Pharmacog. Phytother.*, **2015**, *7*, 27-34. DOI: 10.5897/JPP2014.0340
- ³²⁶Satish, S. B., Yogesh, H. W., Effect of aqueous extract of *Amaranthus spinosus* on hematological parameters of wistar albino rats, *J. Exp. Biol. Agric. Sci.*, **2016**, *4*, 116-120. <https://www.cabdirect.org/cabdirect/abstract/20163102379>
- ³²⁷Hussain, Z., Amresh, G., Singh, S., Rao, C. V., Antidiarrheal and antiulcer activity of *Amaranthus spinosus* in experimental animals, *Pharmaceut. Biol.*, **2009**, *47*, 932-939. DOI:10.1080/13880200902950769
- ³²⁸Panda, S. K., Sarkar, G., Acharjya, M., Panda, P. K., Antiulcer activity of *Amaranthus spinosus* leaf extract and its comparison with famotidine in shay rats, *J. Drug Deliv. Therapeut.*, **2017**, *7*, 96-98. DOI:10.22270/jddt.v7i2.1392
- ³²⁹Chaudhary, M. A., Imran, I., Bashir, S., Mehmood, M. H., Rehman, N., Gilani, A. H., Evaluation of gut modulatory and bronchodilator activities of *Amaranthus spinosus* Linn, *BMC Complement. Alternat. Med.*, **2012**, *12*, Article 166, 11 pages. DOI:10.1186/1472-6882-12-166
- ³³⁰Amuthan, A., Chogtu, B., Bairy, K. L., Sudhakar, M., Prakash, M., Evaluation of diuretic activity of *Amaranthus spinosus* Linn. aqueous extract in Wistar rats, *J. Ethnopharmacol.*, **2012**, *140*, 424-427. DOI: 10.1016/j.jep.2012.01.049
- ³³¹Potlappalli, S., Narumalla, J., Naga, T. P., Govindadas, D., Chikkannasetty, S. S., Study of diuretic activity of aqueous extract of *Amaranthus spinosus* linn on rats, *Int. J. Basic Clin. Pharmacol.*, **2017**, *6*, 141-144. DOI:10.18203/2319-2003.ijbcp20164769
- ³³²Zeashan, H., Amresh, G., Singh, S., Rao, C. V., Protective effect of *Amaranthus spinosus* against *d*-galactosamine/lipopolysaccharide-induced hepatic failure, *Pharmaceut. Biol.*, **2010**, *48*, 1157-1163. DOI:10.3109/13880200903168023

- ³³³Rjeibi, I., Ben Saad, A., Hfaiedh, N., Oxidative damage and hepatotoxicity associated with deltamethrin in rats: The protective effects of *Amaranthus spinosus* seed extract, *Biomed. Pharmacother.*, **2016**, *84*, 853-860. DOI:10.1016/j.biopha.2016.10.010
- ³³⁴Ashok Kumar, B. S., Lakshman, K., Jayaveera, K. N., Vel Murgan, C., Arun Kumar, P. A., Vinod Kumar, R., Hegade, M., Sridhar, S. M., Pain management in mice using methanol extracts of three plants belongs to family *Amaranthaceae*, *Asian Pac. J. Trop. Med.*, **2010**, *3*, 527-530. DOI: 10.1016/S1995-7645(10)60127-7
- ³³⁵Sarker, U., Oba, S., Nutraceuticals, antioxidant pigments and phytochemicals in the leaves of *Amaranthus spinosus* and *Amaranthus viridis* weedy species, *Sci. Rep.*, **2019**, *9*, Article 20413, 10 pages. DOI:10.1038/s41598-019-50977-5
- ³³⁶Susantiningih, T., Ridwan, R., Prijanti, A. R., Sadikin, M., Freisleben, H. J., Schizonticidal effect of a combination of *Amaranthus spinosus* L. and *Andrographis paniculata* Burm. f./Nees extracts in *Plasmodium berghei*-infected mice, *Med. J. Indones.*, **2012**, *21*, 66-70.
<https://mji.ui.ac.id/journal/index.php/mji/article/view/482>
- ³³⁷Das, R. K., Gogoi, N., Babu, P. J., Sharma, P., Mahanta, C., Bora, U., The Synthesis of Gold Nanoparticles Using *Amaranthus spinosus* Leaf Extract and Study of Their Optical Properties, *Adv. Mater. Phys. Chem.*, **2012**, *2*, 275-281. <http://dx.doi.org/10.4236/ampc.2012.24040>
- ³³⁸Muthukumar, H., Matheswaran, M., *Amaranthus spinosus* Leaf Extract Mediated FeO Nanoparticles: Physicochemical Traits, Photocatalytic and Antioxidant Activity, *ACS Sustain. Chem. Eng.*, **2015**, *3*, 3149-3156.
DOI:10.1021/acssuschemeng.5b00722
- ³³⁹Adewolu, M., Adamson, A. A., *Amaranthus spinosus* Leaf Meal as Potential Dietary Protein Source in the Practical Diets for *Clarias gariepinus* (Burchell, 1822) Fingerlings, *Int. J. Zool. Res.*, **2011**, *7*, 128-137. DOI: 10.3923/ijzr.2011.128.137
- ³⁴⁰Kuswaningrum, O., Suwandono, A., Ariyanti, I., Hadisaputro, S., Suhartono, S., The Impact of Consuming *Amaranthus spinosus* L Extract on Prolactin Level and Breast Milk Production in Postpartum Mothers, *Belitung Nurs. J.*, **2017**, *3*, 541-547. DOI: 10.33546/bnj.194
- ³⁴¹Peixoto, P. V., Brust, L. A., Brito, M. F., Franca, T. N., da Cunha, B. R., de Andrade, G. B., *Amaranthus spinosus* (Amaranthaceae) poisoning in sheep in southern Brazil, *Pesq. Vet. Bras.*, **2003**, *23*, 179-184. (Article in Portuguese). <https://doi.org/10.1590/S0100-736X2003000400007>
- ³⁴²Murugan, K., Vanithakumari, G., Sampathraj, R., Effects of combined extracts of *Dolichos biflorus* seeds and *Amaranthus spinosus* roots on the accessory sex organs of male rats, *Ann. Sci. Life*, **1993**, *12*, 351-357. <https://www.ncbi.nlm.nih.gov/PMC3336559>
- ³⁴³Gigani, Y., Vekaria, A., Ali, S. A., Effect of *Abrus precatorius* and *Amaranthus spinosus* combination treatment on fertility in male rats, *J. Pharmacol. Pharmacotherapeut.*, **2012**, *3*, 272-273. DOI:10.4103/0976-500X.99441
- ³⁴⁴Gurumani, M., Balamurugan, K., Evaluation of antifertility potential of ethanolic extract of whole plant of *Amaranthus spinosus* in female albino rats, *World J. Pharm. Pharmaceut. Sci.*, **2014**, *3*, 2162-2167.
<https://www.researchgate.net/publication/287742204>
- ³⁴⁵Haq, A., Malik, A., Khan, A., Shah, M. R., Muhammad, P., Spinose, new coumaroyl flavone glycoside from *Amaranthus spinosus*, *Arch. Pharm. Res.*, **2004**, *27*, 1216-1219. DOI: 10.1007/BF02975884
- ³⁴⁶Stintzing, F. C., Kammerer, D., Schieber, A., Adama, H., Nacoulma, O. G., Carle, R., Betacyanins and phenolic compounds from *Amaranthus spinosus* L. And *Boerhavia erecta* L., *Z. Naturforsch. C.*, **2004**, *59*, 1-8. DOI:10.1515/znc-2004-1-201
- ³⁴⁷Queddeng, M. Q., Phytochemical Constituents of Kalunay Leaves and Roots (*Amaranthus spinosus* L.), *UNP Res. J.*, **2009**, *18*, 37-48. <https://ejournals.ph/article.php?id=6866>
- ³⁴⁸Balakrishnan, S., Natarajan, B., Arumugam, B., Hariram, B. V., Pharmacognostical Evaluation of *Amaranthus spinosus* L., *Pharmacog. J.*, **2011**, *3*, 13-18. DOI: 10.5530/pj.2011.19.3
- ³⁴⁹Jhade, D., Ahirwar, D., Jain, R., Sharma, N. K., Gupta, S., Pharmacognostic standardization, physico- and phytochemical evaluation of *Amaranthus spinosus* Linn. root, *J. Young Pharm.*, **2011**, *3*, 221-225. DOI:10.4103/0975-1483.83770
- ³⁵⁰Mishra, S. B., Verma, A., Mukerjee, A., Vijayakumar, M., Pharmacognostic Standardization and Phytochemical screening of Leaves of *Amaranthus spinosus* L., *Pharmacog. J.*, **2011**, *3*, 34-38. DOI:10.5530/pj.2011.26.6
- ³⁵¹Baral, M., Datta, A., Chakraborty, S., Chakraborty, P., Pharmacognostic studies on stem and leaves of *Amaranthus spinosus* Linn., *Int. J. Appl. Biol. Pharmaceut. Technol.*, **2011**, *2*, 41-47.
<https://www.researchgate.net/publication/236667897>
- ³⁵²Choudhury, A., Evaluation of Physicochemical and Phytochemical Parameters of *Amaranthus spinosus* Leaves, *Int. Res. J. Pharm.*, **2012**, *3*, 210-211.
<https://www.researchgate.net/publication/341713257>
- ³⁵³Sable, K. V., Saswade, R. R., Preliminary phytochemical analysis of *Amaranthus spinosus* leaves, *Int. J. Life Sci.*, **2017**, *5*, 742-745. <http://oaji.net/articles/2017/736-1515177213.pdf>
- ³⁵⁴Tuyen, P. N., Duong, N. T., Lien, D. T., Insights into chemical constituents of *Amaranthus spinosus* L. (Amaranthaceae), *Vietnam J. Chem.*, **2019**, *57*, 245-249. DOI: 10.1002/vjch.201900023
- ³⁵⁵Billah, A. H., Hussain, M. M., Dastagir, M. G., Ismail, M., Quader, A., Isolation of α -spinasterol from *Amaranthus spinosus* stems, *Bol. Latinoam. Caribe Plantas Med. Aromat.*, **2013**, *12*, 15-17.
<https://www.researchgate.net/publication/282504952>
- ³⁵⁶Asha, S., Rekha, R., Sadiq, A. M., Identification of Quercetin Present in Ethanol Leaf Extract of *Amaranthus spinosus* by Chromatographic Techniques, *Eur. J. Biomed. Pharmaceut. Sci.*, **2016**, *3*, 534-539.
https://www.ejbps.com/ejbps/abstract_id/1375
- ³⁵⁷Ang, A. M., Raman, I. C., Characterization of Mucilages from *Abelmoschus manihot* Linn., *Amaranthus spinosus* Linn. and *Talinum triangulare* (Jacq.) Willd. Leaves for Pharmaceutical Excipient Application, *Asian J. Biol. Life Sci.*, **2019**, *8*, 16-24. DOI:10.5530/ajbls.2019.8.3
- ³⁵⁸Sarwar, S., Sabir, M., Raza, S. S., Malik, S. A., Analysis of antimicrobial activity of medicinal plant *Amaranthus viridis*, *Int. J. Innov. Sci. Res.*, **2016**, *20*, 494-499. <http://www.ijisr.issr-journals.org/abstract.php?article=IJISR-15-296-02>
- ³⁵⁹Devi, D. S., Hindumathi, A., Pranathi, S., Gundeti, R., Shekar, S. C., GC-MS Analysis of Phytochemical Constituents and Screening for Antibacterial Activity of the Methanol Leaf Extract of *Amaranthus Viridis* Linn. Against Human Pathogenic Bacteria, *Indian Am. J. Pharmaceut. Res.*, **2017**, *7*, 7773-7779.
<https://zenodo.org/record/2383673#.Xy7gaSgzblU>
- ³⁶⁰Serag, M. M., El-Naga, A. Z., El-Gendey, M. A., Environmental impact of some poisonous weeds in Damietta, Egypt, *East Afr. Sch. J. of Agric. Life Sci.*, **2019**, *2*, 490-495. <https://www.researchgate.net/publication/338175684>
- ³⁶¹Iqbal, M. J., Hanif, S., Mahmood, Z., Anwar, F., Jamil, A., Antioxidant and antimicrobial activities of Chowlai (*Amaranthus viridis* L.) leaf and seed extracts, *J. Med. Plants Res.*, **2012**, *6*, 4450-4455. DOI: 10.5897/JMPR12.822

- ³⁶²Ahmed, S. A., Hanif, S., Iftkhar, T., Phytochemical Profiling with Antioxidant and Antimicrobial Screening of *Amaranthus viridis* L. Leaf and Seed Extracts, *Open J. Med. Microbiol.*, **2013**, *3*, 164-171. DOI: 10.4236/ojmm.2013.33025
- ³⁶³Akbar, M., Sherazi, I. N., Iqbal, M. S., Khalil, T., Waqas, H. M., Antibacterial and Antioxidant Activities of Slender Amaranth Weed, *Planta Daninha*, **2020**, *38*, Article e020192974, 8 pages. DOI: 10.1590/S0100-83582020380100006
- ³⁶⁴Carminate, B., Martin, G. B., Barcelos, R. M., Gontijo, I., de Almeida, M. S., Belinelo, V. J., Evaluation of Antifungal Activity of *Amaranthus viridis* L. (Amaranthaceae) on Fusariosis by *Pipernigrum* L. and on Anthracnose by *Musa* sp., *Agric. J.*, **2012**, *7*, 215-219. DOI:10.3923/aj.2012.215.219
- ³⁶⁵Kumar, B. S., Lakshman, K., Jayaveera, K. K., Shekar, D. S., Muragan, C. S., Manoj, B., Antinociceptive and Antipyretic Activities of *Amaranthus Viridis* Linn. in Different Experimental Models, *Avicenna J. Med. Biotechnol.*, **2009**, *1*, 167-171. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3558137/>
- ³⁶⁶Sahoo, H. B., Sahoo, S. K., Mishra, K., Sagar, R., Evaluation of the wound-healing potential of *Amaranthus viridis* (Linn.) in experimentally induced diabetic rats. *Int. J. Nutr. Pharmacol. Neurol. Dis.*, **2015**, *5*, 50-55. DOI:10.4103/2231-0738.153792
- ³⁶⁷Rahman, F., Afroz, S., Jahan, S., Hosain, M., Khondoker, D. F., Rahman, M., Banik, J., Ahmad, I., Rahmatullah, M., Antihyperglycemic and antinociceptive properties of methanolic extract of whole plants of *Amaranthus viridis* L. (Amaranthaceae). *Adv. Nat. Appl. Sci.*, **2012**, *6*, 1330-1335. <https://www.researchgate.net/publication/288820719>
- ³⁶⁸Kaur, N., Dhuna, V., Kamboj, S. S., Agrewala, J. N., Singh, J., A novel antiproliferative and antifungal lectin from *Amaranthus viridis* Linn seeds, *Protein Pept. Lett.*, **2006**, *13*, 897-905. DOI: 10.2174/092986606778256153
- ³⁶⁹Larbie, C., Appiah-Opong, R., Acheampong, F., Tuffour, I., Uto, T., Torkornoo, D., Marfo, E., Ankamah-Mensah, D., Opoku-Mensah, E., Abotsi, P., Anti-proliferative effect of *Amaranthus Viridis* Linn. on human leukemic cell lines- a preliminary study, *Int. J. Biol. Pharmaceut. Res.*, **2015**, *6*, 236-243. <http://129.122.16.11/bitstream/123456789/10961/1/2015%20IJBP.pdf>
- ³⁷⁰Shihab, U., Mahmud, I., Mynul, H., Amrita, B. Begum, R., *Amaranthus viridis* modulates anti-hyperglycemic pathways in hemi-diaphragm and improves glycogenesis liver function in rats, *J. Pharmacog. Phytother.*, **2016**, *8*, 173-181. DOI: 10.5897/JPP2016.0406
- ³⁷¹Ashok Kumar, B. S., Lakshman, K., Jayaveera, K. N., Sheshadri Shekar, D., Khan, S., Thippeswamy, B. S., Veerapur, V. P., Antidiabetic, antihyperlipidemic and antioxidant activities of methanolic extract of *Amaranthus viridis* Linn in alloxan induced diabetic rats, *Exper. Toxicol. Pathol.*, **2012**, *64*, 75-79. DOI:10.1016/j.etp.2010.06.009
- ³⁷²Girija, K., Lakshman, K., Pruthvi, N., Chandrika, P. U., Antihyperglycemic and hypolipidemic activity of methanolic extract of *Amaranthus viridis* leaves in experimental diabetes, *Indian J. Pharmacol.*, **2011**, *43*, 450-454. DOI: 10.4103/0253-7613.83120
- ³⁷³Sharma, N., Gupta P. C., Rao, C. V., Nutrient Content, Mineral Content and Antioxidant Activity of *Amaranthus viridis* and *Moringa oleifera* Leaves, *Res. J. Med. Plants*, **2012**, *6*, 253-259. DOI: 10.3923/rjmp.2012.253.259
- ³⁷⁴Jadhav, V., Biradar, S. D., Evaluation of Antioxidant Activity of *Amaranthus viridis* L. Methanolic Extract, *Int. J. Pharm. Biol. Sci.*, **2016**, *6*, 150-153. https://www.ijpbs.com/ijpbsadmin/upload/ijpbs_58157deb7619f.pdf
- ³⁷⁵Faiz, N., Priya, V. V., Ponnulakshmi, R., Gayathri, R., Shyamaladevi, B., Madhan, K., Manikannan, M., Selvaraj, J., *In vitro* antioxidant potential of stem of *Amaranthus viridis* - A medicine used in the Ayurvedic system of medicine, *Drug Invent. Today*, **2019**, *11*, 1148-1151. http://jprsolutions.info/article_detail.php?article_id=3477
- ³⁷⁶Sasikumar, V., Subramaniam, A., Aneesh, A., Saravanan, G., Protective Effect of Alkaloids from *Amaranthus Viridis* Linn. against Hydrogen Peroxide Induced Oxidative Damage in Human Erythrocytes (RBC), *Int. J. Clin. Endocrinol. Metab.*, **2015**, *1*, 49-53. <https://www.peertechz.com/articles/IJCEM-1-111.php>
- ³⁷⁷Kumari, S., Elancheran, R., Devi, R., Phytochemical screening, antioxidant, antityrosinase and antigenotoxic potential of *Amaranthus viridis* extract, *Indian J. Pharmacol.*, **2018**, *50*, 130-138. DOI:10.4103/ijp.IJP_77_18
- ³⁷⁸Salvamani, S., Gunasekaran, B., Shukor, M. Y., Shaharuddin, N. A., Sabullah, M. K., Ahmad, S. A., Anti-HMG-CoA Reductase, Antioxidant, and Anti-Inflammatory Activities of *Amaranthus viridis* Leaf Extract as a Potential Treatment for Hypercholesterolemia, *Evid. Based Complement. Alternat. Med.*, **2016**, Article 8090841, 10 pages. <http://dx.doi.org/10.1155/2016/8090841>
- ³⁷⁹Salvamani, S., Gunasekaran, B., Shukor, M. Y., Bakar, Z. A., Ahmad, S. A., Phytochemical investigation, hypocholesterolemic and anti-atherosclerotic effects of *Amaranthus viridis* leaf extract in hypercholesterolemia-induced rabbits, *RSC Adv.*, **2016**, *6*, 32685-32696. <https://doi.org/10.1039/c6ra04827g>
- ³⁸⁰Saravanan, G., Ponnurugan, P., *Amaranthus viridis* Linn. extract ameliorates isoproterenol-induced cardiac toxicity in rats by stabilizing circulatory antioxidant system, *Oxid. Antioxid. Med. Sci.*, **2012**, *1*, 69-73. DOI:10.5455/oams.060312.or.002
- ³⁸¹Saravanan, G., Ponnurugan, P., Sathiyavathi, M., Vadivukkarasi, S., Sengottuvelu, S., Cardioprotective activity of *Amaranthus viridis* Linn.: Effect on serum marker enzymes, cardiac troponin and antioxidant system in experimental myocardial infarcted rats, *Int. J. Cardiol.*, **2013**, *165*, 494-498. DOI:10.1016/j.ijcard.2011.09.005
- ³⁸²Asha, S., Thirunavukkarasu, P., Antiuro lithiatic Activity of *Amaranthus Viridis* on Ethylene Glycol Induced Male Rats, *Inventi Rap. Ethnopharmacol.*, **2013**, *4*, 1-4. <https://www.researchgate.net/publication/264977853>
- ³⁸³Ashok Kumar, B. S., Lakshman, K., Narayan Swamy, V. B., Arun Kumar, P. A., Sheshadri Shekar, D., Manoj, B., Vishwantha, G. L., Hepatoprotective and Antioxidant Activities of *Amaranthus viridis* Linn., *Maced. J. Med. Sci.*, **2011**, *4*, 125-130. DOI:10.3889/MJMS.1857-5773.2011.0163
- ³⁸⁴Ayoka, O. A., Ojo, O. E., Imafidon, E. C., Ademoye, K. A., Oladele, A. A., Neuro-endocrine effects of aqueous extract of *Amaranthus viridis* (Linn.) leaf in male Wistar rat model of cyclophosphamide-induced reproductive toxicity, *Toxicol. Rep.*, **2016**, *3*, 608-619. <http://dx.doi.org/10.1016/j.toxrep.2016.07.007>
- ³⁸⁵Olarewaju, O. A., Alashi, A. M., Aluko, R. E., Trypsin inhibitory activity of aqueous extracts from *Amaranthus viridis*, *Solanum macrocarpon*, and *Telfairia occidentalis* vegetable leaves, *Acta Hort.*, **2019**, *1238*, 129-136. DOI: 10.17660/ActaHortic.2019.1238.14
- ³⁸⁶Dayani Siriwardena, S. A., Karunathilaka, L. P., Karunathilaka, Y. A., Kodithuwakkam N. D., Comparative clinical study on the efficacy of *Amaranthus viridis* Linn (Tanduliyaka) and *Symplocos racemosa* Roxb (Lodhra) on Asrugdara with special reference to Menorrhagia, *Ayurv.*, **2009**, *30*, 421-424. <http://www.ayujournal.org/article.asp?issn=09748520;year=2009;volume=30;issue=4;spage=421>

- ³⁸⁷Kanerkar, U. R., Bhogaonkar, P. Y., Indurwade, N. H., Abortifacient Effect of *Amaranthus viridis* L. Aqueous Root Extract on Albino Rats, *Asian J. Biomed. Pharmaceut. Sci.*, **2016**, *6*, 13-16.
<https://www.alliedacademies.org/articles/abortifacient-effect-of-amaranthus-viridis-l-aqueous-root-extract-on-albino-rats.pdf>
- ³⁸⁸Affy, M. E., Tovi, W. M., Zougrou, N. E., Kouakou, K., Effect of methanolic extract of *Amaranthus viridis* leaves on reproductive functions in Wistar female rats, *J. Drug Deliv. Therapeut.*, **2019**, *9*, 119-126. DOI:10.22270/jddt.v9i6-s.3775
- ³⁸⁹Mani, D., Patel, N. K., Kumar, S., Kumar, A., Effect of FYM on the uptake of cadmium by Amaranth (*Amaranthus viridis* L.), *J. Earth Sci. Environ. Stud.*, **2015**, *1*, 1-5. DOI: 10.25177/JESES.1.1.1
- ³⁹⁰Joseph, J., Reddy, J., Sayantan, D., Effect of nickel uptake on selected growth parameters of *Amaranthus viridis* L., *J. Appl. Nat. Sci.*, **2018**, *10*, 1011-1017. DOI:10.31018/jans.v10i3.1838
- ³⁹¹Pal, S., Comparative Assessment of Cadmium, Chromium Resistance Capacity of *Amaranthus viridis*, *Trianthema portulacastrum*, *Alternanthera philoxeroides* Collected from Contaminated Site and Assessment of Phytoremedial Potential, *Environ. Ecol.*, **2018**, *36*, 409-417. <https://drive.google.com/file/d/1i6eGISi9Q0Ap6i9FZFEB2GhuQ46qCCJz/view>
- ³⁹²Radhika, B., Romali, D., Shraboni, R., Ragni, D., Sarojini, D., Priyanka, B., Arup Kumar, M., Risk of Heavy Metal Poisoning via *Amaranthus viridis* Growing in Potentially Contaminated Site, *Eur. J. Biomed. Pharmaceut. Sci.*, **2019**, *6*, 490-499. https://www.ejbps.com/ejbps/abstract_id/5488
- ³⁹³Phanjom, P., Borthakur, M., Das, R., Dey, S., Bhuyan, T., Green Synthesis of Silver Nanoparticles using Leaf Extract of *Amaranthus Viridis*, *Int. J. Nanotechnol. Appl.*, **2012**, *6*, 55-59. <https://www.researchgate.net/publication/273762430>
- ³⁹⁴Koyyati, R., Nagati, V. B., Nalvothula, R., Merugu, R., Kudle, K. R., Marx, P., Padigya, P. R., Antibacterial activity of silver nanoparticles synthesized using *Amaranthus viridis* twig extract, *Int. J. Res. Pharm. Sci.*, **2014**, *5*, 32-39. <https://pharmascope.org/ijrps/article/view/1156/1090>
- ³⁹⁵Fasuan, T. O., Gbadamosi, S. O., Akanbi, C. T., Modification of amaranth (*Amaranthus viridis*) starch, identification of functional groups, and its potentials as fat replacer, *J. Food Biochem.*, **2018**, *42*, Article e12537, 11 pages.
<https://doi.org/10.1111/jfbc.12537>
- ³⁹⁶Fasuan, T. O., Akanbi, C. T., Betiku, E., Acetylation of *Amaranthus viridis* starch: Modeling and process parameters optimization, *Food Sci. Nutr.*, **2018**, *6*, 1287-1297. <https://doi.org/10.1002/fsn3.677>
- ³⁹⁷Emmanuel, A. M., Roger, K. K., Toussaint, D. G., Koffi, K., Acute and subacute toxicity of the aqueous extract of *Amaranthus viridis* (Amaranthaceae) leaves in rats, *J. Phytopharmacol.*, **2018**, *7*, 366-372. http://www.phytopharmajournal.com/Vol7_Issue4_03.pdf
- ³⁹⁸Ragasa, C. Y., Austria, J. P., Subosa, A. F., Torres, O. B., Shen, C. C., Chemical Constituents of *Amaranthus viridis*, *Chem. Nat. Compd.*, **2015**, *51*, 146-147. DOI: 10.1007/s10600-015-1224-9
- ³⁹⁹Srivastava, R., Rurum, A. M., Mishra, P., Shehu, I. A., Rajak, C., Preliminary phytochemical investigation and TLC fingerprint profile of *Amaranthus* herbs with nutraceutical potential, *Pharm. Innov. J.*, **2018**, *7*, 224-229. <http://www.thepharmajournal.com/archives/?year=2018&vol=7&issue=5&ArticleId=2043>
- ⁴⁰⁰Roy, S., Dutta, A. K., Chakraborty, D. P., Amasterol, an ecdysone precursor and a growth inhibitor from *Amaranthus viridis*, *Phytochemistry*, **1982**, *21*, 2417-2420. [https://doi.org/10.1016/0031-9422\(82\)85224-2](https://doi.org/10.1016/0031-9422(82)85224-2)
- ⁴⁰¹Suraweera, W., Warrell, D., Whitaker, R., Menon, G., Rodrigues, R., Fu, S. H., Begum, R., Sati, P., Piyasena, K., Bhatia, M., Brown, P., Jha, P., Trends in snakebite deaths in India from 2000 to 2019 in a nationally representative mortality study, *eLife*, **2020**, *9*, Article e54076, 37 pages. DOI:10.7554/eLife.54076
- ⁴⁰²Upasani, S. V., Beldar, V. G., Tatiya, A. U., Upasani, M. S., Surana, S. J., Patil, D. S., Ethnomedicinal plants used for snakebite in India: a brief overview, *Integr. Med. Res.*, **2017**, *6*, 114-130. DOI: 10.1016/j.imr.2017.03.001
- ⁴⁰³Laddha, K. S., Ghosh, D., Isolation Of 20 -Hydroxyecdysone From Indian Medicinal Plant *Achyranthes Aspera* And Development of Simple HPLC Analysis, *Nat. Prod. Ind. J.*, **2005**, *1*, 1-4. <https://www.tsijournals.com/abstract/526.html>
- ⁴⁰⁴Saraf, A., Samant, A., Pharmacognostic and Phytochemical Studies for the Establishment of Quality Parameters of Leaf of *Achyranthes aspera* Linn., *Int. J. Pharmacog. Phytochem. Res.*, **2014**, *6*, 122-127. <https://www.researchgate.net/publication/284253916>
- ⁴⁰⁵Salama, F., El-Ghani, M. A., El-Tayeh, N., Amro, A., El-Naggar, S., Adaptive Responses of *Aerva javanica* to Severe Aridity in the Egyptian Deserts, *Egypt. J. Bot.*, **2018**, *58*, 171-184. DOI: 10.21608/ejbo.2018.2557.1144
- ⁴⁰⁶Chabane, D., Mouhoub, F., Establishment of vitroplants from female flowers of *Aerva javanica* (Burm. F) Juss ex Schult. *Acta Hort.*, **2017**, *1155*, 607-612. DOI: 10.17660/ActaHortic.2017.1155.88
- ⁴⁰⁷Ikram, N., Dawar, S., Soil amendment with *Aerva javanica* (Burm. F.) Juss. ex Schult. in the control of root rot fungi of cowpea [*Vigna unguiculata* (L.) Walp.] and mung bean [*Vigna radiata* (L.)], *Acta Agrobot.*, **2012**, *65*, 69-74. DOI: 10.5586/aa.2012.009
- ⁴⁰⁸Singh, A., Kandasamy, T., Odhav, B., *In vitro* propagation of *Alternanthera sessilis* (sessilejooyweed), a famine food plant, *Afr. J. Biotechnol.*, **2009**, *8*, 5691-5695. DOI: 10.5897/AJB09.307
- ⁴⁰⁹Gayathri, B. M., Balasuriya, K., Gunawardena, G. S., Rajapakse, R. P., Dharmaratne, H. R., Toxicological studies of the water extract of green leafy vegetable Sessilejooy weed (*Alternanthera sessilis*), *Curr. Sci.*, **2006**, *91*, 1517-1520. <https://www.researchgate.net/publication/283720405>
- ⁴¹⁰Belkhode, S., Panhekar, D., Kalambe, A., Seasonal Variation of Heavy Metal Concentration in *Alternanthera Sessilis*, *Indian J. Appl. Res.*, **2016**, *6*, 397-399. DOI: 10.36106/ijar
- ⁴¹¹Noori, M., Talebi, M., Nasiri, Z., Seven *Amaranthus* L. (Amaranthaceae) Taxa Flavonoid Compounds from Tehran Province, Iran, *Int. J. Mod. Bot.*, **2015**, *5*, 9-17. DOI:10.5923/j.ijmb.20150501.02
- ⁴¹²Suma, S., Ambika, S. R., Kazinczi, G., Narwal, S. S., Allelopathic plants. 6. *Amaranthus spp.*, *Allelopathy J.*, **2002**, *10*, 1-12. <https://www.researchgate.net/publication/286992211>
- ⁴¹³Srivastava, R., Nutritional quality of some cultivated and wild species of *Amaranthus* L., *Int. J. Pharmaceut. Sci. Res.*, **2011**, *2*, 3152-3156. <https://www.researchgate.net/publication/267960456>
- ⁴¹⁴Marin, D. I., Bolohan, C., Mihalache, M., Rusu, T., Research on *Amaranthus cruentus* L. and *Amaranthus hypochondriacus* L. Grown in South-Eastern Romania (Mora Domneasca – Ilfov), *Sci. Pap. Ser. A*, **2011**, *54*, 297-303. <https://www.researchgate.net/publication/308784667>
- ⁴¹⁵Kornarzynski, K., Dziwulska-Hunek, A., Kornarzynska-Gregorowicz, A., Sujak, A., Effect of Electromagnetic Stimulation of Amaranth Seeds of Different Initial Moisture on the Germination Parameters and Photosynthetic Pigments Content, *Sci. Rep.*, **2018**, *8*, Article14023, 12 pages. DOI:10.1038/s41598-018-32305-5
- ⁴¹⁶Hlinkova, A., Bednarova, A., Havrlentova, M., Supova, J., Cicova, I., Evaluation of fatty acid composition among selected amaranth grains grown in two consecutive years,

- Biologia*, **2013**, 68, 641-650. DOI: 10.2478/s11756-013-0190-6
- ⁴¹⁷Goncalves Netto, A., Nicolai, M., Carvalho, S. J., Borgato, E. A., Christoffoleti, P. J., Multiple resistance of *amaranthus palmeri* to ALS and EPSPS inhibiting herbicides in the state of Mato Grosso, Brazil, *Planta Daninha*, **2016**, 34, 581-587. DOI:10.1590/S0100-83582016340300019
- ⁴¹⁸Woo, M. L., Chemical Constituents of *Amaranthus retroflexus*, *Bot. Gaz.*, **1919**, 68, 313-344. DOI: 10.1086/332569
- ⁴¹⁹Francischini, A. C., Constantin, J., Oliveira, R. S., Santos, G., Franchini, L. H., Biffe, D. F., Resistance of *Amaranthus retroflexus* to acetolactate synthase inhibitor herbicides in Brazil, *Planta Daninha*, **2014**, 32, 437-446. DOI:10.1590/S0100-83582014000200022
- ⁴²⁰Wang, H., Guo, W., Zhang, L., Zhao, K., Ge, L., Lv, X., Liu, W., Wang, J., Multiple resistance to thifensulfuronmethyl and fomesafen in redroot pigweed (*Amaranthus retroflexus* L.) from China, *Chil. J. Agric. Res.*, **2017**, 77, 311-317. DOI:10.4067/S0718-58392017000400311
- ⁴²¹Dinu, M., Anghel, A. I., Oлару, O. T., Șeremet, O., Calalb, T., Cojocaru-Toma, M., Negreș, S., Hovanet, V., Zbarcea, C. E., Ancuceanu, R., Toxicity investigation of an extract of *Amaranthus retroflexus* L. (*Amaranthaceae*) leaves, *Farmacia*, **2017**, 2, 289-294.
<https://www.researchgate.net/publication/316958026>
- ⁴²²Qi, Y., Yan, B., Fu, G., Guan, X., Du, L., Li, J., Germination of Seeds and Seedling Growth of *Amaranthus retroflexus* L. following Sublethal Exposure of Parent Plants to Herbicides, *Sci. Rep.*, **2017**, 7, Article 157, 8 pages. DOI:10.1038/s41598-017-00153-4
- ⁴²³Kepeczynski, J., Sznigir, P., Response of *Amaranthus retroflexus* L. seeds to gibberellic acid, ethylene and abscisic acid depending on duration of stratification and burial, *Plant Growth Regul.*, **2013**, 70, 15–26. DOI: 10.1007/s10725-012-9774-3
- ⁴²⁴Vyn, J. D., Swanton, C. J., Weaver, S. E., Sikkema, P. H., Control of *Amaranthus tuberculatus* var. *rudis* (common water hemp) with pre- and post-emergence herbicides in *Zea mays* L. (maize), *Crop. Protection*, **2006**, 25, 1051-1056. <https://doi.org/10.1016/j.cropro.2006.01.016>
- ⁴²⁵Syahri, R., Widaryanto, E., Wicaksono, K. P., Bioactive compound from mangoes leaves extract as potential soil bioherbicide to control amaranth weed (*Amaranthus spinosus* Linn.), *J. Degrade. Min. Land Manage.*, **2017**, 4, 829-836. DOI:10.15243/jdmlm.2017.043.829
- ⁴²⁶Sanon, A., Beguiristain, T., Cebon, A., Berthelin, J., Ndoye, I., Leyval, C., Sylla, S., Duponnois, R., Changes in soil diversity and global activities following invasions of the exotic invasive plant, *Amaranthus viridis* L., decrease the growth of native Sahelian *Acacia* species, *FEMS Microbiol. Ecol.*, **2009**, 70, 118–131. DOI:10.1111/j.1574-6941.2009.00740.x
- ⁴²⁷Francischini, A. C., Constantin, J., Oliveira, R. S., Santos, G., Braz, G. B., Dan, H. A., First report of *Amaranthus viridis* resistance to herbicides, *Planta Daninha*, **2014**, 32, 571-578. DOI: 10.1590/S0100-83582014000300013
- ⁴²⁸Moyin-Jesu, E. I., Use of selected agro-industrial biomass for enhancing seed nitrogen, ash and crude protein quality of *Amaranthus viridis* L., *Emir. J. Food Agric.*, **2007**, 19, 13-21. <https://www.ejfa.me/index.php/journal/article/view/406>
- ⁴²⁹Ratnasooriya, W. D., Obeysekera, M., Jayakody, J., Ratnasooriya, C., Weerasekera, K. R., Lack of diuretic activity of hot water extract of whole plant of *Amaranthus viridis* L. in rats, *J. Pharm. Negat. Results*, **2012**, 3, 26-28. DOI:10.4103/0976-9234.99646
- ⁴³⁰Piyasena, K. G., Wickramarachchi, W. A., Kumar, N. S., Jayasinghe, L., Fujimoto, Y., Two phytotoxic azaphilone derivatives from *Chaetomium lobosum*, a fungal endophyte isolated from *Amaranthus viridis* leaves, *Mycology*, **2015**, 6, 158-160. DOI: 10.1080/21501203.2015.1089332

Personal notice from the author

I grew up in an agricultural family, and most local *Amaranthaceae* plants are very common in our fields and gardens. We have always known their allelopathic effects, especially that of *Amaranthus retroflexus* (Table 8). We unrooted them when they grew close to crops, but we highly esteemed them as food for our livestock. When I started the literature search for this article, a young plant of *A. retroflexus* started growing in our garden. So, I planted very young plant of Chile pepper (*Capsicum annum*, *Solanaceae*) near it. Both plants grew very well, the plant of *A. retroflexus* started blossoming, but the pepper plant, despite reaching impressive size, was not flowering. I added flowering promoter to both plants, and *A. retroflexus* flowered more rapidly. The pepper plants were very green but no flowers or fruits. When *A. retroflexus* was about 100 cm high, I unrooted it. Within a week, the pepper plants were flowering so impressively and now (second week of August), they are heavily loaded with fruits.

Received: 15.08.2020.

Accepted: 31.10.2020.