



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)


NOTE

THE OAT-LIKE GRASS *TRISETOPSIS ASPERA* (MUNRO EX THWAITES) RÖSER & A.WÖLK (POACEAE): A NEW RECORD FOR THE FLORA OF CENTRAL WESTERN GHATS OF KARNATAKA, INDIA

H.U. Abhijit, Y.L. Krishnamurthy & K. Gopalakrishna Bhat

26 January 2021 | Vol. 13 | No. 1 | Pages: 17601–17603

DOI: [10.11609/jott.6181.13.1.17601-17603](https://doi.org/10.11609/jott.6181.13.1.17601-17603)



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact ravi@threatenedtaxa.org

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Member



Publisher & Host





The oat-like grass *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk (Poaceae): a new record for the flora of central Western Ghats of Karnataka, India

H.U. Abhijit¹ , Y.L. Krishnamurthy² & K. Gopalakrishna Bhat³

^{1,2} Department of P.G. Studies and research in applied Botany, Kuvempu University, Jnanasahyadri, Shankaraghatta, Karnataka 577451, India.

³ Taxonomy Research Centre, Department of Botany, Poorna Prajna College, Volakadu Road, Udupi, Karnataka 576101, India.
¹abhicrogon@gmail.com, ²murthy_ylk@yahoo.co.in (corresponding author), ³kakunje_bhat@yahoo.co.in

During the survey of grasses of Baba Budangiri Hills, Western Ghats of Chikkamagaluru District, Karnataka (13.431°N & 75.758°E), some interesting grass specimens were collected from the montane highlands associated with *Chrysopogon zeylanicus* (Steud.) Thwaites, *Arundinella pumila* (Hochst.) Steud. and *Agrostris pilosula* Trin. Initially, these were identified as *Helictotrichon aspera* by referring to Flora of the Presidency of Madras (Fischer 1934–36) and The Grasses of Burma, Ceylon, India and Pakistan (Bor 1960). The identity of this grass was later confirmed by matching our sample with photograph of the type specimen. A scrutiny of literature revealed that this species has now been transferred to the genus *Trisetopsis* by M. Röser and A. Wölk (Wölk & Röser 2013) as the morphological and phylogenetic studies by them revealed that *Helictotrichon* s.l. is polyphyletic and heterogeneous (Wölk & Röser 2017). The genus *Trisetopsis* is characterized by its apical, bifid lemma. This species, *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk, was hitherto known only from Kerala and Tamil Nadu (Sreekumar & Nair 1991; Kabeer & Nair 2009). Sreekumar & Nair (1991) reported this

species as *H. virescens* (Nees ex Steud.) Henrard. They followed Henrard (1940) and Sevenstert & Veldkamp (1983) to treat *H. aspera* as a synonym of *H. virescens*; however, in this work by following Kellogg et al. (2020), it is considered as *Trisetopsis aspera* and reported here as an addition to the grass flora of Karnataka. A brief description along with photographs is provided to facilitate easy recognition of this grass. The herbarium specimens are deposited in Herbarium of Department of Applied Botany, Kuvempu University, Shankaraghatta, Shivamogga, Karnataka.

Trisetopsis aspera (Munro ex Thwaites) Röser & A. Wölk in Taxon 66(1): 38. 2017 *Avena aspera* Munro ex Thwaites, Enum. Pl. Zeyl. 372. 1864. *Helictotrichon asperum* (Munro ex Thwaites) Bor in Indian Forest Rec., Bot. n.s., 1: 68. 1938; Bor, Grasses Burma, Ceylon, India & Pakistan: 438. 1960. *Avenastrum asperum* (Munro ex Thwaites) Vierh. in Verh. Ges. Deutsch. Naturf. 85(2;1): 672. 1914; Fischer in Gamble, Fl. Madras: 1802. 1934. *Helictotrichon virescens* sensu Sreek. & V.J. Nair Fl. Kerala: Grasses: 351. 1991,

Editor: V. Sampath Kumar, Botanical Survey of India, Coimbatore, India.

Date of publication: 26 January 2021 (online & print)

Citation: Abhijit, H.U., Y.L. Krishnamurthy & K.G. Bhat (2021). The oat-like grass *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk (Poaceae): a new record for the flora of central Western Ghats of Karnataka, India. *Journal of Threatened Taxa* 13(1): 17601–17603. <https://doi.org/10.11609/jott.6181.13.1.17601-17603>

Copyright: © Abhijit et al. 2021. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: DST- Inspire (IF170707).

Competing interests: The authors declare no competing interests.

Acknowledgements: The authors wish to thank the authorities of Karnataka Forest Department for giving the necessary permission to collect the specimens from the forest. The first author acknowledges the Department of Science and Technology (DST), Inspire for financial assistance and also Mr. Vishwajith H U, Mr. Pavan kumar K S, Mr. Krishna Karanth K M, Mr. Bharath B T and Mr. Srinivasa Murthy M V for their support during the research work.





Image 1. *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk. A—Habitat | B—habit | C—raceme | D—leaf | E—spikelet | F—lower & upper glume | G—lemma & palea | H—stamens | I—dissected spikelet. © H.U. Abhijit.




Image 2. Herbarium of *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk.

p.p. quoad syn. *H. asperum*, non (Nees ex Steud.) Henrard.

Tufted perennials; culms up to 120cm high. Leaf blades up to 40 × 0.5 cm, glabrous or pubescent; sheaths glabrous or pubescent; ligule membranous, up to 4mm long. Panicles up to 30cm long, effuse, nodding. Spikelets 10–14 mm long (excluding awns); florets 3–5, uppermost floret rudimentary and sometimes reduced to awn. Glumes unequal, lanceolate-oblong, herbaceous, 3-nerved, acute to acuminate; lower 5–7 mm long;

upper 8–10 mm long. Lemmas lanceolate, scabrous, 7–9-nerved, lowest 8.5–10.5 mm long, bidentate at tip and awned from back near middle; awn geniculate, 10–13 mm long, scabrous. Palea 6–6.8 mm long, ciliate on keels. Stamens 3; anthers 2.5–2.8 mm long. Styles 2; stigmas plumose. Caryopsis linear-elliptic, 3.5–3.8 mm long, pubescent. (Image 1B- I)

Flowering and fruiting: September–December

Habitat and Ecology: Grasslands of high altitude about 1,200m (Image 1A).

Distribution: India: Kerala, Tamil Nadu and in the present work from Karnataka. Endemic.

Specimens examined: KUAB455, 03.i.2019, India, Karnataka, Chikkamagaluru District, Baba Budangiri Hills, 13.431°E & 75.758°N, coll. H.U. Abhijit.

References

- Bor, N.L. (1960). *The Grasses of Burma, Ceylon, India and Pakistan (excluding Bambusae)*. Pergamon Press, Oxford. 767pp.
- Fischer, C.E.C. (1934). Gramineae, pp. 1689–1864. In: Gamble, J.S. (ed.). *The Flora of the Presidency of Madras*. Part X. Adlard & Son, London.
- Fischer, C.E.C. (1934). Gramineae, pp. 1687–1864. In: Gamble, J.S. (ed.). *The Flora of the Presidency Madras*. Part X & XI. Vol. 3 (2nd Edition 1967). Botanical survey of India, Calcutta, 1802pp.
- Henrard, J.T. (1940). Notes on the nomenclature of some grasses. *Blumea* 3: 411–480.
- Kabeer, K.A. & V.J. Nair (2009). *Flora of Tamil Nadu - Grasses*. Botanical Survey of India, 525pp.
- Kellogg, E.A., J.A. Richard, K.S. Bawa, K.N. Gandhi, B.R. Kailash, K.N. Ganeshiah, U.B. Shrestha & P. Raven (2020). Checklist of the grasses of India. *PhytoKeys* 163: 1–560.
- Sevenstert, J.G. & J.F. Veldkamp (1983). A revision of *Helictotrichon* (Gramineae) in Malaysia. *Blumea* 28: 329–342.
- Sreekumar, P.V. & V.J. Nair (1991). *Flora of Kerala - Grasses*. Botanical Survey of India. 470pp.
- Wölk, A. & M. Röser (2013). The new genus *Trisetopsis* and new combinations in oat-like grasses (Poaceae). *Schlechtendalia* 25: 57–61.
- Wölk, A. & M. Röser (2017). Hybridization and long-distance colonization in oat-like grasses of South and East Asia, including an amended circumscription of *Helictotrichon* and the description of the new genus *Tzveleviochloa* (Poaceae). *Taxon* 66(1): 20–43.





www.threatenedtaxa.org

PLATINUM
OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

January 2021 | Vol. 13 | No. 1 | Pages: 17455–17610
Date of Publication: 26 January 2021 (Online & Print)
DOI: 10.11609/jott.2021.13.1.17455-17610

Communications

Diversity and distribution of snakes in Trashigang Territorial Forest Division, eastern Bhutan

– Bal Krishna Koirala, Karma Jamtsho, Phuntsho Wangdi, Dawa Tshering, Rinchen Wangdi, Lam Norbu, Sonam Phuntsho, Sonam Lhendup & Tshering Nidup, Pp. 17455–17469

Freshwater fishes of Cauvery Wildlife Sanctuary, Western Ghats of Karnataka, India

– Naren Sreenivasan, Neethi Mahesh & Rajeev Raghavan, Pp. 17470–17476

Fish communities and associated habitat variables in the upper Subansiri River of Arunachal Pradesh, eastern Himalaya, India

– Sutanu Satpathy, Kuppusamy Sivakumar & Jeyaraj Antony Johnson, Pp. 17477–17486

Diversity and distribution of odonates in Rani Reserve Forest, Assam, India

– Dipti Thakuria & Jatin Kalita, Pp. 17487–17503

An assessment of the population status of the threatened medicinal plant *Illicium griffithii* Hook.f. & Thomson in West Kameng District of Arunachal Pradesh, India

– Tashi Dorjee Bapu & Gibji Nimasow, Pp. 17504–17512

Short Communications

The discovery of a melanistic Leopard *Panthera pardus delacouri* (Linnaeus, 1758) (Mammalia: Carnivora: Felidae) at Bukit Kudung in Jeli, Kelantan, Peninsular Malaysia: conservation and ecotourism

– Kamarul Hambali, Nor Fakhira Muhamad Fazli, Aainaa Amir, Norashikin Fauzi, Nor Hizami Hassin, Muhamad Azahar Abas, Muhammad Firdaus Abdul Karim & Ai Yin Sow, Pp. 17513–17516

On the epidemiology of helminth parasites in Hangul Deer *Cervus hanglu hanglu* (Mammalia: Artiodactyla: Cervidae) of Dachigam National Park, India

– Naziya Khurshid, Hidayatulla Tak, Ruqeyya Nazir, Kulsum Ahmad Bhat & Muniza Manzoor, Pp. 17517–17520

Histopathological findings of infections caused by canine distemper virus, *Trypanosoma cruzi*, and other parasites in two free-ranging White-nosed Coatis *Nasua narica* (Carnivora: Procyonidae) from Costa Rica

– Jorge Rojas-Jiménez, Juan A. Morales-Acuña, Milena Argüello-Sáenz, Sílvia E. Acevedo-González, Michael J. Yabsley & Andrea Urbina-Villalobos, Pp. 17521–17528

On a new species of *Macrobrachium* Spence Bate (Decapoda: Palaemonidae) from Ayeyarwady River, Myanmar

– H.H.S. Myo, K.V. Jayachandran & K.L. Khin, Pp. 17529–17536

Review of the tiger beetle genus *Calomera* Motschulsky, 1862 (Coleoptera: Cicindelidae) of the Philippines

– Milton Norman Medina, Alexander Anichtchenko & Jürgen Wiesner, Pp. 17537–17542

Rediscovery of Martin's Duskhawker *Anaciaeschna martini* (Selys, 1897) (Odonata: Aeshnidae) from Western Ghats, peninsular India, with notes on its current distribution and oviposition behavior

– Kalesh Sadasivan, Manoj Sethumadavan, S. Jeevith & Baiju Kochunarayanan, Pp. 17543–17547

A note on the current distribution of reedtail damselfly *Protosticta rufostigma* Kimmins, 1958 (Odonata: Zygoptera: Platystictidae) from Western Ghats, and its addition to the odonate checklist of Kerala

– Kalesh Sadasivan & Muhamed Jafer Palot, Pp. 17548–17553

Assessment of threat status of the holly fern *Cyrtomium micropterum* (Kunze) Ching (Polypodiopsida: Dryopteridaceae) in India using IUCN Regional guidelines

– C. Bagathsingh & A. Benniamin, Pp. 17554–17560

Notes

First report of the Asiatic Brush-tailed Porcupine *Atherurus macrourus* (Linnaeus, 1758) (Mammalia: Rodentia: Hystricidae) from West Bengal, India

– Suraj Kumar Dash, Abhisek Chettri, Dipanjan Naha & Sambandam Sathyakumar, Pp. 17561–17563

Record of the world's biggest pangolin? New observations of bodyweight and total body length of the Indian Pangolin *Manis crassicaudata* Gray, 1827 (Mammalia: Pholidota: Manidae) from Mannar District, Sri Lanka

– Priyan Perera, Hirusha Randimal Algewatta & Buddhika Vidanage, Pp. 17564–17568

First record of *Touit melanonotus* (Wied, 1820) (Aves: Psittaciformes: Psittacidae) in Cantareira State Park, Brazil: new colonization or simply unnoticed?

– Marcos Antônio Melo & David de Almeida Braga, Pp. 17569–17573

Is *Bombus pomorum* (Panzer, 1805) (Hymenoptera: Apidae) a new bumblebee for Siberia or an indigenous species?

– Alexandr Byvaltsev, Svyatoslav Knyazev & Anatoly Afinogenov, Pp. 17574–17579

Some new records of scarab beetles of the genus *Onthophagus* Latreille, 1802 (Coleoptera: Scarabaeidae) from northern Western Ghats, Maharashtra, with a checklist

– Aparna Sureshchandra Kalawate, Banani Mukhopadhyay, Sonal Vithal Pawar & Vighnesh Durgaram Shinde, Pp. 17580–17586

Ecological importance of two large heritage trees in Moyar River valley, southern India

– Vedagiri Thirumurugan, Nehru Prabakaran, Vishnu Sreedharan Nair & Chinnasamy Ramesh, Pp. 17587–17591

Bulbophyllum spathulatum (Orchidaceae), a new record for Bhutan

– Pema Zangpo, Phub Gyeltshen & Pankaj Kumar, Pp. 17592–17596

On the occurrence and distribution of the narrowly endemic Andaman Lantern Flower *Ceropegia andamanica* (Apocynaceae: Ceropegieae)

– M. Uma Maheshwari & K. Karthigeyan, Pp. 17597–17600

The oat-like grass *Trisetopsis aspera* (Munro ex Thwaites) Röser & A.Wölk (Poaceae): a new record for the flora of central Western Ghats of Karnataka, India

– H.U. Abhijit, Y.L. Krishnamurthy & K. Gopalakrishna Bhat, Pp. 17601–17603

Star Grass Lily *Iphigenia stellata* Blatter (Colchicaceae) – a new addition to the flora of Gujarat, India

– Mitesh B. Patel, Pp. 17604–17606

A new record of pyrenocarpous lichen to the Indian biota

– N. Rajaprabu, P. Ponmurugan & Gaurav K. Mishra, Pp. 17607–17610

Member



Publisher & Host

