

# STARDOM™

Jamesbrittenia



BRAND  
NEW 23



**STARDOM™ White Fusion**

- Enhanced resistance to foliage disease
- Day neutral / benefits from long days
- Heat tolerant and continues flowering through the summer
- Perfect for combinations
- Mature height and width 8" x 12" (20cm x 30cm)

More varieties in the STARDOM™ series:



**STARDOM™ Pink**



**STARDOM™ Purple**



**STARDOM™ Goldstar**

For additional technical growing information:

STARDOM™ [CLICK HERE](#)



# STARDOM™

Jamesbrittenia



## PROPAGATION TIPS

- Stick on priority – Number 1 out of 4 categories



Flowering time  
Spring / Summer  
Flowering



Width  
(6 - 8")  
15 - 20 cm



Height  
(10 - 14")  
25 - 36 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21°-22° C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65°-68° F (18°-20° C)		Weeks 3-4 100 ppm N	Day of sticking & Week 2



**Irrigation specification** - Remove from mist as soon as possible. Mist enough to maintain turgid cuttings – keep on dry side.



Apply broad spectrum fungicide the day of sticking to control leaf disease. Stardom may require Daminozide in propagation @ 1500 -2000 ppm if grown warm and under low light.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.5	0.80	72°-74° F (22°-23° C)	50 ppm N in mist			Day of sticking	
Week 2	5.8 to 6.5	.80 to .90	72°-74° F (22°-23° C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.5	.90 to 1.0	Cool to 68°-70° F (20°-21° C)	Feed 100 ppm				
Week 4	5.8 to 6.5	.90 to 1.0	65°-68° F (18°-20° C)	Feed 100 ppm		Daminozide as needed		

For additional technical growing information:

STARDOM™ [CLICK HERE](#)



# STARDOM™

Jamesbrittenia



## FINISHING TIPS

- Maintain consistent soil moisture –plants should not dry to severe wilt
- Cool temperature finishing for best quality and optimum flower color

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 65°-68° F (18°-20° C)	Pinch – Recommended	100 - 200 ppm N	Daminozide at 1500 to 2500 ppm on more vigorous varieties
7 to 8 wks (15 cm) 6 ½"	1 ppp	Not Required – Day Neutral	Soil EC 0.8 - 1.0 pH 5.8 to 6.5	Paclobutrazol Drench at 1 - 2 ppm to finish
7 to 8 wks (20 cm) 8"	3 ppp			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide at 1500 to 2500 ppm on young finished plants - Paclobutrazol drench application at 1 to 2 ppm for finishing if required for the more vigorous varieties
- Maintain good airflow and allow plants to dry before nightfall
- Scout for Aphids & Thrips on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.8 to 6.5	0.8 to 1.0	65°-68° F (18°-20° C)	feed 100 to 150 ppm	Not required	Daminozide as needed 1500 to 2500 ppm	Spray fungicide after transplant	
<b>Week 6 to 12</b>	5.8 to 6.5	0.8 to 1.0	65°-68° F (18°-20° C)	feed 100 to 200 ppm	Not required	Paclobutrazol Drench at 1 - 2 ppm		Paclobutrazol 1 – 2 ppm drench when plants reach 85% of desired size

For additional technical growing information:

STARDOM™ [CLICK HERE](#)

